

ISSN 0019-5588


# Indian Journal of Pure & Applied Mathematics

---

DEVOTED PRIMARILY TO ORIGINAL RESEARCH  
IN PURE AND APPLIED MATHEMATICS

Including  
CONTENTS & INDEX

VOLUME 20/12  
DECEMBER 1989



# INDIAN JOURNAL OF PURE AND APPLIED MATHEMATICS

*Published monthly by the*

INDIAN NATIONAL SCIENCE ACADEMY

Editor of Publications

PROFESSOR D. V. S. JAIN

Department of Physical Chemistry, Panjab University

Chandigarh 160 014

PROFESSOR J. K. GHOSH

Indian Statistical Institute

203, Barrackpore Trunk Road

Calcutta 700 035

PROFESSOR A. S. GUPTA

Department of Mathematics

Indian Institute of Technology

Kharagpur 721 302

PROFESSOR M. K. JAIN

Department of Mathematics

Indian Institute of Technology

Hauz Khas

New Delhi 110 016

PROFESSOR S. K. JOSHI

Director

National Physical Laboratory

New Delhi 110 012

PROFESSOR V. KANNAN

Dean, School of Mathematics &

Copmputer/Information Sciences

University of Hyderabad

P O Central University

Hyderabad 500 134

Assistant Executive Secretary

(Associate Editor/Publications)

DR. M. DHARA

Subscriptions :

For India, Pakistan, Sri Lanka, Nepal, Bangladesh and Burma, Contact :

Associate Editor, Indian National Science Academy, Bahadur Shah Zafar Marg,  
New Delhi 110002, Telephone : 3311865, Telex : 31-61835 INSA IN.

For other countries, Contact :

M/s J. C. Baltzer AG, Scientific Publishing Company, Wettsteinplatz 10, CH-4058 Basel,  
Switzerland, Telephone : 61-268925, Telex : 63475.

*The Journal is indexed in the Science Citation Index; Current Contents (Physical, Chemical & Earth Sciences); Mathematical Reviews; INSPEC Science Abstracts (Part A); as well as all the major abstracting services of the World.*

PROFESSOR N. MUKUNDA

Centre for Theoretical Studies

Indian Institute of Science

Bangalore 560 012

DR PREM NARAIN

Director

Indian Agricultural Statistics

Research Institute, Library Avenue

New Delhi 110 012

PROFESSOR I. B. S. PASSI

Centre for advanced study in Mathematics

Panjab University

Chandigarh 160 014

PROFESSOR PHOOLAN PRASAD

Department of Applied Mathematics

Indian Institute of Science

Bangalore 560 012

PROFESSOR M. S. RAGHUNATHAN

Senior Professor of Mathematics

Tata Institute of Fundamental Research

Homi Bhabha Road

Bombay 500 005

PROFESSOR T. N. SHOREY

School of Mathematics

Tata Institute of Fundamental Research

Homi Bhabha Road

Bombay 400 005

Assistant Editor

SRI R. D. BHALLA



# PSEUDOLINEARITY AND EFFICIENCY VIA DINI DERIVATIVES

SHASHI AGGARWAL

*Department of Mathematics, Miranda House, University of Delhi, Delhi 110007*

AND

DAVINDER BHATIA

*Department of Mathematics, S.G.T.B. Khalsa College, University of Delhi, Delhi 110007*

(Received 17 May 1988; after revision 15 February 1989; accepted 27 July 1989)

In this paper, characterization of pseudolinear functions in terms of Dini derivatives is given. Necessary and sufficient conditions for efficiency in terms of Dini derivatives are derived for multiobjective programming problems involving pseudolinear functions.

## INTRODUCTION

Nonlinear multiobjective programming problems involving pseudolinear functions have been studied by Chew and Choo<sup>1</sup>. They have characterized pseudolinear functions by means of proportional functions under the assumption of differentiability. Rockafellar<sup>5</sup> has pointed out that the functions involved may not be always differentiable and so characterization of pseudolinear functions without assuming differentiability is required.

Kaul *et al.*<sup>4</sup> have defined semilocally pseudolinear functions in terms of right derivatives generalizing pseudolinear functions and have extended results of Chew and Choo<sup>1</sup> concerning efficiency for multiobjective programming problems involving functions that are semilocally pseudolinear. It was shown by Kaul *et al.*<sup>4</sup> that a function  $f$  is semilocally pseudolinear on a set  $\Gamma \subseteq R^n$  iff  $\exists$  a real valued function  $p$  called proportional function of  $f$ , defined on  $\Gamma \times \Gamma$  such that  $p(x, y) > 0$  and  $f(y) = f(x) + p(x, y) df^+(x, y-x)$  for all  $x, y$  in  $\Gamma$ .

The purpose of this paper is to define pseudolinear functions in terms of Dini derivatives<sup>2,3</sup> which generalize the class of semilocally pseudolinear functions. In this paper, we consider the multiobjective pseudolinear programming problem of the form

$$(P) \quad \text{Max } f(x) = (f_1(x), \dots, f_k(x))$$

subject to

$$x \in X = \{x \in S \mid g_j(x) \geq 0, j = 1, \dots, m\}$$

where it is assumed that  $f_i, i = 1, \dots, k$  and  $g_j, j = 1, \dots, m$  are real pseudolinear functions defined on a convex subset  $S$  of  $R^N$ .

Here an alternative characterization of pseudolinearity is provided which makes use of Dini Derivatives instead of right derivatives. The useful feature about Dini derivatives is that they always exists whereas right derivatives need not. Such a characterization seems necessary because of the existence of functions of the type given in the example presented in section 2, which motivated the authors to make a present study.

Theorems 2 and 3 develop necessary and sufficient conditions for the existence of an efficient solution for the above mentioned problem (P), whereas Kaul *et al.*<sup>4</sup> have derived only sufficient condition.

## 2. DINI DERIVATIVES AND PSEUDOLINEAR FUNCTIONS

**Definition 1**—Let  $f$  be a real valued function defined over  $S$ , a convex subset of  $R^N$ . Let  $x \in R^N$ ,  $v \in R^N$  with  $v^T v = 1$ . Dini derivatives of  $f$  in the direction  $v$  at  $x$  are defined as follows :

$$D_v^{+u} f(x) = \lim_{n \rightarrow \infty} \sup_{\{t_n\}} \left\{ \frac{f(x + t_n v) - f(x)}{t_n} : 0 < t_n \leq 1/n \right\}$$

$$D_v^{+l} f(x) = \lim_{n \rightarrow \infty} \inf_{\{t_n\}} \left\{ \frac{f(x + t_n v) - f(x)}{t_n} : 0 < t_n \leq 1/n \right\}$$

$$D_v^{-u} f(x) = \lim_{n \rightarrow \infty} \sup_{\{t_n\}} \left\{ \frac{f(x - t_n v) - f(x)}{-t_n} : 0 < t_n \leq 1/n \right\}$$

$$D_v^{-l} f(x) = \lim_{n \rightarrow \infty} \inf_{\{t_n\}} \left\{ \frac{f(x - t_n v) - f(x)}{-t_n} : 0 < t_n \leq 1/n \right\}.$$

Here  $D_v^{+u} f(x)$  is the upper right derivative,  $D_v^{+l} f(x)$  is the lower right derivative,  $D_v^{-u} f(x)$  is the upper left derivative and  $D_v^{-l} f(x)$  is lower left derivative evaluated at  $x$  in the direction  $v$ . Limits can be infinite in the above definition. It may easily be proved, by using the definitions, that

(I) Dini derivatives always exist (finite or infinite) for any function  $f$  and satisfy

$$D_v^{+u} f(x) \geq D_v^{+l} f(x), D_v^{-u} f(x) \geq D_v^{-l} f(x)$$

$$D_v^{+u} (-f)(x) = - D_v^{+l} f(x), D_v^{+l} (-f)(x) = - D_v^{+u} f(x)$$

(II) If  $D_v^{+l} f(x) = D_v^{+u} f(x)$  (or if  $D_v^{-l} f(x) = D_v^{-u} f(x)$ ) then the common value, written  $D_v^+ f(x)$  (or  $D_v^- f(x)$ ) is just the right (or left) derivative of  $f$  at  $x$  in the



direction  $v$ . In general,  $D_v^- f(x) \leq D_v^+ f(x)$  and if  $D_v^- f(x) = D_v^+ f(x)$ , then  $f$  has the derivative at  $x$  in the direction  $v$  and the common value will be denoted by  $D_v f(x)$ .

If  $f$  is a function of one variable, then usually, we take the directional vector  $v$  to be the scalar 1 and we write

$$D_1^{+u} f(x) \equiv D^{+u} f(x), \quad D_1^{+l} f(x) \equiv D^{+l} f(x)$$

$$D_1^+ f(x) \equiv D^+ f(x), \quad D_1^- f(x) \equiv D^- f(x).$$

Diewert<sup>2</sup> introduces pseudoconcave functions as :

*Definition 2*—A function  $f$  defined over a convex subset  $S$  of  $R^N$  is said to be pseudoconcave over  $S$  iff for every  $x^0 \in S$ ,  $v \in R^N$  satisfying

$$v^T v = 1, \quad t > 0, \quad x^0 + t v \in S, \quad D_v^{+u} f(x^0) \leq 0$$

implies

$$f(x^0 + t v) \leq f(x^0).$$

In the light of the above definition 2, we may now define pseudoconvex functions as follows :

*Definition 3*—A function  $f$  defined over a convex subset  $S$  of  $R^N$  is said to be pseudoconvex over  $S$  iff  $-f$  is pseudoconcave over  $S$  i. e. for every  $x^0 \in S$ ,  $v \in R^N$  satisfying

$$v^T v = 1, \quad t > 0, \quad x^0 + t v \in S, \quad D_v^{+l} f(x^0) \geq 0 \text{ implies } f(x^0 + t v) \geq f(x^0).$$

Hence a pseudolinear function in terms of Dini derivatives may be defined as follows

*Definition 4*—A function  $f$  defined over a convex subset  $S$  of  $R^N$  is said to be pseudolinear over  $S$  iff it is both pseudoconcave and pseudoconvex according to definitions 2 and 3 above.

*Example*—Let

$$f(x) = \begin{cases} 1 & , -2 \leq x \leq -1 \\ 0 & , -1 < x \leq 0 \\ 1/2^{n+1} & , 1/2^{n+1} \leq x < 1/2^n, n = 0, 1, 2, \dots \end{cases}$$

be a function defined over  $[-2, 1[$ .

Some of the Dini derivatives computed at different points of the domain of the function are as follows :

$$D^{+u} f(-1) = -\infty, D^{+l} f(-1) = -\infty$$

$$D^{+u} f(0) = 1, D^{+l} f(0) = 1/2.$$

It may be easily verified that

$$D^{+u} f(-1) < 0 \Rightarrow f(-1+t) < f(-1), t > 0, (-1+t) \in ]-1, 0]$$

$$D^{+l} f(0) > 0 \Rightarrow f(0+t) > f(0), t > 0, 0+t \in ]0, 1[.$$

Thus  $f$  is pseudolinear according to definition 4 over  $[-2, 1[$ . It is obvious that  $f$  does not have the right derivative at  $x = 0$  and hence  $f$  is not semilocally pseudolinear Kaul *et al.*<sup>4</sup> over  $[-2, 1]$ .

*Assumption*—In the sequel, we assume all the functions and Dini derivatives to be finite.

*Theorem 1*—Let  $f$  be a function defined over a convex subset  $S$  of  $R^N$ . Then the following statements are equivalent.

- (i)  $f$  is pseudolinear over  $S$ .
- (ii) There exist real functions  $p$  and  $q$  defined over  $S \times S$  such that  $p(x^\circ, x^\circ + tv) > 0$ ,  $q(x^\circ, x^\circ + tv) > 0$  and

$$\begin{aligned} f(x^\circ + tv) &= f(x^\circ) + p(x^\circ, x^\circ + tv) D_v^{+u} f(x^\circ) \\ &\quad + q(x^\circ, x^\circ + tv) D_v^{+l} f(x^\circ) \end{aligned} \quad \dots(1)$$

for any  $x^\circ \in S$ ,  $v \in R^N$  satisfying  $v^T v = 1$ ,  $t > 0$ ,  $x^\circ + tv \in S$ .

*PROOF* : (i)  $\Rightarrow$  (ii). Let  $x^\circ \in S$ ,  $v \in R^N$  satisfying  $v^T v = 1$ ,  $t > 0$ ,  $x^\circ + tv \in S$  and  $f$  be pseudolinear over  $S$ . Therefore

$$\left. \begin{aligned} &\text{if } D_v^{+u} f(x^\circ) \leq 0 \text{ then } f(x^\circ + tv) \leq f(x^\circ) \\ &\text{and if } D_v^{+l} f(x^\circ) \geq 0 \text{ then } f(x^\circ + tv) \geq f(x^\circ) \end{aligned} \right] \quad \dots(2)$$

As the right derivative of  $f$  may not exist at every point of  $S$ , we have two cases.

*Case 1*: When right derivative at  $x^\circ$  does not exist, i. e.

$$D_v^{+u} f(x^\circ) \neq D_v^{+l} f(x^\circ)$$

*Case 2*: When right derivative at  $x^\circ$  exists i. e.

$$D_v^{+u} f(x^\circ) = D_v^{+l} f(x^\circ) = D_v^{+} f(x^\circ).$$



Hence in case (1), relation (2) leads to

$$(a) \quad D_v^{+u} f(x^0) < 0 \Rightarrow D_v^{+l} f(x^0) < 0 \text{ and } f(x^0 + tv) < f(x^0)$$

$$(b) \quad D_v^{+u} f(x^0) = 0 \Rightarrow D_v^{+l} f(x^0) < 0 \text{ and } f(x^0 + tv) < f(x^0)$$

$$(c) \quad D_v^{+l} f(x^0) > 0 \Rightarrow D_v^{+u} f(x^0) > 0 \text{ and } f(x^0 + tv) > f(x^0)$$

$$(d) \quad D_v^{+l} f(x^0) = 0 \Rightarrow D_v^{+u} f(x^0) > 0 \text{ and } f(x^0 + tv) > f(x^0)$$

since  $D_v^{+l} f(x) \leq D_v^{+u} f(x)$  for all  $x \in S$  and moreover for any  $x \in S$ ,  $v \in R^N$  satisfying  $v^T v = 1$ ,  $t > 0$ ,  $x + tv \in S$  if  $f(x + tv) = f(x)$  then  $D_v^{+u} f(x) = D_v^{+l} f(x) = 0$ .

Now we establish statement (ii) in each of the above four possibilities. In possibilities (a) and (c) above we may define

$$p(x^0, x^0 + tv) = \frac{f(x^0 + tv) - f(x^0)}{D_v^{+u} f(x^0)}$$

and

$$q(x^0, x^0 + tv) = \frac{f(x^0 + tv) - f(x^0)}{D_v^{+l} f(x^0)}$$

and the results follows.

In case of possibility (b), we can define  $p(x^0, x^0 + tv)$  to be any positive real number and

$$q(x^0, x^0 + tv) = \frac{f(x^0 + tv) - f(x^0)}{D_v^{+l} f(x^0)}.$$

Similarly, in case of possibility (d), we define

$$p(x^0, x^0 + tv) = \frac{f(x^0 + tv) - f(x^0)}{D_v^{+u} f(x^0)}$$

and  $q(x^0, x^0 + tv)$  to be any positive real number. Thus in all the four possibilities, statement (ii) holds.

In case 2, relation (1) can be written as

$$f(x^0 + tv) = f(x^0) + p'(x^0, x^0 + tv) D_v^{+} f(x^0)$$

where  $p'(x^0, x^0 + tv) = p(x^0, x^0 + tv) + q(x^0, x^0 + tv)$  and can be defined as

$$p'(x^0, x^0 + tv) = \begin{cases} \frac{f(x^0 + tv) - f(x^0)}{D_v^+ f(x^0)} & \text{if } D_v^+ f(x^0) \neq 0 \\ K, \text{ a positive real number, if } D_v^+ f(x^0) = 0. \end{cases}$$

The positiveness of the function  $p'$  can be proved exactly in the same manner as proved for the functions  $p$  and  $q$  in Case 1.

(ii)  $\Rightarrow$  (i) : Suppose that there are functions  $p$  and  $q$  defined over  $S \times S$  such that  $p(x^0, x^0 + tv) > 0$ ,  $q(x^0, x^0 + tv) > 0$  and relation (1) holds.

Then  $D_v^{+u} f(x^0) \leq 0$  and the fact that  $D_v^{+l} f(x^0) \leq D_v^{+u} f(x^0)$  imply that  $f(x^0 + tv) \leq f(x^0)$  showing that  $f$  is pseudoconcave over  $S$ . Also  $D_v^{+l} f(x^0) \geq 0$  and again the fact that  $D_v^{+u} f(x^0) > D_v^{+l} f(x^0)$  imply that  $f(x^0 + tv) \geq f(x^0)$  showing that  $f$  is pseudoconvex over  $S$ .

Hence  $f$  is pseudolinear over  $S$ .

Q.E.D.

### 3. NECESSARY AND SUFFICIENT CONDITIONS FOR THE EXISTENCE OF EFFICIENT SOLUTION

Throughout this section  $f_i$ ,  $i = 1, \dots, k$ ,  $g_j$ ,  $j = 1, \dots, m$  will be real pseudolinear functions according to definition 4 over a convex subset  $S$  of  $R^N$  with positive functions  $p_i$ ,  $q_i$ ,  $i = 1, \dots, k$  and  $p_j$ ,  $q'_j$ ,  $j = 1, \dots, m$  respectively.

Consider the following multiobjective pseudolinear programming problem :

$$(P) \quad \text{Max } f(x) = (f_1(x), \dots, f_k(x))$$

subject to

$$x \in X = \{x \in S \mid g_j(x) \geq 0, j = 1, 2, \dots, m\}.$$

For a point  $x$  in  $X$ , we shall denote by  $I(x)$ , the set of all  $j$  such that  $g_j(x) = 0$

**Definition 5**—A vector  $v \in R^N$  satisfying  $v^T v = 1$  is called a feasible direction for  $X$  at  $x^* \in X$  if there exists a  $\delta > 0$  such that  $x^* + tv \in X$  for all  $0 < t \leq \delta$ .

The set of all feasible directions at  $x^* \in X$  is denoted by  $D(x^*)$  and

$$D(x^*) = \{v \in R^N : v^T v = 1, \exists \delta > 0 \text{ such that}$$

$$x^* + tv \in X \text{ for } 0 < t \leq \delta\}.$$

**Definition 6**—A point  $x^* \in X$  is said to be efficient solution of problem (P) if there exists no  $v \in R^N$  with  $v^T v = 1$  such that

$$f_i(x^* + tv) \geq f_i(x^*), \quad i = 1, \dots, k$$



and  $f_t(x^* + tv) > f_t(x^*)$  for at least one  $i$ ,  
 where  $0 < t \leq \delta$  for some  $\delta > 0$  such that  $x^* + tv \in X$ .

Note 1 : For  $x^* \in X$ , denote

$$p_i^* = p_i(x^*, x^* + t^*v), \quad q_i^* = q_i(x^*, x^* + t^*v), \quad i = 1, \dots, k$$

$$p_j'^* = p_j'(x^*, x^* + t^*v), \quad q_j'^* = q_j'(x^*, x^* + t^*v), \quad j = 1, \dots, m$$

where we can define<sup>2</sup>

$$v = \frac{(x - x^*)}{[(x - x^*)^T (x - x^*)]^{1/2}}, \quad x = x^* + t^*v \in X \text{ for}$$

$$t^* = [(x - x^*)^T (x - x^*)]^{1/2} > 0.$$

Lemma—Let  $x^*$  be a feasible solution for problem (P), then

$$p_j'^* D_v^{+u} g_j(x^*) + q_j'^* D_v^{+l} g_j(x^*) > 0 \text{ for } j \in I(x^*)$$

and

$$p_j'^* D_v^{+u} g_j(x^*) + q_j'^* D_v^{+l} g_j(x^*) > -\infty, j \notin I(x^*) \text{ implies } v \in D(x^*).$$

PROOF : Let  $j \in I(x^*)$  and  $p_j'^* D_v^{+u} g_j(x^*) + q_j'^* D_v^{+l} g_j(x^*) > 0$  for some direction  $v$ . Then  $g_j(x^*) = 0$  since  $j \in I(x^*)$ . Suppose that there exists  $\delta_j > 0$  such that  $0 < t_j \leq \delta_j$  and  $g_j(x^* + t_j v) < 0$ . Then

$$\begin{aligned} & p_j'^* D_v^{+u} g_j(x^*) + q_j'^* D_v^{+l} g_j(x^*) \\ &= p_j'^* \limsup_{t \rightarrow 0} \frac{g_j(x^* + tv) - 0}{t} + q_j'^* \liminf_{t \rightarrow 0^+} \frac{g_j(x^* + tv) - 0}{t} \leq 0 \end{aligned}$$

which contradicts  $p_j'^* D_v^{+u} g_j(x^*) + q_j'^* D_v^{+l} g_j(x^*) > 0$ .

Thus there exists some  $\delta_j > 0$  such that

$$g_j(x^* + tv) \geq 0 \text{ for } 0 < t \leq \delta_j, j \in I(x^*). \quad \dots(3)$$

Now let  $j \notin I(x^*)$  and  $p_j'^* D_v^{+u} g_j(x^*) + q_j'^* D_v^{+l} g_j(x^*) > -\infty$  for some direction  $v$ . Since  $j \notin I(x^*)$ , therefore  $g_j(x^*) > 0$ . Suppose that there exists  $\delta_j > 0$  such that  $0 < t_j \leq \delta_j$  and  $g_j(x^* + t_j v) < 0$ . Then

$$p_j'^* D_v^{+u} g_j(x^*) + q_j'^* D_v^{+l} g_j(x^*) = p_j'^* \limsup_{t \rightarrow 0^+}$$

(equation continued on p. 1180)

$$+ \frac{g_j(x^* + tv) - g_j(x^*)}{t} + q_j'^* \lim_{t \rightarrow 0^+} \inf \frac{g_j(x^* + tv) - g_j(x^*)}{t} \rightarrow -\infty$$

which contradicts  $p_j'^* D_v^{+u} g_j(x^*) + q_j'^* D_v^{+l} g_j(x^*) > -\infty$ .

Thus there exists some  $\delta_j > 0$  such that

$$g_j(x^* + tv) \geq 0, 0 < t \leq \delta_j, j \in I(x^*) \quad \dots(4)$$

(3) and (4) imply that  $v \in D(x^*)$ .

Q.E.D.

**Theorem 2**—If  $x^*$  is an efficient solution of problem (P) and constraints are assumed to satisfy conditions of Lemma then there exists  $\lambda_i > 0, i = 1, \dots, k$  and  $\mu_j > 0, j \in I(x^*)$  such that

$$\sum_{i=1}^k \lambda_i [p_i^* D_v^{+u} f_i(x^*) + q_i^* D_v^{+l} f_i(x^*)] + \sum_{j \in I(x^*)} \mu_j [p_j'^* D_v^{+u} g_j(x^*) + q_j'^* D_v^{+l} g_j(x^*)] = 0. \quad \dots(5)$$

**PROOF :** Let  $x^*$  be an efficient solution of problem (P) and constraints satisfy the conditions of Lemma. Then by Lemma,  $v \in D(x^*)$  and thus there is a  $\delta > 0$  such that  $x^* + tv \in X$  for  $0 < t \leq \delta$ .

We now assert that for  $1 \leq r \leq k$  the system

$$(A) \quad \begin{cases} [p_j'^* D_v^{+u} g_j(x^*) v + q_j'^* D_v^{+l} g_j(x^*) v]^T (x^* + tv - x^*) \geq 0, \\ \quad \quad \quad j \in I(x^*) \quad \dots(6) \\ [p_i^* D_v^{+u} f_i(x^*) v + q_i^* D_v^{+l} f_i(x^*) v]^T (x^* + tv - x^*) \geq 0, \\ \quad \quad \quad i = 1, \dots, k, i \neq r \quad \dots(7) \\ [p_r^* D_v^{+u} f_r(x^*) v + q_r^* D_v^{+l} f_r(x^*) v]^T (x^* + tv - x^*) > 0 \end{cases}$$

...

has no solution  $x^* + tv \in S$  for  $0 < t \leq \delta$ .

Let, if possible,  $x^* + tv \in S$  for some  $t \in ]0, \delta]$  be a solution of the system A. Now as  $f_i$ 's are pseudolinear over  $S$  with positive functions  $p_i$ 's and  $q_i$ 's, therefore relation (1) yields

$$f_i(x^* + tv) - f_i(x^*) = p_i(x^*, x^* + tv) D_v^{+u} f_i(x^*) + q_i(x^*, x^* + tv) D_v^{+l} f_i(x^*) \quad i = 1, \dots, k \quad \dots(9)$$



and clearly  $x = x^* + t v \in X$ .

As in Note 1, we may now define

$$v = \frac{x - x^*}{[(x - x^*)^T (x - x^*)]^{1/2}},$$

$$x = x^* + t^* v \text{ for } t^* = [(x - x^*)^T (x - x^*)]^{1/2} > 0.$$

Hence (9) implies

$$\begin{aligned} f_i(x) - f_i(x^*) &= p_i^* D_v^{+u} f_i(x^*) + q_i^* D_v^{+l} f_i(x^*), i = 1, \dots, k \\ &\geq 0, i = 1, \dots, k, i \neq r \text{ (using (7))}. \end{aligned} \quad \dots(10)$$

$$\text{Similarly } f_r(x) - f_r(x^*) > 0 \text{ (using (8))}. \quad \dots(11)$$

(10) and (11) contradict that  $x^*$  is efficient for problem (P). Hence system  $A$  has no solution  $x^* + t v \in S$  for  $0 < t \leq \delta$ . Therefore, by Farkas' Lemma given by Mangasarian<sup>5</sup>, there exist  $\lambda_{r_i} \geq 0$ ,  $\mu_{r_j} \geq 0$ ,  $r = 1, \dots, k$  such that

$$\begin{aligned} &\sum_{j \in I(x^*)} \mu_{r_j} [p_j^{r*} D_v^{+u} g_j(x^*) v + q_j^{r*} D_v^{+l} g_j(x^*) v]^T \\ &\quad + \sum_{\substack{i=1 \\ i \neq r}}^k \lambda_{r_i} [p_i^* D_v^{+u} f_i(x^*) v + q_i^* D_v^{+l} f_i(x^*) v]^T \\ &\quad - [p_r^* D_v^{+u} f_r(x^*) v + q_r^* D_v^{+l} f_r(x^*) v]^T = 0. \end{aligned} \quad \dots(12)$$

Summing (12) over  $r = 1, 2, \dots, k$ , we get

$$\begin{aligned} &\sum_{i=1}^k \lambda_i [p_i^* D_v^{+u} f_i(x^*) v + q_i^* D_v^{+l} f_i(x^*) v]^T \\ &\quad \sum_{j \in I(x^*)} \mu_j [p_j^{r*} D_v^{+u} g_j(x^*) v + q_j^{r*} D_v^{+l} g_j(x^*) v]^T = 0 \end{aligned} \quad \dots(13)$$

where

$$\lambda_i = 1 + \sum_{r \neq i} \lambda_{r_i} > 0, \mu_j = \sum_{r=1}^k \mu_{r_j} \geq 0.$$

Relation (13) yields relation (5).

Q.E.D.

*Theorem 3*—Let  $x^*$  be a feasible solution of the problem (P) and let there exist  $\lambda_i > 0$ ,  $i = 1, \dots, k$ ,  $\mu_j \geq 0$ ,  $j \in I(x^*)$  such that relation (5) holds. Then  $x^*$  is efficient for problem (P).

PROOF: Let, if possible,  $x^*$  be not an efficient solution of problem (P). Then there exists some  $v \in K^N$  with  $v^T v = 1$  such that

$$\left. \begin{aligned} f_i(x^* + tv) &\geq f_i(x^*), \quad i = 1, \dots, k, \quad i \neq r \\ f_r(x^* + tv) &> f_r(x^*) \\ g_j(x^* + tv) &\geq 0, \quad j = 1, \dots, m \end{aligned} \right\} \quad \dots (14)$$

where  $0 < t \leq \delta$  for some  $\delta > 0$  such that  $x^* + tv \in S$ .

Further on using relation (1) and defining  $p_i^*, q_i^*, p_j'^*, q_j'^*$  as in Note 1, relation (14) gives

$$\sum_{i=1}^k \lambda_i (p_i^* D_v^{+u} f_i(x^*) + q_i^* D_v^{+l} f_i(x^*)) > 0$$

and

$$p_j'^* D_v^{+u} g_j(x^*) + q_j'^* D_v^{+l} g_j(x^*) \geq 0, \quad j \in I(x^*)$$

Hence

$$\begin{aligned} 0 &\leq \sum_{j \in I(x^*)} [p_j'^* D_v^{+u} g_j(x^*) + q_j'^* D_v^{+l} g_j(x^*)] \\ &= - \sum_{i=1}^k \lambda_i [p_i^* D_v^{+u} f_i(x^*) + q_i^* D_v^{+l} f_i(x^*)] \\ &< 0 \end{aligned}$$

which is an obvious contradiction.

Hence  $x^*$  is efficient for problem (P).

Q.E.D.

Note 2: More general results where in problem (P) the objective functions are pseudoconcave and constraints are quasiconcave using Dini derivatives are developed in authors subsequent paper.

#### ACKNOWLEDGEMENT

The authors wish to thank Professor R. N. Kaul, Department of Mathematics, University of Delhi, Delhi, India, for his constant encouragement in preparing this paper. Thanks are also due to the referees for their valuable suggestions.



## REFERENCES

1. K. L. Chew and E. U. Choo, *Math. Prog.* **28** (1984), 226-39.
2. S. Schaible and W. T. Ziembea, *Generalized Concavity in Optimization and Economics*. Academic Press, New York, 1981, pp. 51-93.
3. E. Hewitt and K. Stromberg, *Real and Abstract Analysis*. Springer-Verlag, New York, 1969, pp. 256-57.
4. R. N. Kaul, Vinod Lyall and Surjeet Kaur, *European J. Ops. Res.* **36** (1988), 402-409.
5. R. T. Rockafellar, *Convex Analysis*, Princeton University Press, Princeton, New Jersey, 1970.
6. O. L. Mangasarian, *Nonlinear Programming*, McGraw-Hill Book Co, Inc., New York, 1969; 16-17.

# ON THE EXISTENCE OF UNITY IN LEHMER'S $\psi$ -PRODUCT RING

V. SITARAMAIAH

Department of Mathematics, Pondicherry Engineering College  
Pillaichavady, Pondichery 605104

(Received 19 April 1988; accepted 17 July 1989)

Let  $T$  be a non-empty subset of  $Z^+ \times Z^+$  where  $Z^+$  is the set of positive integers and  $\psi : T \rightarrow Z^+$  be a mapping such that for each  $n \in Z^+$ ,  $\psi(x, y) = n$  has a finite number of solutions. If  $f$  and  $g$  are two arithmetic functions, then the binary operation  $\psi$  on the set of arithmetic function  $F$  is defined by

$$(f \psi g)(n) = \sum_{\psi(x, y) = n} f(x) g(y)$$

for any  $n \in Z^+$ . If  $(F, +, \psi)$  is a commutative ring, where  $+$  denotes the usual pointwise addition and  $\psi(x, y) \geq \max\{x, y\}$  for all  $(x, y) \in T$ , we prove that the ring  $(F, +, \psi)$  possesses the unity if and only if for each  $k \in Z^+$ ,  $\psi(x, k) = k$  has a solution. In such a case the unity can be explicitly determined.

## INTRODUCTION

An arithmetic function is a complex-valued function whose domain is the set of positive integers  $Z^+$ . Let  $F$  denote the set of all arithmetic functions. Let  $T$  be a non-empty subset of  $Z^+ \times Z^+$  and let  $\psi : T \rightarrow Z^+$  be a mapping satisfying the following postulates:

- (I) For each  $n \in Z^+$ ,  $\psi(x, y) = n$  has a finite number of solutions.
- (II) If  $(x, y) \in T$ , then  $(y, x) \in T$  and  $\psi(x, y) = \psi(y, x)$ .
- (III)  $(y, z) \in T$  and  $(x, \psi(y, z)) \in T$  if and only if " $(x, y) \in T$  and  $(\psi(x, y), z) \in T$ "; whenever one of these conditions holds, we have  $\psi(x, \psi(y, z)) = \psi(\psi(x, y), z)$ .
- (IV)  $\psi(1, 1) = 1$  and for each  $k \in Z^+$ ,  $\psi(x, k) = k$  has a solution.
- (V) For each  $k \in Z^+$ ,  $k = \max\{x \in Z^+ : \psi(x, y) = k \text{ for some } y \in Z^+\}$  or equivalently  $\psi(x, y) \geq \max\{x, y\}$  for all  $(x, y) \in T$ .

If we define the binary operation  $\psi$  on  $F$  by

$$(f \psi g)(n) = \sum_{\psi(x, y) = n} f(x) g(y)$$

for each  $n \in Z^+$ ,  $f, g \in F$ , then using the postulates (I), (II) and (III) it is easily seen that  $(F, +, \psi)$  is a commutative ring.

The following is proved in Sita Ramaiah<sup>1</sup>:

*Lemma 1.1* (Sita Ramaiah<sup>1</sup>, Lemma 2.1)—An arithmetic function  $g$  is an identity with respect to  $\psi$  if and only if, for any fixed  $k$ ,  $n \in Z^+$

$$\sum_{\psi(x,k)=n} g(x) = \begin{cases} 1, & \text{if } n = k \\ 0 & \text{if } n \neq k. \end{cases} \quad \dots(1.1)$$

Let

$$S_k = \{x : \psi(x, k) = k\} \quad \dots(1.2)$$

$$ik = \min S_k \quad \dots(1.3)$$

and

$$S = \{ik : k = 1, 2, \dots\}. \quad \dots(1.4)$$

The number  $ik$  in (1.3) exists by postulate (IV).

If the equation  $\psi(x, k) = n$  has a unique solution in  $S$  if  $n = k$  and no solution in  $S$  if  $n \neq k$ , it was mentioned in Theorem 2 of (Sita Ramaiah<sup>1</sup>) that the characteristic function  $X$  of  $S$  would be the unity of the commutative ring  $(F, +, \psi)$ .

It was proved in Sita Ramaiah<sup>2</sup> (Theorem 3.1) that if  $g$  is the unity of  $(F, +, \psi)$ , then  $g$  is integer-valued and  $g(ik) = 1$ , where  $ik$  is as given in (1.3). In addition, if  $g$  is non-negative, then  $g$  must be the characteristic function of the set  $S$  given in (1.4) (See Sita Ramaiah<sup>2</sup>, Theorem 3.2). Also, an example of a ring  $(F, +, \psi)$  was given (Sita Ramaiah<sup>2</sup>, Example 3.1) in which the unity assumed negative-values also.

Suppose  $g$  is the unity of  $(F, +, \psi)$ . We shall now investigate the question of determining  $g$  from the relation (1.1). From (1.1), we have

$$\sum_{x \in S_k} g(x) = 1$$

so that for  $k \in S_k$

$$g(k) = 1 - \sum_{\substack{x < k \\ x \in S_k}} g(x) \quad \dots(1.5)$$

where  $S_k$  is as given in (1.2). Let  $k \notin S_k$ . Let  $x_r$  be the largest element in  $S_k$  so that  $x_r < k$ . From (1.1), we obtain

$$\sum_{\psi(x, x_r)=k} g(x) = 0. \quad \dots(1.6)$$

Now,  $x_r \in S_k$  implies that  $\psi(k, x_r) = k$ . Hence from (1.6),

$$g(k) = - \sum_{\substack{\psi(x, x_r)=k \\ x < k}} g(x). \quad \dots(1.7)$$



If  $g(1) = 1$  and  $g(x)$  has been defined for  $1 \leq x < k$ , then the relations (1.5) and (1.7) completely determine the value of  $g(k)$  for any  $k \in \mathbb{Z}^+$ . The task is to prove the otherway. We show that (See Lemma 2.2) the relation (1.7) implies that  $g(k) = 0$  whenever  $k \notin S_k$ . Using this and (1.5) we ultimately show that (see Theorem 2.1)  $g$  is the unity of  $(F, +, \psi)$ , thus establishing the existence of the unity in the ring  $(F, +, \psi)$ , in the presence of the postulates (I) through (V).

## 2. MAIN RESULTS

First we prove the following :

*Lemma 2.1*—We have

- (i)  $a, b \in S_k \Rightarrow \psi(a, b) \in S_k$ .
- (ii) If  $a \in S_k$ , then  $S_a \subseteq S_k$ .
- (iii)  $\psi(a, b) = k$  implies that  $S_a \subseteq S_k$  and  $S_b \subseteq S_k$ .
- (iv)  $S_k = S_{x_r}$ , where  $x_r$  is the largest element in  $S_k$ .

PROOF : (i)  $k = \psi(a, k) = \psi(a, \psi(b, k)) = \psi(\psi(a, b), k)$ .

(ii) If  $x \in S_a$ , then  $\psi(x, a) = a$ . Hence

$$\psi(x, k) = \psi(x, \psi(a, k)) = \psi(\psi(x, a), k) = \psi(a, k) = k,$$

so that  $x \in S_k$ .

(iii) Let  $\psi(a, b) = k$ . Let  $x \in S_a$ . Then  $\psi(x, a) = a$ . Hence

$$\psi(x, k) = \psi(x, \psi(a, b)) = \psi(\psi(x, a), b) = \psi(a, b) = k.$$

Hence  $x \in S_k$  so that  $S_a \subseteq S_k$ . Similarly  $S_b \subseteq S_k$ .

(iv) If  $k \in S_k$ , then  $x_r = k$ . So we may assume that  $k \notin S_k$ . Hence  $x_r < k$ . Since  $x_r \in S_k$ , by (ii),  $S_{x_r} \subseteq S_k$ . Let  $x \in S_k$ . Then  $\psi(x, x_r) \in S_k$  by (i). Also,  $\psi(x, x_r) \geq x_r$ . Since  $x_r$  is the largest element in  $S_k$ ,  $\psi(x, x_r) = x_r$ . Hence  $x \in S_{x_r}$ , implying that  $S_k \subseteq S_{x_r}$ .

*Lemma 2.2*—If  $g$  is the unity of  $(F, +, \psi)$ , then  $g(k) = 0$  whenever  $k \notin S_k$ .

PROOF : If  $k \in S_k$  for every  $k$ , then there is nothing to prove. We assume that  $k \notin S_k$  for some  $k$ . Let  $t$  be the least positive integer such that  $t \notin S_t$  clearly  $t \geq 2$  since  $1 \in S_1$ . For  $1 \leq j < t$ ,  $j \in S_j$ . We have from (1.7),

$$g(t) = - \sum_{\substack{\psi(x, x_r) = t \\ x < t}} g(x)$$

where  $x_r$  is the largest in  $S_t$ . Now  $x < t$  implies that  $x \in S_x$ . Also, by (iii) of Lemma 2.1,  $\psi(x, x_r) = t$  implies that  $S_x \subseteq S_t$ . Hence  $x \in S_t$ . Since  $x_r \in S_t$ ,

(i) of Lemma 2.1 implies that  $t = \psi(x, x_r) \in S_t$  and this is false. Hence the sum on the right-hand side defining  $g(t)$  is an empty sum. Hence  $g(t) = 0$ . We assume that  $g(x) = 0$  whenever  $x \notin S_x$  and  $t \leq x < k$ .

Let  $k \notin S_k$ . By (1.7), we have

$$g(k) = - \sum_{\substack{\psi(x, x_r)=k \\ x < k}} g(x)$$

where  $x_r$  is the largest element in  $S_k$ . If  $1 \leq x < t$  and  $\psi(x, x_r) = k$ , then  $x \in S_x \subseteq S_k$ , implying that  $k = \psi(x, x_r) \in S_k$ . We may assume that  $x \geq t$ . For  $t \leq x < k$ , if  $x \notin S_x$ ,  $g(x) = 0$ , by our induction hypothesis. Arguing as before we obtain

$$g(k) = - \sum_{\substack{\psi(x, x_r)=k \\ t \leq x < k \\ x \in S_x}} g(x) = \text{empty sum} = 0$$

since  $k \notin S_k$ . We now prove

*Theorem 2.1*—Let  $g$  be defined by

$$g(k) = \begin{cases} 1 - \sum_{\substack{x < k \\ x \in S_k}} g(x), & \text{if } k \in S_k \\ 0, & \text{if } k \notin S_k. \end{cases} \quad \dots(2.1)$$

Then  $g$  is the unity of  $(F, +, \psi)$ .

**PROOF :** We shall prove that  $g$  satisfies (1.1).

Let  $x_r$  be the largest element in  $S_k$ . By (iv) of Lemma 2.1,  $S_k = Sx_r$ . Since  $x_r \in Sx_r$ , from (2.1), it is clear that

$$1 = \sum_{x \in Sx_r} g(x) = \sum_{x \in S_k} g(x). \quad \dots(2.2)$$

It remains to prove that if  $n$  and  $k$  are positive integers with  $n < k$ , then

$$\sum_{\psi(y, n)=k} g(y) = 0. \quad \dots(2.3)$$

We distinguish the following cases (in what follows we tacitly assume the results of Lemma 2.1):

*Case 1*—Let  $n \in S_n$ . Since  $g(y) = 0$  if  $y \notin S_y$ , we may assume that  $y \in S_y$  in the sum on the left-hand side of (2.3). Now,  $\psi(y, n) = k$  implies that  $S_y \subseteq S_k$  and  $S_n \subseteq S_k$ . Since  $y \in S_y$  and  $n \in S_n$ ,  $y, n \in S_k$ . Hence  $k = \psi(y, n) \in S_k$ . Thus we have  $n \in S_n \subseteq S_k$  and  $k \in S_k$ . Let  $S_k = \{x_1, x_2, \dots, x_r\}$  with  $x_1 < x_2 < \dots < x_r = k$ . Since  $n \in S_n \subseteq S_k$ , we assume that  $n = x_i$  where  $x_i \in S_{x_i}$ .

First we show that  $\sum_{\psi(y, x_i) = x_{i+1}} g(y) = 0$ . Since  $g(y) = 0$ , if  $y \notin S_y$ , in this sum, we may assume that  $y \in S_y$ . Now  $\psi(y, x_i) = x_{i+1}$  implies that  $y \in S_y \subseteq S_{x_{i+1}}$  and  $S_{x_i} \subseteq S_{x_{i+1}}$ . Since  $x_i \in S_{x_i}$ , we have  $x_i \in S_{x_{i+1}}$ ; this together with  $y \in S_{x_{i+1}}$  implies that  $x_{i+1} = \psi(y, x_i) \in S_{x_{i+1}}$ .

Let  $y \in S_{x_{i+1}}$  and  $y \notin S_{x_i}$ . Since  $x_i \in S_{x_{i+1}}$ ,  $\psi(y, x_i) \in S_{x_{i+1}}$ . Also,  $\psi(y, x_i) \geq x_i$ . Hence  $\psi(y, x_i) = x_i$  or  $x_{i+1}$ , since  $x_{i+1}$  is the largest element in  $S_{x_{i+1}}$  and no element of  $S_{x_{i+1}}$  can exist in the interval  $(x_i, x_{i+1})$  as  $S_{x_{i+1}} \subseteq S_k$ . Hence  $\psi(y, x_i) = x_{i+1}$  since  $\psi(y, x_i) = x_i$  implies that  $y \in S_{x_i}$ . Therefore by (2.2), since  $S_{x_i} \subseteq S_{x_{i+1}}$ , we have

$$0 = \sum_{\substack{y \in S_{x_{i+1}} \\ y \notin S_{x_i}}} g(y) = \sum_{\psi(y, x_i) = x_{i+1}} g(y).$$

Let us assume that

$$\sum_{\psi(y, x_i) = x_{i+s}} g(y) = 0$$

for all  $s$  with  $1 \leq s < t \leq r - i - 1$ . We shall prove that

$$\sum_{\psi(y, x_i) = x_{i+t}} g(y) = 0, \text{ if } x_{i+t} \in S_{x_{i+t}}.$$

Since  $\psi(y, x_i) = x_{i+t}$  implies that  $S_{x_i} \subseteq S_{x_{i+t}}$

and  $x_i \in S_{x_i}$ , we have  $x_i \in S_{x_{i+t}}$ .

If  $y \notin S_{x_i}$ ,  $y \in S_{x_{i+t}}$  then  $\psi(y, x_i) \in S_{x_{i+t}}$  and  $\psi(y, x_i) > x_i$ .

Hence  $\psi(y, x_i) = x_{i+1}$  or. Therefore

$$0 = \sum_{\substack{y \in S_{x_{i+t}} \\ y \notin S_{x_i}}} g(y) = \sum_{\substack{y \in S_{x_{i+t}} \\ \psi(y, x_i) = x_{i+1}}} g(y) + \dots + \sum_{\substack{y \in S_{x_{i+t}} \\ \psi(y, x_i) = x_{i+t}}} g(y). \quad \dots(2.4)$$

If  $x_{i+1} \notin S_{x_{i+t}}$ , then  $\psi(y, x_i) = x_{i+1}$  does not occur. Hence the first sum on the right-hand side of (2.4) vanishes. Similar remark applies to the other sums on the right-hand side of (2.4). Hence we may assume that  $x_{i+s} \in S_{x_{i+t}}$ , for  $s = 1, 2, \dots, t$ . So,  $S_{x_{i+s}} \subseteq S_{x_{i+t}}$ , for  $s = 1, 2, \dots, t$ . The variable  $y$  in each of the sums on the right-hand side of (2.4) can be assumed to be in  $S_y$  since  $g(y) = 0$  if  $y \notin S_y$ . This, implies that ' $y \in S_{x_{i+t}}$ ' need not be mentioned in each of the sums on the right-hand side of (2.4). Hence (2.4) can be written as



$$\begin{aligned}
 0 = \sum_{\substack{y \in S_{x_{i+t}} \\ y \notin S_{x_i}}} g(y) &= \sum_{\psi(y, x_i) = x_{i+1}} g(y) + \dots + \sum_{\psi(y, x_i) = x_{i+t-1}} g(y) \\
 &\quad + \sum_{\psi(y, x_i) = x_{i+t}} g(y). \quad \dots(2.5)
 \end{aligned}$$

By our induction hypothesis, the first  $t-1$  sums on the right side of (2.5) vanish and then we obtain  $0 = \sum_{\psi(y, x_i) = x_{i+t}} g(y)$ . The induction is complete. Therefore,

we have  $\sum_{\psi(y, x_i) = k} g(y) = 0$ , since  $k = x_r$ .

*Case 2— $n \notin S_n$ .* Suppose the equation  $\psi(y, n) = k$  has a solution  $y$  with  $y \in S_y$ . Then  $y \in S_y \subseteq S_k$  and  $S_n \subseteq S_k$ . Let  $t$  be the largest element in  $S_n$  so that  $t < n$ . If  $\psi(y, n) = k$ , then we also have

$$k = \psi(y, n) = \psi(y, \psi(t, n)) = \psi(\psi(y, t), n)$$

so that  $\psi(y, t)$  is also a solution of the equation  $\psi(x, n) = k$ . Let  $Y_1, Y_2, \dots, Y_r$  be all the elements of  $S_k$  which satisfy  $\psi(y_i, n) = k$ , and  $y_i \geq t$  for  $i = 1, 2, \dots, r$  and  $y_1 < y_2 < \dots < y_r$ . (We may note here that  $y_r = x_r$  the largest element in  $S_k$ ). Also, no  $y_i = t$ . For if  $y_i = t$ , then  $k = \psi(y_i, n) = \psi(t, n) = n$ . But  $n < k$ . So,  $y_i > t$  for  $i = 1, 2, \dots, r$ . We have

$$\sum_{\psi(y, n) = k} g(y) = \sum_{\substack{\psi(y, n) = k \\ \psi(y, t) = y_1}} g(y) + \sum_{\substack{\psi(y, n) = k \\ \psi(y, t) = y_2}} g(y) + \dots + \sum_{\substack{\psi(y, n) = k \\ \psi(y, t) = y_r}} g(y).$$

Now,  $\psi(y, t) = y_i$  and  $\psi(y_i, n) = k$  imply that

$$k = \psi(y_i, n) = \psi(\psi(y, t), n) = \psi(y, \psi(t, n)) = \psi(y, n)$$

since  $t \in S_n$ . Hence we have

$$\sum_{\psi(y, n) = k} g(y) = \sum_{\psi(y, t) = y_1} g(y) + \dots + \sum_{\psi(y, t) = y_r} g(y) \quad \dots(2.6)$$

since  $t < y_i$ ,  $i = 1, 2, \dots, r$  and  $t \in S_t$ , each sum on the right-hand side of (2.6) is a sum considered in Case 1 and hence vanishes. Thus

$$\sum_{\psi(y, n) = k} g(y) = 0.$$

The proof of Theorem 2.1 is complete.

We state without proof the following :

*Theorem 2.2—*Suppose  $\psi$  satisfies the postulates (I), (II) and (III) of §1 so that  $(F, +, \psi)$  is a commutative ring. If  $\psi(x, y) \geq \max\{x, y\}$  for all  $(x, y) \in T$ , then

the unity of  $(F, +, \psi)$  exists if and only if for each  $k \in \mathbb{Z}^+$ , equation  $\psi(x, k) = k$  has a solution. In such a case, the unity is given by (2.1).

*Remark 2.1 :* If  $\psi(x, y) < \max\{x, y\}$  for some  $(x, y) \in T$ , then the conclusion of Theorem 2.2 need not hold. For example, let  $T = \{(1, 2), (2, 1)\} \cup \{(k, k) : k \geq 2\}$  and  $\psi$  on  $T$  be defined by  $\psi(1, 2) = \psi(2, 1) = 1$  and  $\psi(k, k) = k$  for  $k \geq 2$ . Then  $\psi$  satisfies the postulates I, II and III of §1 and clearly for each  $k \in \mathbb{Z}^+$ ,  $\psi(x, k) = k$  has a solution. Note that  $\psi(2, 1) = 1 < 2 = \max\{2, 1\}$ . It can be easily shown that (for example using (1.1)) that  $(F, +, \psi)$  does not possess the unity.

*Remark 2.2 :* Let  $T = \{(2k, 2k), (2k-1, 2k), (2k, 2k-1) : k \in \mathbb{Z}^+\}$ . We define  $\psi : T \rightarrow \mathbb{Z}^+$  by  $\psi(x, y) = \min\{x, y\}$  for all  $(x, y) \in T$ . It can be shown that  $\psi$  satisfies the postulates I, II and III of §1 so that  $(F, +, \psi)$  is a commutative ring. Also using (1.1) it is not difficult to show that the function of defined by  $g(2k) = 1$  and  $g(2k-1) = 0$  for  $k = 1, 2, 3, \dots$ , is the unity of  $(F, +, \psi)$ . Clearly the condition  $\psi(x, y) \geq \max\{x, y\}$  for all  $(x, y) \in T$  is violated.

#### ACKNOWLEDGEMENT

The problems of this paper have been proposed to me by Professor N. V. Subrahmanyam more than ten years ago. I express my gratitude to him. Also, I wish to thank Dr S. Rame Gowda, Principal, Pondicherry Engineering College, Pondicherry for his constant encouragement.

#### REFERENCES

1. V. Sita Ramaiah, *Indian J. pure appl. Math.* **16** (1985), 994-1008.
2. V. Sita Ramaiah, *Indian J. pure appl. Math.* **19** (1988), 1-10.

## ITERATIVE METHODS OF SOLUTIONS FOR LINEAR AND QUASI LINEAR COMPLEMENTARITY PROBLEMS

R. N. MUKHERJEE AND H. L. VERMA

*Department of Applied Mathematics, Institute of Technology, Banaras Hindu  
University, Varanasi 221005*

(Received 12 August 1988; after revision 3 July 1989; accepted 27 July 1989)

The present work has been conceived out of a need to extend the method of Ahn to give an iterative procedure for approximating solution of a 'quasi-linear complementarity problem' (QLCP). We also give the bound for spectral radius of them odified matrix in the context of QLCP. Further we give more results on the modification of the algorithm of Pang for finding the solution of QLCP for a pair  $(M, q)$ , where  $M$  is symmetric and positive definite. The fixed parameter approach of Pang also has been modified to incorporate the variable parameter method in successive iteration process.

### 1. INTRODUCTION

The numerical method for LCP carries with it two methods, e. g., direct and indirect methods. Because of the complexity, the use of direct method is restricted for large size problems. Therefore iterative methods are well suited for such problems. The present work attempts to develop the procedure for finding approximate solutions of Quasi-Linear-Complementarity problems by iterative technique. Essentially this extends the earlier algorithm of Ahn<sup>1</sup> used for solution method for Linear Complementarity problems. Secondly, we also give an extension of a method of Pang<sup>5</sup> to incorporate QLCP for the same purpose. Our attempt next would be to briefly indicate the essential procedure of Pang<sup>5</sup> because we would refer that latter in our extensions.

Consider the symmetric LCP  $(q, M)$  :

$$q + Mx \geq 0, \quad x \geq 0, \quad \text{and} \quad x^T (q + Mx) = 0$$

where  $q \in R^n$  and  $M \in R^{n \times n}$  are given and  $x \in R^n$ . Let  $(B, C)$  be a  $Q$ -splitting of the matrix  $M$ , i. e.  $M = B + C$  is a  $Q$ -matrix [the LCP  $(q, B)$  has a solution for all vectors  $q$ ]. Let  $E^k$  be a non-negative diagonal matrix with  $E_{ii}^k < 1$ . Define the point to set algorithmic map  $A^k$  as follows: for all vectors  $x$ ,

$$A^k(x^k) = \text{solution set of the LCP } (q + Cx, B, E^k x^k). \quad \dots(1.1)$$

The latter LCP  $(r, B, s)$  is to find  $y$  so that

$$r + By \geq 0, \quad y \geq s \quad \text{and} \quad (y - s)^T (r + By) = 0.$$



The LCP  $(r, B, s)$  can be converted into the LCP  $(r + Bs, B)$  if we translate the variable  $x = y - s$  since  $B$  is a  $Q$ -matrix, the set  $A^k(x^k)$  is non-empty for all vectors  $x$ . Moreover a vector  $x^*$  solves the LCP  $(q, M)$  if and only if it is fixed point of the map  $A^k$  i. e.  $x^* \in A^k(x^*)$ .

We define an iterative technique for solving the LCP  $(q, M)$  given the diagonal matrix  $E^k$  and the  $Q$ -splitting  $(B, C)$  of the matrix  $M$ . Let  $x^0 \geq 0$  be an arbitrary non-negative vector. In general  $x^k \geq 0, k \geq 0$  let  $x^{k+1}$  be any vector in the set  $A^k(x^k)$ .

The motivation for using the map  $A^k(x^k)$  lies in the fact that the matrix  $E^k$  may satisfy the bound  $E_{ii}^k < 1$  after the iteration proceeds onwards after a fixed index  $k_0$ . This idea is compatible with the usual approach in the contraction mapping case where a fixed power of a mapping may be a contraction although the original map may not be a contraction.

If  $B$  is a  $P$ -matrix (A real matrix  $A \in R^{n \times n}$  is said to be a  $P$ -matrix if it has positive principal minors) then the set  $A^k(x^k)$  is singleton for all  $x$ . In this case, each  $x^{k+1}$  will be uniquely defined.

Pang<sup>4</sup> has given necessary and sufficient conditions on the matrix  $M$  on the convergence property, i. e. for all vector  $q$  and all starting vector  $x^0 \geq 0$ , each sequence  $\{x^k\}$  generated by the iterative technique will converge to some solution of the LCP  $(q, M)$ .

A Quasi-linear Complementarity (QLCP) can be stated as follows

Find  $z \in R^n$  such that

$$z - Qz \geq 0, Mz + q \geq 0, (z - Qz)^T (Mz + q) = 0. \quad \dots(1.2)$$

With splitting as indicated previously the point to-set algorithmic map takes the shape.

$$A^k(x^k) = \text{Solution set of LCP } (q + BQx^k + Cx^k, B, E^k(1 - Q)x^k) \quad \dots(1.3)$$

Variantly  $x^*$  solves the QLCP (1.2) if and only if  $(1 - Q)x^*$  is a point in range of the set valued map  $A$ , i. e. a point in  $A(x^*)$ . The result connected with the convergence of various dual iterative techniques for the solution of strictly convex quadratic program

$$\min_{(1-Q)x \geq 0} f(x) = q^T x + \frac{1}{2} x^T Mx \quad \dots(1.4)$$

can be derived by the methods of Pang<sup>5</sup>.

We explain some matrix notations as follows : If  $A$  is an  $n \times m$  matrix,  $\alpha$  and  $\beta$  are subsets of  $\{1, \dots, n\}$  and  $\{1, \dots, m\}$ , respectively, by  $A_{\alpha\beta}$  we denote the sub-matrix of  $A$  whose rows and columns are indexed by  $\alpha$  and  $\beta$  respectively. If  $\alpha = \{1, \dots, n\}$ ,

we denote by  $A_\beta$  the submatrix whose columns of  $A$  are indexed by  $\beta$ ; similar definition applies to  $A_\alpha$ .

## 2. PRELIMINARIES

We restate again the QLCP as :

find  $z \in R^n$ , such that

$$z - Qz \geq 0, Mz + q \geq 0, (z - Qz)^T (Mz + q) = 0 \quad \dots(2.1)$$

where  $M$  is an  $n \times n$  real and non-symmetric matrix,  $q$  is  $n \times 1$  vector. If we take  $Q = 0$  in QLCP we get the same LCP as in Ahn<sup>1</sup>.

First of all we describe the notations which occur in the QLCP. All matrices and vectors are real. A matrix  $A$  with  $m$ -rows,  $n$ -columns is denoted by  $R^{m \times n}$ . Row  $i$  of matrix  $A$  is denoted by  $A_i$  and column  $j$  by  $A_j$  and the element in row  $i$  and column  $j$  by  $A_{ij}$ . The transpose of a matrix is denoted by super script  $T$ , such as the transpose of the matrix  $A$  is given by  $A^T$ ,  $|A|$  denotes the matrix obtained from the real matrix  $A \in R^{m \times n}$  by replacing each element  $A_{ij}$  by its absolute value.

If  $x \in R^n$ ,  $x_+$  denotes the vector with elements

$$(x_+)_j = \max \{0, x_j\}; j = 1, 2, \dots, n.$$

For any  $x$  and  $y$  in  $R^n$ , it can be easily shown that

$$(i) \quad (x + y)_+ \leq x_+ + y_+,$$

$$(ii) \quad x \leq y \Rightarrow x_+ \leq y_+.$$

A real matrix  $A \in R^{n \times n}$  is said to be a  $Z$ -matrix (a  $P$ -matrix) if it has non-positive off diagonal entries (positive primal minors).

A square matrix with non-positive off diagonal elements and with a non-negative inverse is called an  $M$ -matrix. It can be easily shown that a matrix which is both a  $Z$ -matrix and a  $P$ -matrix is an  $M$ -matrix (or Minkowski matrix).

Given any real matrix  $A \in R^{n \times n}$ , we define its comparison matrix

$$A_c = (C_{ij})$$

by

$$C_{ii} = |A_{ii}|$$

and

$$C_{ij} = -|A_{ij}|, i \neq j, i, j = 1, 2, \dots, n$$

This definition is due to Verga<sup>6</sup>

## 3. ITERATIVE ALGORITHM

For solving QLCP (2.1) we describe the general fundamental algorithm.

*Lemma 3.1*—Let  $M \in R^{n \times n}$  and  $E$  be any positive diagonal matrix, then,

$$z - Qz \geq 0, Mz + q \geq 0, (z - Qz)^T (Mz + q) = 0$$

$$\Leftrightarrow z = \{(1 - Q)z - \omega E(Mz + q)\}_+, \text{ for all or some } \omega > 0.$$

Its proof is same as in Mangasarian<sup>2</sup>. This result can be transformed to a fixed point problem for solving the equation  $z = f(z)$

where

$$f(z) = \{(1 - Q)z - \omega E(Mz + q)\}_+.$$

This result readily leads to the following general algorithm suggested by Mangasarian<sup>2</sup>. We modify this algorithm at certain steps.

*Algorithm 3.1*—Let  $z^0 \geq 0$ , compute

$$\begin{aligned} z^{k+1} = & \lambda [(1 - Q)z^k - \omega E^k(Mz^k + q + K^k(1 - Q)(z^{k+1} - z^k))]_+ \\ & + (1 - \lambda)(1 - Q)z^k \end{aligned} \quad \dots(3.1)$$

where

$$k = 0, 1, \dots, 0 < \lambda \leq 1, \omega > 0$$

and  $\{E^k\}$  and  $\{K^k\}$  are bounded sequences of matrices in  $R^{n \times n}$ , with each  $E^k$  being a positive diagonal matrix satisfying  $E^k \geq \alpha I$ , for some  $\alpha > 0$

where  $I$  is the identity matrix.

For the symmetric case Mangasarian has established convergence criteria of this general algorithm. We simplify this algorithm by setting

$$\lambda = 1, E^k = E, K^k = K, \text{ for each } k.$$

*Remark* : As has been indicated in the conclusion, we can relax the above criteria for fixing the matrix powers  $E^k$  and  $K^k$  as constant matrices to derive certain variable parameter algorithm as well.

*Algorithm 3.2*—Let  $z^0 \geq 0$ , compute,

$$\begin{aligned} z^{k+1} = & [(1 - Q)z^k - \omega E(Mz^k + q + K(1 - Q)(z^{k+1} - z^k))]_+ \\ k = & 0, 1, \dots \end{aligned} \quad \dots(3.2)$$

where  $\omega > 0$ ,  $E$  is a positive diagonal matrix and  $K$  is either strictly upper triangular or lower triangular matrix. Convergence properties for non-symmetric situations can not be established relying on the descent function of the form

$$\frac{1}{2} x^T M x + q^T x$$

so the recursive relation between two successive iterations will be utilized here.



#### 4. CONVERGENCE PROPERTIES

First of all we develop the fundamental recursive inequality for Algorithm 3.2 which will be the basis of convergence. This inequality is derived from the inequality properties of  $x_+$  and  $y_+$ .

*Lemma 4.1*—The  $k$ th and  $(k + 1)$ th solutions  $z^k$  and  $z^{k+1}$  satisfy the partial ordering recursive inequality :

$$\begin{aligned} \|z^{k+1} - z^k\| &\leq (I - \omega E \|K\| I - Q)^{-1} \|(1 - Q)(I + \omega EK) \\ &\quad - \omega EM\|z^k - z^{k-1}\|. \end{aligned} \quad \dots(4.1)$$

From this Lemma we can produce a condition for the sequence  $\{z^k\}$  of Algorithm 3.2 to be bounded and have an accumulation point which solves the QLCP (2.1).

If we put  $Q = 0$  in (4.1) we have

$$\|z^{k+1} - z^k\| \leq (I - \omega E \|K\|)^{-1} \|I - \omega E(M - K)\|z^k - z^{k-1}\|$$

which is the standard form given by Ahn<sup>1</sup>.

*Theorem 4.1*—Suppose that the given iteration parameter  $\omega$ ,  $E$ ,  $K$  and the underlying matrix  $M$  satisfy

$$\mu((I - \omega E \|K\|)^{-1} \|I - \omega E(M - K)\|) < 1 \quad \dots(4.2)$$

where  $\mu(\cdot)$  denotes the spectral radius; then the sequence  $\{z^k\}$  of Algorithm 3.2 converges to a solution  $z^*$  of QLCP.

The proof is similar to Ahn<sup>1</sup>.

Here also we can find the same spectral radius which is established by Ahn<sup>1</sup>, simply by taking  $Q = 0$ , in (4.2), viz.,

$$\mu((I - \omega E \|K\|)^{-1} \|I - \omega E(M - K)\|) < 1.$$

We shall start the next section in which we modify the same result on the convergence of iterative methods for the symmetric QLCP. Pang had developed necessary and sufficient condition (for a fixed parameter) for the convergence of iterative method and for solving each individual LCP. We shall extend the method of Pang to incorporate it for the treatment of QLCP.

#### 5. NON-DEGENERATE CASE

We classify our analysis into two cases which depends on the nature of the matrix, i. e. either the matrix is non-degenerate or positive semi-definite. Since we know that the matrix  $M$  is non-degenerate if all its principal minors are non-zero and the same case for the non-degeneracy in linear complementarity theory<sup>3</sup> the matrix  $M$  is non-degenerate if and only if the LCP  $(q, M)$  has a finite number of solutions for all vectors  $q$ .

The following theorem is the main result of this section.

**Theorem 5.1**—Let  $M$  be symmetric and non-degenerate matrix. Let  $(B, C)$  be regular  $Q$ -splitting of the matrix  $M$ . Let  $E^k$  be a non-negative diagonal matrix, with  $E_{ii}^k < 1$ , for all  $i$ . Then the following statements are equivalent:

- (A) for some vector  $q$  and any initial vector  $x^0 \geq 0$ , any sequence  $\{x^k\}$  satisfying  $(1 - Q) x^{k+1} \in A^k(x^k)$  is bounded and thus has at least one accumulation point, moreover, any such point solves the QLCP  $(q, M)$ .
- (B) for some vector  $q$ , the quadratic function  $f(x) = q^T x + \frac{1}{2} x^T Mx$  is bounded below for  $(1 - Q) x \geq 0$ .
- (C) for some vector  $q$  and any initial vector  $x^0 > 0$ , any sequence  $\{x^k\}$  satisfying  $(1 - Q) x^{k+1} \in A^k(x^k)$  converges to solution of the QLCP  $(q, M)$ .

Proof can be given in a line of the arguments given in Pang<sup>5</sup>.

## 6. CONCLUDING REMARKS

As one of the concluding remarks we would like to point out that in the case of quasi-linear complementarity problems the algorithm which was developed in section 3, for the iterative solution technique can as well be generalized for variable parameters such as the case when the assumptions  $E = E^k$  and  $K = K^k$  are relaxed and we take uniformly bounded (by matrix norm) matrices  $E^k$  and  $K^k$  in the iterative process of the algorithm itself. The variable parameter algorithms are still possible to find the fixed point for the set-valued maps  $A(x^k)$ .

## ACKNOWLEDGEMENT

The authors are grateful to the referee for his valuable comments which enabled the authors to improve over the earlier version of the manuscript.

## REFERENCES

1. B. H. Ahn, *Optimization Theory Applic.* **33** (1981), 175-85.
2. O. L. Mangasarian, *Optimization Theory Applic.* **22** (1977), 465-85.
3. K. G. Murthy, *Linear Algebra Applic.* **5** (1972), 65-108.
4. J. S. Pang, *J. Optimization Theory Applic.* **42** (1984), 1-18.
5. J. S. Pang, *J. Optimization Theory Applic.* **49** (1986), 108-34.
6. R. S. Verga, *Linear Algebra Applic.* **13** (1974), 1-9.

## ON SOME NEW DISCRETE INEQUALITIES IN TWO INDEPENDENT VARIABLES

B. G. PACHPATTE

*Department of Mathematics and Statistics, Marathwada University  
Aurangabad 431004, (Maharashtra)*

(Received 16 December 1988; accepted 23 May 1989)

The aim of this paper is to establish some new discrete inequalities in two independent variables which can be used as handy tools in the qualitative analysis of a new class of finite difference equations involving two independent variables.

### 1. INTRODUCTION

The fundamental role played by the discrete inequalities in the development of the theory of finite difference equations and numerical analysis is well known. A large number of papers dealing with discrete inequalities and their applications have appeared during the last few years, see<sup>1-10</sup> and some of the references given therein. Although stimulating research works have been undertaken in this direction, there are still a number of interesting classes of multidimensional finite difference equations which needs new types of discrete inequalities in their analysis. Our objective here is to present some new discrete inequalities in two independent variables which can be used as handy tools in the qualitative analysis of a new class of finite difference equations in two independent variables. In order to convey the importance of our results to the literature, we present applications of some of our inequalities to the study of boundedness, uniqueness and continuous dependence of the solutions of a new class of fourth order finite difference equations in two independent variables.

### 2. STATEMENT OF RESULTS

We first summarise some basic notations and definitions which will be used throughout this paper. Let  $N_0 = \{0, 1, 2, \dots\}$ . The expression  $u(0) + \sum_{s=0}^{n-1} b(s)$  represents a solution of the linear difference equation  $\Delta u(n) = b(n)$  for  $n \in N_0$ , where  $\Delta$  is the operator defined by  $\Delta u(n) = u(n+1) - u(n)$ . The expression  $u(0) + \sum_{s=0}^{n-1} b(s)$  represents a solution of the linear difference equation  $u(n+1) = b(n)u(n)$  for  $n \in N_0$ . We use the usual convention of writing  $\sum_{s \in \Phi} b(s) = 0$

and  $\prod_{s \in \Phi} b(s) = 1$ , if  $\Phi$  is the empty set. We also use the the following notations of the operators

$$\Delta_1 u(m, n) = u(m+1, n) - u(m, n),$$

$$\Delta_2 u(m, n) = u(m, n+1) - u(m, n)$$

for  $m, n \in N_0$ . We often use the letters  $m$  and  $n$  to denote the two independent variables which are members of  $N_0$ .

For convenience we list the following hypotheses :

- (H<sub>1</sub>)  $u(m, n)$  and  $h(m, n)$  are real-valued nonnegative functions defined for  $m, n \in N_0$ .
- (H<sub>2</sub>)  $p_1(m, n)$ ,  $p_2(m, n)$ ,  $p_3(m, n)$  are real-valued positive functions defined for  $m, n \in N_0$ .
- (H<sub>3</sub>)  $a(m, n)$  is real-valued, positive and nondecreasing function in both the variables  $m$  and  $n$  in  $N_0$ .
- (H<sub>4</sub>)  $u(m, n) \geq u_0 \geq 0$ ,  $u_0$  is a constant,  $h(m, n) > 0$  are real-valued functions defined for  $m, n \in N_0$ .
- (H<sub>5</sub>)  $g(u)$  is continuous, nondecreasing real-valued function defined on an interval  $I = [u_0, \infty)$ ,  $u_0 \geq 0$  is a constant, and  $g(u) > 0$  on  $(u_0, \infty)$ ,  $g(u_0) = 0$ .
- (H<sub>6</sub>)  $q_1(m, n)$ ,  $q_2(m, n)$ ,  $q_3(m, n)$  are real-valued positive functions defined for  $m, n \in N_0$ .
- (H<sub>7</sub>)  $W(u)$  is continuous, nondecreasing and submultiplicative real-valued function defined on an interval  $I$ , and  $W(u) > 0$  on  $(u_0, \infty)$ ,  $W(u_0) = 0$ .

A useful two independent variable discrete inequality is embodied in the following theorem.

*Theorem 1*—Suppose (H<sub>1</sub>) and (H<sub>2</sub>) are true. If

$$u(m, n) \leq c + \sum_{x=0}^{m-1} \frac{1}{p_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \times \sum_{t=0}^{y-1} h(s, t) u(s, t) \quad \dots(1)$$

for  $m, n \in N_0$ , where  $c$  is a nonnegative constant, then

$$u(m, n) \leq c \prod_{x=0}^{m-1} \left[ 1 + \frac{1}{p_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \right]$$



$$\times \sum_{t=0}^{y-1} h(s, t) \Big] \quad \dots(2)$$

for  $m, n \in N_0$ .

A slightly different version of Theorem 1 is given in the following theorem.

*Theorem 2*—Suppose  $(H_1)$ ,  $(H_2)$  and  $(H_3)$  are true. If

$$\begin{aligned} u(m, n) \leq a(m, n) + \sum_{x=0}^{m-1} \frac{1}{p_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \\ \times \sum_{t=0}^{y-1} h(s, t) u(s, t) \end{aligned} \quad \dots(3)$$

for  $m, n \in N_0$ , then

$$\begin{aligned} u(m, n) \leq a(m, n) \prod_{x=0}^{m-1} \left[ 1 + \frac{1}{p_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \right. \\ \left. \times \sum_{t=0}^{y-1} h(s, t) \right] \end{aligned} \quad \dots(4)$$

for  $m, n \in N_0$ .

Another interesting and useful discrete inequality is established in the following theorem

*Theorem 3*—Suppose  $(H_2)$ ,  $(H_4)$  and  $(H_5)$  are true. If

$$\begin{aligned} u(m, n) \leq c + \sum_{x=0}^{m-1} \frac{1}{p_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \\ \times \sum_{t=0}^{y-1} h(s, t) g(u(s, t)) \end{aligned} \quad \dots(5)$$

for  $m, n \in N_0$ , where  $c$  is a nonnegative constant, then for  $0 \leq m \leq m_1$ ,  $0 \leq n \leq n_1$ ,  $m, m_1, n, n_1 \in N_0$ ,

$$\begin{aligned} u(m, n) \leq G^{-1} \left[ G(c) + \sum_{x=0}^{m-1} \frac{1}{p_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \right. \\ \left. \times \sum_{t=0}^{y-1} h(s, t) \right] \end{aligned} \quad \dots(6)$$

where

$$G(r) = \int_{r_0}^r \frac{dy}{g(y)}, \quad r \geq r_0 \text{ with } r_0 > u_0 \quad \dots(7)$$

$G^{-1}$  is the inverse of  $G$  and  $m_1, n_1 \in N_0$  are chosen so that

$$G(c) + \sum_{x=0}^{m-1} \frac{1}{p_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \sum_{t=0}^{y-1} \\ \times h(s, t) \in \text{Dom}(G^{-1})$$

for  $m, n \in N_0$  and  $0 \leq m \leq m_1, 0 \leq n \leq n_1$ .

We next establish the following more general inequality which may be convenient in some applications.

*Theorem 4*—Suppose  $(H_1)$ ,  $(H_2)$ ,  $(H_6)$  and  $(H_7)$  are true. If

$$u(m, n) \leq c + \sum_{x=0}^{m-1} \frac{1}{p_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \\ \times \sum_{t=0}^{y-1} h(s, t) u(s, t) \\ + \sum_{x=0}^{m-1} \frac{1}{q_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{q_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{q_3(s, y)} \\ \times \sum_{t=0}^{y-1} k(s, t) W(u(s, t)) \quad \dots(8)$$

for  $m, n \in N_0$ , where  $c$  is a nonnegative constant and  $k(m, n)$  is a real-valued non-negative function defined for  $m, n \in N_0$ , then for  $0 \leq m \leq m_2, 0 \leq n \leq n_2, m, m_2, n, n_2 \in N_0$

$$u(m, n) \leq Q(m, n) \Omega^{-1} \left[ \Omega(c) + \sum_{x=0}^{m-1} \frac{1}{q_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{q_2(s, n)} \right. \\ \left. \times \sum_{y=0}^{n-1} \frac{1}{q_3(s, y)} \sum_{t=0}^{y-1} k(s, t) W(Q(s, t)) \right] \quad \dots(9)$$

where

$$Q(m, n) = \prod_{x=0}^{m-1} \left[ 1 + \frac{1}{p_1(x, n)} \sum_{s=0}^{x-1} \frac{p}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \right. \\ \left. \times \sum_{t=0}^{y-1} h(s, t) \right] \quad \dots(10)$$

and

$$\Omega(r) = \int_{r_0}^r \frac{dy}{W(y)}, \quad r \geq u_0 \text{ with } r_0 > u_0 \quad \dots(11)$$

$\Omega^{-1}$  is the inverse of  $\Omega$  and  $m_2, n_2 \in N_0$  are chosen so that

$$\Omega(c) + \sum_{x=0}^{m-1} \frac{1}{q_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{q_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{q_3(s, y)} \\ \times \sum_{t=0}^{y-1} k(t, s) W(Q(t, s)) \in \text{Dom}(\Omega^{-1})$$

for  $m, n \in N_0$  and  $0 \leq m \leq m_2, 0 \leq n \leq n_2$ .

### 3. PROOFS OF THEOREMS 1-4

In order to establish the inequality (2) in Theorem 1, we first assume that  $c > 0$  and define a function  $z(m, n)$  by

$$z(m, n) = c + \sum_{x=0}^{m-1} \frac{1}{p_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \\ \times \sum_{t=0}^{y-1} h(s, t) w(s, t). \quad \dots(12)$$

From (12) it is easy to observe that

$$z(0, n) = z(m, 0) = c \quad \dots(13)$$

and

$$p_1(m, n) \Delta_1 z(m, n) = \sum_{s=0}^{m-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \sum_{t=0}^{y-1} h(s, t) w(s, t) \quad \dots(14)$$

$$p_2(m, n) \Delta_1 [p_1(m, n) \Delta_1 z(m, n)] = \sum_{y=0}^{n-1} \frac{1}{p_3(m, y)} \sum_{t=0}^{y-1} h(m, t) u(m, t) \quad \dots(15)$$

$$p_3(m, n) \Delta_2 [p_2(m, n) \Delta_1 [p_1(m, n) \Delta_1 z(m, n)]] = \sum_{t=0}^{n-1} h(m, t) u(m, t) \quad \dots(16)$$

$$\Delta_2 [p_3(m, n) \Delta_2 [p_2(m, n) \Delta_1 [p_1(m, n) \Delta_1 z(m, n)]]] = h(m, n) u(m, n). \quad \dots(17)$$

Using the fact that  $(u, m, n) \leq z(m, n)$  in (17) we have

$$\Delta_2 (p_3(m, n) \Delta_2 [p_2(m, n) \Delta_1 [p_1(m, n) \Delta_1 z(m, n)]]] \leq h(m, n) z(m, n). \quad \dots(18)$$

From the definition of  $u(m, n)$  we observe that  $z(m, n) \leq z(m, n+1)$  for  $m, n \in N_0$ . Using this fact in (18) we see that

$$\begin{aligned} & \frac{p_3(m, n+1) \Delta_2 [p_2(m, n+1) \Delta_1 [p_1(m, n+1) \Delta_1 z(m, n+1)]]}{z(m, n+1)} \\ & - \frac{p_3(m, n) \Delta_2 [p_2(m, n) \Delta_1 [p_1(m, n) \Delta_1 z(m, n)]]}{z(m, n+1)} \leq h(m, n). \end{aligned} \quad \dots(19)$$

From (19) and the fact that  $p_3(m, n) \Delta_2 [p_2(m, n) \Delta_1 [p_1(m, n) \Delta_1 z(m, n)]] \geq 0$  from (16), we observe that

$$\begin{aligned} & \frac{p_3(m, n+1) \Delta_2 [p_2(m, n+1) \Delta_1 [p_1(m, n+1) \Delta_1 z(m, n+1)]]}{z(m, n+1)} \\ & - \frac{p_3(m, n) \Delta_2 [p_2(m, n) \Delta_1 [p_1(m, n) \Delta_1 z(m, n)]]}{z(m, n)} \leq h(m, n). \end{aligned} \quad \dots(20)$$

Now keeping  $m$  fixed in (20), set  $n = t$  and sum over  $t = 0, 1, 2, \dots, n-1$  and use the fact that  $p_3(m, 0) \Delta_2 [p_2(m, 0) \Delta_1 [p_1(m, 0) \Delta_1 z(m, 0)]] = 0$ , from (16), to obtain the estimate

$$\frac{p_3(m, n) \Delta_2 [p_2(m, n) \Delta_1 [p_1(m, n) \Delta_1 z(m, n)]]}{z(m, n)} \leq \sum_{t=0}^{n-1} h(m, t). \quad \dots(21)$$

From (21) and in view of the facts that  $z(m, n) \leq z(m, n+1)$  and  $p_2(m, n) \Delta_1 [p_1(m, n) \Delta_1 z(m, n)] \geq 0$ , we observe that

$$\frac{p_2(m, n+1) \Delta_1 [p_1(m, n+1) \Delta_1 z(m, n+1)]}{z(m, n+1)}$$



$$\begin{aligned}
 & - \frac{p_2(m, n) \Delta_1 [p_1(m, n) \Delta_1 z(m, n)]}{z(m, n)} \\
 & \leq \frac{1}{p_3(m, n)} \sum_{t=0}^{n-1} h(m, t). \quad \dots (22)
 \end{aligned}$$

Keeping  $m$  fixed in (22) set  $n = y$  and sum over  $y = 0, 1, 2, \dots, n-1$  and use the fact that  $p_2(m, 0) \Delta_1 [p_1(m, 0) \Delta_1 z(m, 0)] = 0$  from (15), to obtain the estimate

$$\frac{p_2(m, n) \Delta_1 [p_1(m, n) \Delta_1 z(m, n)]}{z(m, n)} \leq \sum_{y=0}^{n-1} \frac{1}{p_3(m, y)} \sum_{t=0}^{y-1} h(m, t). \quad \dots (23)$$

From (23) and in view of the facts that  $z(m, n) \leq z(m+1, n)$  and  $p_1(m, n) \Delta_1 z(m, n) \geq 0$  from (14), we observe that

$$\begin{aligned}
 & \frac{p_1(m+1, n) \Delta_1 z(m+1, n)}{z(m+1, n)} - \frac{p_1(m, n) \Delta_1 z(m, n)}{z(m, n)} \\
 & \leq \frac{1}{p_2(m, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(m, y)} \sum_{t=0}^{y-1} h(m, t). \quad \dots (24)
 \end{aligned}$$

Now keeping  $n$  fixed in (24), set  $m = s$  and sum over  $s = 0, 1, 2, \dots, m-1$  and use the fact that  $p_1(0, n) \Delta_1 z(0, n) = 0$  from (14), to obtain the estimate

$$\frac{\Delta_1 z(m, n)}{z(m, n)} \leq \frac{1}{p_1(m, n)} \sum_{s=0}^{m-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \sum_{t=0}^{y-1} h(s, t). \quad \dots (25)$$

From (25) we see that

$$\begin{aligned}
 z(m+1, n) & \leq z(m, n) \left[ 1 + \frac{1}{p_1(m, n)} \sum_{s=0}^{m-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \right. \\
 & \quad \times \left. \sum_{t=0}^{y-1} h(s, t) \right]. \quad \dots (26)
 \end{aligned}$$

Now keeping  $n$  fixed in (26), set  $n = x$  and substitute  $x = 0, 1, 2, \dots, m-1$  successively and use the fact that  $z(0, n) = c$  from (13), to obtain the estimate

$$z(m, n) \leq c \prod_{x=0}^{m-1} \left[ 1 + \frac{1}{p_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \right]$$

$$\times \sum_{t=0}^{y-1} h(s, t) \Big].$$

Substituting this bound on  $z(m, n)$  on the right side of (1) we obtain the inequality in (2).

Now suppose  $c = 0$ . Then from (1) we see that the inequality

$$\begin{aligned} u(m, n) &\leq \epsilon + \sum_{x=0}^{m-1} \frac{1}{p_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \\ &\quad \times \sum_{t=0}^{y-1} h(s, t) u(s, t) \end{aligned}$$

holds for every arbitrary positive number  $\epsilon$  and  $m, n \in N_0$ , which by the above argument yields the estimate

$$\begin{aligned} u(m, n) &\leq \epsilon \prod_{x=0}^{m-1} \left[ 1 + \frac{1}{p_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \right. \\ &\quad \times \left. \sum_{t=0}^{y-1} h(s, t) \right]. \end{aligned} \quad \dots(27)$$

Since  $u(m, n) \geq 0$  and  $\epsilon > 0$  is arbitrary number independent of  $m, n$  then from (27) it follows that  $u(m, n) = 0$ . This completes the proof of Theorem 1.

Since  $a(m, n)$  is positive and nondecreasing, we observe from (3) that

$$\begin{aligned} \frac{u(m, n)}{a(m, n)} &\leq 1 + \sum_{x=0}^{m-1} \frac{1}{p_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \\ &\quad \times \sum_{t=0}^{y-1} h(s, t) \frac{u(s, t)}{a(s, t)}. \end{aligned}$$

Now an application of Theorem 1 yields the required bound in (4) and the proof of Theorem 2 is complete.

In order to establish the inequality (6) in Theorem 3, let  $\epsilon > 0$  and  $u_*(m, n) = u(m, n) + \epsilon \geq u_0$  for all  $m, n \in N_0$ . Then from (5) we see that

$$\begin{aligned}
u_{\epsilon}(m, n) &\leq c + \epsilon + \sum_{x=0}^{m-1} \frac{1}{p_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \\
&\quad \times \sum_{t=0}^{y-1} h(s, t) g(u_{\epsilon}(s, t) - \epsilon) \\
&\leq c + \epsilon + \sum_{x=0}^{m-1} \frac{1}{p_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \\
&\quad \times \sum_{t=0}^{y-1} h(s, t) g(u_{\epsilon}(s, t)). \quad \dots (28)
\end{aligned}$$

Define a function  $z(m, n)$  by

$$\begin{aligned}
z(m, n) &= c + \epsilon + \sum_{x=0}^{m-1} \frac{1}{p_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \\
&\quad \times \sum_{t=0}^{y-1} h(s, t) g(u_{\epsilon}(s, t)). \quad \dots (29)
\end{aligned}$$

From (29) it is easy to observe that

$$z(m, 0) = z(0, n) = c + \epsilon \quad \dots (30)$$

and

$$p_1(m, n) \Delta_1 z(m, n) = \sum_{s=0}^{m-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \sum_{t=0}^{y-1} h(s, t) g(u_{\epsilon}(s, t)). \quad \dots (31)$$

$$p_2(m, n) \Delta_1 [p_1(m, n) \Delta_1 z(m, n)] = \sum_{y=0}^{n-1} \frac{1}{p_3(m, y)} \sum_{t=0}^{y-1} h(m, t) g(u_{\epsilon}(m, t)) \quad \dots (32)$$

$$p_3(m, n) \Delta_2 [p_2(m, n) \Delta_1 [p_1(m, n) \Delta_1 z(m, n)]] = \sum_{t=0}^{n-1} h(m, t) g(u_{\epsilon}(m, t)) \quad \dots (33)$$

$$\Delta_2 [p_3(m, n) \Delta_2 [p_2(m, n) \Delta_1 [p_1(m, n) \Delta_1 z(m, n)]]] = h(m, n) g(u_{\epsilon}(m, n)). \quad \dots (34)$$

Using the fact that  $u_4(m, n) \leq z(m, n)$  in (34) we have

$$\Delta_2 [p_3(m, n) \Delta_2 [p_2(m, n) \Delta_1 [p_1(m, n) \Delta_1 z(m, n)]]] \leq h(m, n) g(z(m, n)). \quad \dots(35)$$

From the definition of  $z(m, n)$  in (29) we observe that  $z(m, n) \leq z(m, n+1)$  for  $m, n \in N_0$ . Using this and the fact that

$$p_3(m, n) \Delta_2 [p_2(m, n) \Delta_1 [p_1(m, n) \Delta_1 z(m, n)]] \geq 0$$

from (33), we observe from (35) that

$$\begin{aligned} & \frac{p_3(m, n+1) \Delta_2 [p_2(m, n+1) \Delta_1 [p_1(m, n+1) \Delta_1 z(m, n+1)]]}{g(z(m, n+1))} \\ & - \frac{p_3(m, n) \Delta_2 [p_2(m, n) \Delta_1 [p_1(m, n) \Delta_1 z(m, n)]]}{g(z(m, n))} \leq h(m, n). \end{aligned} \quad \dots(36)$$

Now by following exactly the same steps as in the proof of Theorem 1 below the inequality (20) up to the inequality (25) with suitable changes, we obtain

$$\frac{\Delta_1 z(m, n)}{g(z(m, n))} \leq \frac{1}{p_1(m, n)} \sum_{s=0}^{m-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \sum_{t=0}^{y-1} h(s, t). \quad \dots(37)$$

From (7) and (37) we have

$$\begin{aligned} G(z(m+1, n)) - G(z(m, n)) &= \int_{z(m, n)}^{z(m+1, n)} \frac{dy}{g(y)} \leq \frac{\Delta_1 z(m, n)}{g(z(m, n))} \\ &\leq \frac{1}{p_1(m, n)} \sum_{s=0}^{m-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \\ &\quad \times \sum_{t=0}^{y-1} h(s, t). \end{aligned} \quad \dots(38)$$

Now keeping  $n$  fixed in (38), set  $m = x$  and sum over  $x = 0, 1, 2, \dots, m-1$  to obtain the estimate

$$\begin{aligned} G(z(m, n)) &\leq G(c + \epsilon) + \sum_{x=0}^{m-1} \frac{1}{p_1(x, n)} \sum_{s=0}^{n-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \\ &\quad \times \sum_{t=0}^{y-1} h(s, t). \end{aligned} \quad \dots(39)$$



The bound in (6) now follows by substituting the bound for  $z(m, n)$  from (39) in (28) and letting  $\epsilon \rightarrow 0$ . The subintervals of  $N_0$  for  $m, n$  are obvious and the proof of Theorem 3 is complete.

In order to prove the inequality (9) in Theorem 4, let  $\epsilon > 0$  and  $u_\epsilon(m, n) = u(m, n) + \epsilon \geq u_0$  for  $m, n \in N_0$ . Then from (8) we see that

$$\begin{aligned}
 u_\epsilon(m, n) &\leq c + \epsilon + \sum_{x=0}^{m-1} \frac{1}{p_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \\
 &\quad \times \sum_{t=0}^{y-1} h(s, t) (u_\epsilon(s, t) - \epsilon) + \sum_{x=0}^{m-1} \frac{1}{q_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{q_2(s, n)} \\
 &\quad \times \sum_{y=0}^{n-1} \frac{1}{q_3(s, n)} \sum_{t=0}^{y-1} k(s, t) W(u_\epsilon(s, t) - \epsilon) \\
 &\leq c + \epsilon + \sum_{x=0}^{m-1} \frac{1}{p_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \\
 &\quad \times \sum_{t=0}^{y-1} h(s, t) u_\epsilon(s, t) + \sum_{x=0}^{m-1} \frac{1}{q_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{q_2(s, n)} \\
 &\quad \times \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \sum_{t=0}^{y-1} k(s, t) W(u_\epsilon(s, t)). \quad \dots(40)
 \end{aligned}$$

Define

$$\begin{aligned}
 a(m, n) &= c + \epsilon + \sum_{x=0}^{m-1} \frac{1}{q_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{q_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{q_3(s, y)} \\
 &\quad \times \sum_{t=0}^{y-1} k(s, t) W(u_\epsilon(s, t)) \quad \dots(41)
 \end{aligned}$$

then (40) can be restated as

$$\begin{aligned}
 u_\epsilon(m, n) &\leq a(m, n) + \sum_{x=0}^{m-1} \frac{1}{p_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{p_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{p_3(s, y)} \\
 &\quad \times \sum_{t=0}^{y-1} h(s, t) u_\epsilon(s, t).
 \end{aligned}$$

Since  $a(m, n)$  is positive and nondecreasing function in both the variables  $m$  and  $n$ , we have from Theorem 2

$$u_\epsilon(m, n) \leq a(m, n) Q(m, n) \quad \dots(42)$$

where  $Q(m, n)$  is as defined in (10). Since  $W$  is submultiplicative, we have

$$W(u_\epsilon(m, n)) \leq W(a(m, n)) W(Q(m, n)). \quad \dots(43)$$

From (41) and (43) we have

$$\begin{aligned} a(m, n) &\leq c + \epsilon + \sum_{x=0}^{m-1} \frac{1}{q_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{q_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{q_3(s, y)} \\ &\quad \times \sum_{t=0}^{y-1} k(s, t) W(Q(s, t)) W(a(s, t)). \end{aligned}$$

Now by following the proof of Theorem 3 with suitable modifications we obtain

$$\begin{aligned} a(m, n) &\leq \Omega^{-1} [\Omega(c + \epsilon) + \sum_{x=0}^{m-1} \frac{1}{q_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{q_2(s, n)} \\ &\quad \times \sum_{y=0}^{n-1} \frac{1}{q_3(s, y)} \sum_{t=0}^{y-1} k(s, t) W(Q(s, t))]. \quad \dots(44) \end{aligned}$$

The desired bound in (9) now follows by substituting (44) in (42) and letting  $\epsilon \rightarrow 0$ . The subintervals of  $N_0$  for  $m$  and  $n$  are obvious. This completes the proof of Theorem 4.

#### 4. SOME APPLICATIONS

In this section, we present some applications of our results to the study of boundedness, uniqueness and continuous dependence of the solutions of a new class of nonlinear finite difference equations in two independent variables. Each of these applications could be stated formally as a theorem. This has not been done so as not to obscure the essential ideas with technical details.

*Example 1*—As a first application, we obtain a bound on the solution of a nonlinear fourth order finite difference equation

$$\Delta_2 [a_3(m, n) \Delta_2 [a_2(m, n) \Delta_1 [a_1(m, n) \Delta_1 u(m, n)]]] = f(m, n, u(m, n)) \quad \dots(45)$$

with the given boundary conditions at  $m = 0, n = 0$

$$u(0, n) = \phi_1(n)$$

$$a_1(0, n) \Delta_1 u(0, n) = \phi_2(n)$$

$$a_2(m, 0) \Delta_1 [a_1(m, 0) \Delta_1 u(m, 0)] = \psi_1(m)$$

$$a_3(m, 0) \Delta_2 [a_2(m, 0) \Delta_1 [a_1(m, 0) \Delta_1 u(m, 0)]] = \psi_2(m). \quad \dots(46)$$

Here  $a_1, a_2, a_3$  are real valued positive functions defined on  $N_0^2, f: N_0^2 \times R \rightarrow R$ , where  $R$  denotes the set of real numbers;  $\phi_1(n), \phi_2(n), \psi_1(m), \psi_2(m)$  are real-valued nonnegative functions defined for  $m, n \in N_0$ . We assume that

$$|f(m, n, u)| \leq h(m, n) |u| \quad \dots(47)$$

where  $h(m, n)$  is a real-valued nonnegative function defined for  $m, n \in N_0$ . It is easy to observe that the problem (45) – (46) is equivalent to the equation

$$u(m, n) = b(m, n) + \sum_{x=0}^{m-1} \frac{1}{a_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{a_2(s, n)} \sum_{y=0}^{s-1} \frac{1}{a_3(s, y)} \\ \times \sum_{t=0}^{y-1} f(s, t, u(s, t)) \quad \dots(48)$$

where

$$b(m, n) = \phi_1(n) + \sum_{x=0}^{m-1} \frac{1}{a_1(x, n)} \phi_2(n) + \sum_{x=0}^{m-1} \frac{1}{a_1(x, n)} \\ \sum_{s=0}^{x-1} \frac{1}{a_2(s, n)} \psi_1(s) + \sum_{x=0}^{m-1} \frac{1}{a_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{a_2(s, n)} \\ \times \psi_2(s) \sum_{y=0}^{s-1} \frac{1}{a_3(s, y)}. \quad \dots(49)$$

Suppose that

$$|b(m, n)| \leq k \quad \dots(50)$$

where  $k$  is a nonnegative constant. Using (47), (50) in (48) we have

$$|u(m, n)| \leq k + \sum_{x=0}^{m-1} \frac{1}{a_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{a_2(s, n)} \sum_{y=0}^{s-1} \frac{1}{a_3(s, y)} \\ \times \sum_{t=0}^{y-1} h(s, t) |u(s, t)|.$$

Now an application of Theorem 1 yields the bound on the solution  $u(m, n)$  of (45)-(46) in terms of the known functions.

*Example 2*—As a second application, we shall discuss the uniqueness of the solution of the problem (45)–(46). We assume that the function  $f$  in (45) satisfies

$$|f(m, n, u) - f(m, n, \bar{u})| \leq h(m, n) |u - \bar{u}| \quad \dots(51)$$

where  $h(m, n)$  is as in Example 1. The problem (45)–(46) is equivalent to the equation (48). Then for any two solutions  $u$  and  $\bar{u}$  of (45)–(46) we have

$$\begin{aligned} |u(m, n) - \bar{u}(m, n)| &\leq \epsilon + \sum_{x=0}^{m-1} \frac{1}{a_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{a_2(s, n)} \\ &\quad \times \sum_{y=0}^{n-1} \frac{1}{a_3(s, y)} \sum_{t=0}^{y-1} h(s, t) |u(s, t) - \bar{u}(s, t)| \quad \dots(52) \end{aligned}$$

where  $\epsilon > 0$  is arbitrary constant. The assumption (51) is used to get the inequality in (52). Now an application of Theorem 1 yields

$$\begin{aligned} &|u(m, n) - \bar{u}(m, n)| \\ &\leq \epsilon \left\{ \prod_{x=0}^{n-1} \left[ 1 + \frac{1}{a_1(x, n)} \sum_{s=x}^{x-1} \frac{1}{a_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{a_3(s, y)} \sum_{t=0}^{y-1} h(s, t) \right] \right\}. \end{aligned}$$

Since  $\epsilon > 0$  is arbitrary we have  $u = \bar{u}$  i. e. there is at most one solution of the problem (45) – (46).

*Example 3*—Our third application is an example of continuous dependence of the solution on the equation and boundary data. Consider the problem (45)–(46) in Example 1 and the problem

$$\Delta_2 [a_3(m, n) \Delta_2 [a_2(m, n) \Delta_1 [a_1(m, n) \Delta_1 z(m, n)]]] = F(m, n, z(m, n)) \quad \dots(53)$$

with the given boundary conditions at  $m = 0, n = 0$

$$\begin{aligned} z(0, n) &= \bar{\phi}_2(n) \\ a_1(0, n) \Delta_1 z(0, n) &= \bar{\psi}_2(n) \\ a_2(m, 0) \Delta_1 [a_1(m, 0) \Delta_1 z(m, 0)] &= \bar{\psi}_1(m) \\ a_3(m, 0) \Delta_2 [a_2(m, 0) \Delta_1 [a_1(m, 0) \Delta_1 z(m, 0)]] &= \bar{\psi}_2(m). \quad \dots(54) \end{aligned}$$

Here  $a_1, a_2, a_3$  are as in Example 1,  $F: N_0^2 \times R \rightarrow R$ ,  $\bar{\phi}_1(n), \bar{\phi}_2(n), \bar{\psi}_1(m), \bar{\psi}_2(m)$  are real-valued nonnegative functions defined for  $m, n \in N_0$ . The equations equivalent to (45) – (46) and (53) – (54) are (48) and



$$z(m, n) = \bar{b}(m, n) + \sum_{x=0}^{m-1} \frac{1}{a_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{a_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{a_3(s, y)} \\ \times \sum_{t=0}^{y-1} F(s, t, z(s, t)) \quad \dots(55)$$

where  $\bar{b}(m, n)$  is obtained from the definition of  $b(m, n)$  by replacing  $\phi_1(n)$ ,  $\phi_2(n)$ ,  $\psi_1(m)$ ,  $\psi_2(m)$  in the right side in (49) by  $\bar{\phi}_1(n)$ ,  $\bar{\phi}_2(n)$ ,  $\bar{\psi}_1(m)$ ,  $\bar{\psi}_2(m)$  respectively. From (48) and (55) we have

$$u(m, n) - z(m, n) \\ = b(m, n) - \bar{b}(m, n) \\ + \sum_{x=0}^{m-1} \frac{1}{a_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{a_2(s, n)} \sum_{y=0}^{y-1} \frac{1}{a_3(s, y)} \\ \times \sum_{t=0}^{y-1} \{f(s, t, u(s, t)) - F(s, t, z(s, t))\}. \quad \dots(56)$$

Suppose that the function  $f$  in (45) satisfies the condition (51) and further we assume that

$$|b(m, n) - \bar{b}(m, n)| \leq \epsilon \quad \dots(57)$$

$$\sum_{x=0}^{m-1} \frac{1}{a_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{a_2(s, n)} \sum_{y=0}^{y-1} \frac{1}{a_3(s, y)} \sum_{t=0}^{y-1} |f(s, t, z(s, t)) \\ - F(s, t, z(s, t))| \leq \epsilon \quad \dots(58)$$

where  $\epsilon > 0$  is arbitrary constant. By subtracting and adding  $f(s, t, z(s, t))$  in the braces on the right side of equation (56) and using (51), (57), (58), we obtain

$$|u(m, n) - z(m, n)| \leq 2\epsilon + \sum_{x=0}^{m-1} \frac{1}{a_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{a_2(s, n)} \\ \sum_{y=0}^{n-1} \frac{1}{a_3(s, y)} \sum_{t=0}^{y-1} h(s, t) |u(s, t) - z(s, t)|. \quad \dots(59)$$

Now an application of Theorem 1 yields

$$|u(m, n) - z(m, n)|$$

$$\leq 2\epsilon \left\{ \prod_{x=0}^{m-1} \left[ 1 + \frac{1}{a_1(x, n)} \sum_{s=0}^{x-1} \frac{1}{a_2(s, n)} \sum_{y=0}^{n-1} \frac{1}{a_3(s, y)} \right. \right. \\ \left. \left. \sum_{t=0}^{y-1} h(s, t) \right] \right\}. \quad \dots(60)$$

If  $h(m, n)$  is bounded on some compact set  $0 \leq m \leq m_0$ ,  $0 \leq n \leq n_0$ ,  $m, m_0, n, n_0 \in N_0$ , then the quantity in braces on the right in (60) is bounded by some constant  $M$  on the set  $0 \leq m \leq m_0$ ,  $0 \leq n \leq n_0$ . Therefore  $|u(m, n) - z(m, n)| \leq 2M\epsilon$  on the set  $0 \leq m \leq m_0$ ,  $0 \leq n \leq n_0$ ; so the solution  $u(m, n)$  of (45) – (46) depends continuously on  $f$  and the boundary data. If  $\epsilon \rightarrow 0$ , then  $|u(m, n) - z(m, n)| \rightarrow 0$  on this set.

#### REFERENCES

1. Y. V. Bykov and V. G. Linenko, *Differentsial 'nye Uravenija* 9 (1973), 349-54.
2. V. B. Demidovich, *Differentsial 'nye Uravenija* 5 (1969), 1247-55.
3. P. Henrici, *Discrete Variable Methods in Ordinary Differential Equations*, Wiley, New York, 1962.
4. L. V. Masolockaja, *Differentsial 'nye Uravenija* 34 (1967), 1976-83.
5. B. G. Pachpatte, *Mich. Math. J.* 18 (1971), 385-91.
6. B. G. Pachpatte, *Univ. Beograd Publ. Elek. Fak. Ser. Mat. Fiz.* No. 577-No. 598 (1977), 65-73.
7. B. G. Pachpatte, *Bull. Inst. Math. Acad. Sinica* 5 (1977), 121-28.
8. B. G. Pachpatte, *Tamkang J. Math.* 12 (1981), 21-33.
9. S. Sugiyama, *Bull. Sci. Engr. Research Lab. Waseda Univ.* 45 (1969), 140-44.
10. D. Wiltent and J. S. W. Wong, *Monatsh. Math.* 69 (1964), 362-67.

## PERIODIC BOUNDARY VALUE PROBLEMS FOR AN INFINITE SYSTEM OF NONLINEAR SECOND ORDER DIFFERENTIAL EQUATIONS

K. NARSIMHA REDDY

*Department of Mathematics, P. G. College, Osmania University, Secunderabad 500003*

*(Received 3 August 1989; after revision 7 April 1989; accepted 8 May 1989)*

Results on existence of solutions to Periodic Boundary Value Problems for an infinite System of nonlinear second order differential equations have been discussed and a uniqueness result is presented.

### 1. INTRODUCTION

There is a large literature on the existence of solutions for periodic boundary value problems (PBVP's for short) of nonlinear scalar differential equations<sup>7-10,12,14</sup>. Results on the existence of solutions for first and second order PBVP's have been obtained earlier<sup>7,8,10,14</sup> by combining two basic techniques, namely the method of upper and lower solutions and the alternative method. In deed, a number of existence theorems for periodic solutions have been obtained by the technique of upper and lower solutions, and others have been and are being obtained by the alternative method. Kannan and Lakshmikantham<sup>6,7</sup> showed that the use of a result proved by Cesari and Kannan<sup>3</sup> by the alternative method could remarkably improve the arguments by upper and lower solutions. New existence theorems could be obtained and previous ones could be obtained by a more uniform approach. Further work in this direction was done by Rao and Vatsala<sup>15</sup>. For an exposition on the alternative method, though preceeding the work in Cesari and Kannan<sup>3</sup>, we refer to Cesari<sup>2</sup>. Also we refer the readers to Bernfeld and Lakshnikantham<sup>1</sup> for a general treatment of boundary value problems. Extremal solutions for nonlinear boundary value problems have been obtained by many workers<sup>4,5,8,9,11,16,17</sup> by employing monotone iterative scheme and this method is found to be constructive. PBVP's for infinite systems of first and second order equations have been studied earlier<sup>8,11,14,15,17</sup> and these results extend the results obtained for scalar equations.

In this paper we consider, as it was done earlier<sup>4,5,7,16</sup>, an infinite system of second order ordinary differential equations with periodic boundary conditions and discuss the questions of existence and uniqueness of solutions again by combining the method of upper and lower solutions with the result in Cesari and Kannan<sup>3</sup> from the alternative method. The organization of the present paper is as follows : Our notations and terminology are fairly consistent and can be understood by referring to earlier workers<sup>5,11,15,16</sup>. However for the sake of completeness we describe them briefly in section 2. Section 3 deals with existence results employing the alternative method. In

section 4, we develop a monotone iterative technique to obtain coupled extremal quasi-solutions for these systems. The uniqueness results are discussed in section 5.

## 2. NOTATIONS AND TERMINOLOGY

Let  $I$  be the interval  $[0, 2\pi]$  and  $E = R^\infty = R \times R \times \dots$  in which  $R = (-\infty, \infty)$ . Also let  $Z^+$  denote the set of all positive integers.

We consider the PBVP

$$-u_i'' = f_i(t, u, u_i'), \quad t \in I, i \in Z^+ \quad \dots(2.1)$$

$$u(0) = u(2\pi), u'(0) = u'(2\pi)$$

where  $f: I \times E \times R \rightarrow E$  is continuous.

For each  $i \in Z^+$ , we define two sets  $P_i$  and  $Q_i$  such that  $Z^+ = \{i\} \cup P_i \cup Q_i$ . Hereafter  $P(q)$  denotes the generic element of  $P_i(Q_i)$  respectively whenever the set is nonempty. Moreover the vector  $u \in E$  may be written as  $u = (u_i, [u]_p, [u]_q)$ , for  $p \in P_i, q \in Q_i$ . Then the PBVP (2.1) becomes

$$\left. \begin{aligned} -u_i'' &= f_i(t, u_i, [u]_p, [u]_q, u_i') \\ u(0) &= u(2\pi), u'(0) = u'(2\pi). \end{aligned} \right\} \quad \dots(2.2)$$

Without further mention we assume that  $i \in Z^+$  and all the inequalities between vectors hold component wise.

We state the following assumptions which will be used in our subsequent discussion :

$$(A_0) \quad \alpha, \beta \in C^2[I, E], \quad \alpha(t) \leq \beta(t), \quad t \in I$$

$$(A_1) \quad (i) \quad \alpha(0) = \alpha(2\pi), \alpha'(0) \geq \alpha'(2\pi) \text{ and } -\alpha_i'' \leq f_i(t, \sigma, \alpha_i')$$

for all  $\sigma$  such that  $\alpha(t) \leq \sigma \leq \beta(t)$  and  $\sigma_i = \alpha_i(t), t \in (0, 2\pi]$

$$(ii) \quad \beta(0) = \beta(2\pi), \beta'(0) \leq \beta'(2\pi) \text{ and } -\beta_i'' \geq f_i(t, \sigma, \beta_i')$$

for all  $\sigma$  such that  $\alpha(t) \leq \sigma \leq \beta(t)$  and  $\sigma_i = \beta_i(t), t \in (0, 2\pi]$

$$(A_2) \quad \text{For } t \in I, \alpha(t) \leq u(t) \leq \beta(t) \text{ and } v_i \in R \text{ we have}$$

$$|f_i(t, u, v_i)| \leq \begin{cases} h_i(|v_i|) & \text{if } |v_i| \leq d_i \\ h_i(d_i) & \text{if } |v_i| > d_i \end{cases}$$

for some  $d_i > e_i = 1/2\pi \max \{ \alpha_i(0) - \beta_i(2\pi), |\alpha_i(2\pi) - \beta_i(0)| \}$ ,  $h_i: [0, \infty) \rightarrow (0, \infty)$  is continuous for each  $i$ .



Also there exists  $N > 0$  depending only on  $\alpha$ ,  $\beta$  and  $h$  such that

$$\int_{e_i}^{l_i} \frac{s \, ds}{h_i(s)} > \max_I \beta_i(t) - \min_I \alpha_i(t).$$

where

$$l_i = \min \{d_i, N_i\}$$

(A<sub>3</sub>)  $f$  is completely continuous on  $I \times E \times R$ .

(A<sub>4</sub>) For  $t \in I$ ,  $\alpha(t) \leq v \leq u \leq \beta(t)$  and  $|z_i| \leq \bar{d}_i$

$$f_i(t, u_i, [u]_p, [u]_q, z_i) - f_i(t, v_i, [u]_p, [u]_q, z_i) \geq -M_i(u_i - v_i)$$

for some  $M_i > 0$ . Where  $\bar{d}_i = \max \left\{ N_i, \max_I |\alpha'_i(t)|, \max_I |\beta'_i(t)| \right\}$

(A<sub>5</sub>)  $f$  possess a mixed quasi-monotone property (*mq mp*) that is  $f_i(t, u_i, [u]_p, [u]_q, z_i)$  is monotone nondecreasing in  $[u]_p$  and monotone nonincreasing in  $[u]_q$ .

The functions  $\alpha, \beta \in C^2[I, E]$  with  $\alpha(t) \leq \beta(t)$  on  $I$  are said to be coupled lower and upper quasi-solutions of (2.1) respectively if

$$-\alpha''_i \leq f_i \left( t, \alpha_i, [\alpha]_p, [\beta]_q, \alpha'_i \right), \alpha(0) = \alpha(2\pi), \alpha'(0) \geq \alpha'(2\pi)$$

$$-\beta''_i \geq f_i \left( t, \beta_i, [\beta]_p, [\alpha]_q, \beta'_i \right), \beta(0) = \beta(2\pi), \beta'(0) \leq \beta'(2\pi).$$

The functions  $x, y \in C^2[I, E]$  are said to be coupled quasi-solutions of (2.1) if

$$-x''_i = f_i \left( t, x_i, [x]_p, [y]_q, x'_i \right), x(0) = x(2\pi), x'(0) = x'(2\pi)$$

$$-y''_i = f_i \left( t, y_i, [y]_p, [x]_q, y'_i \right), y(0) = y(2\pi), y'(0) = y'(2\pi).$$

In the special case where all  $Q_i$ 's are empty, quasi-solutions are just solutions and in the case  $P_i$ 's are empty, the quasi-solutions that result are most useful since they may be obtained most easily. We can also define coupled minimal and maximal quasi-solutions analogously.

We state the lemma which is a modified version of a known result.

**Lemma 2.1**—Let the assumptions (A<sub>0</sub>) and (A<sub>2</sub>) hold, then for any solution  $u \in C^2[I, E]$  of (2.1) with  $\alpha(t) \leq u(t) \leq \beta(t)$  on  $I$ , we have

$$|u'(t)| \leq N \text{ on } I.$$

The proof of this lemma follows from Lemma 1.1 of Das and Devasahayam.

## 3. EXISTENCE RESULTS

In this section we discuss the existence of solutions of PBVP's by the device, already used many times in the method of upper and lower solutions, of defining the function  $F$  as follows :

$$F_t(t, u, v_t) = f_t(t, p(t, u), v_t) + r_t(t, u)$$

where

$$p_t(t, u) = \max \{ \alpha_t(t), \min(u_t, \beta_t(t)) \}$$

and

$$r_t(t, u) = \begin{cases} \frac{\beta_t(t) - u_t}{1 + u_t^2}, & \text{if } u_t > \beta_t(t) \\ 0, & \text{if } \alpha_t(t) \leq u_t \leq \beta_t(t) \\ \frac{\alpha_t(t) - u_t}{1 + u_t^2}, & \text{if } u_t < \alpha_t(t). \end{cases}$$

Consider the following PBVP

$$-u'' = F_t\left(t, u, u'_t\right), u(0) = u(2\pi), u'(0) = u'(2\pi). \quad \dots(3.1)$$

*Lemma 3.1*—Let  $(A_0)$  and  $(A_1)$  hold and let  $u$  be a solution of (3.1). Then

$$\alpha(t) \leq u(t) \leq \beta(t) \text{ on } I.$$

**PROOF :** First we claim that  $\alpha(t) \leq u(t)$  on  $I$ . Suppose not, then we can find a  $t_0 \in I$  and an  $\epsilon > 0$  such that for some  $k \in Z^+$

$$\alpha_k(t_0) = u_k(t_0) + \epsilon, \alpha_t(t) \leq u_t(t) + \epsilon \quad t \in I. \quad \dots(3.2)$$

If  $t_0 \in (0, 2\pi)$ , we have  $\alpha'_k(t_0) = u'_k(t_0)$  and  $\alpha''_k(t_0) \leq u''_k(t_0)$ . From (3.2)  $\alpha_k(t_0) > u_k(t_0)$  and hence  $p_k(t_0, u(t_0)) = \alpha_k(t_0)$ . In view of  $(A_1)$  (i) and using the definition of  $F$  we have

$$\begin{aligned} f_k(t_0, \sigma, \alpha'_k(t_0)) &\geq -\alpha''_k(t_0) \\ &\geq -u''_k(t_0) = F_k(t_0, u(t_0), u'_k(t_0)) \\ &> f_k(t_0, p(t_0, u(t_0)), \alpha'_k(t_0)). \end{aligned}$$

Since  $\alpha(t) \leq p(t, u(t)) \leq \beta(t)$  and  $p_k(t_0, u(t_0)) = \alpha_k(t_0)$ , we get a contradiction by choosing  $\sigma = p(t_0, u(t_0))$ .

If  $t_0 = 0$ , from (3.2) we obtain

$$\alpha_k(0) = u_k(0) + \epsilon = \alpha_k(2\pi) \text{ and } \alpha'_k(0) \leq u'_k(0) \text{ and } \alpha'_k(2\pi) \geq u'_k(2\pi),$$

since  $\alpha_k(0) = \alpha_k(2\pi)$ ,  $u_k(0) = u_k(2\pi)$ .

And in view of  $(A_1)$  (i) it follows that  $\alpha'_k(2\pi) = u'_k(2\pi)$  and as before

$$f_k(2\pi, \sigma, \alpha'_k(2\pi)) = -\alpha''_k(2\pi) = -u''_k(2\pi) = F_k(2\pi, u(2\pi), u'_k(2\pi))$$

$$> f_k(2\pi, p(2\pi, u(2\pi)), \alpha'_k(2\pi)).$$

Since  $\alpha(2\pi) = p(2\pi, u(2\pi)) \leq \beta(2\pi)$  and  $p_k(2\pi, u(2\pi)) = \alpha_k(2\pi)$ , we again get a contradiction by choosing  $\sigma = p(2\pi, u(2\pi))$ . On similar lines we can prove that  $u(t) \leq \beta(t)$  on  $I$  and this completes the proof.

**Lemma 3.2**—Let the assumptions  $(A_0) - (A_2)$  hold. Then there exist  $\alpha_0, \beta_0$  such that the following are true :

$$\left( A_0^* \right) \alpha_0, \beta_0 \in C^2[I, E], \alpha_0(t) < \beta_0(t), t \in I$$

$$\left( A_1^* \right) \text{ (i) } \alpha_0(0) = \alpha_0(2\pi), \alpha'_0(0) \geq \alpha'_0(2\pi) \text{ and}$$

$$-\alpha''_{0,i} < F_i\left(t, \bar{\sigma}, \alpha'_{0,i}\right) \text{ for } \bar{\sigma} \text{ such that}$$

$$\alpha_0(t) \leq \bar{\sigma} \leq \beta_0(t) \text{ and } \bar{\sigma}_t = \alpha_{0,t}(t), t \in (0, 2\pi].$$

$$\text{(ii) } \beta_0(0) = \beta_0(2\pi), \beta'_0(0) \leq \beta'_0(2\pi) \text{ and}$$

$$-\beta''_{0,i} > F_i\left(t, \bar{\sigma}, \beta'_{0,i}\right) \text{ for all } \bar{\sigma} \text{ such that}$$

$$\alpha_0(t) \leq \bar{\sigma} \leq \beta_0(t) \text{ and } \bar{\sigma}_t = \beta_{0,t}(t), t \in (0, 2\pi].$$

$\left( A_2^* \right)$  The condition  $(A_2)$  holds with  $F$  replacing  $f$  and with respect to the pair  $(\alpha_0, \beta_0)$ .

**PROOF :** Let  $a_i > 0$ ,  $b_i > 0$  for all  $i \in Z^+$  be numbers and define  $\alpha_{0,i}(t) = \alpha_i(t) - a_i$  and  $\beta_{0,i}(t) = \beta_i(t) + b_i$ . Then it is easy to see following the proof of Lemma 2.2 Rao and Vatsala<sup>15</sup> that  $\left( A_0^* \right)$  and  $\left( A_1^* \right)$  hold. However for the sake of completeness we establish  $\left( A_2^* \right)$ .

For  $t \in I$ ,  $\alpha(t) \leq u(t) \leq \beta(t)$  and  $u'_i \in R$ , we have

$$|F_i(t, u, u'_i)| = |f_i(t, u, u'_i)| = \begin{cases} h_i(|u'_i|), & \text{if } |u'_i| \leq d_i \\ h_i(d_i), & \text{if } |u'_i| > d_i. \end{cases}$$

Since  $h_i(s)$  is a positive constant for  $s \geq d_i$ , there exists an  $N_i^* > d_i$  such that

$$\int_{e_i^*}^{N_i^*} \frac{s \, ds}{h_i(s)} - \int_{e_i^*}^{N_i^*} \frac{s \, ds}{h_i(s)} > \max_I \beta_{0,i}(t) - \min_I \alpha_{0,i}(t)$$

where

$$e_i^* = \min\{e_i, \bar{e}_i\} \text{ and } \bar{e}_i = \frac{1}{2\pi} \max[|\alpha_{0,i}(0) - \beta_{0,i}(2\pi)|, |\alpha_{0,i}(2\pi) - \beta_{0,i}(0)|].$$

This proves that  $(A_2)$  holds with  $F_i$  replacing  $f_i$ ,  $i \in Z^+$ .

For  $\sigma, \bar{\sigma}$  such that  $\alpha_{0,i}(t) \leq \sigma$ ,  $\bar{\sigma} \leq \beta_{0,i}(t)$ ,  $\sigma_i = \alpha_{0,i}$  and  $\bar{\sigma}_i = \beta_{0,i}$ , define

$$G_i(t, u) = \begin{cases} F_i(t, \bar{\sigma}, \beta'_{0,i}) + \frac{\beta_{0,i} - u}{1 + u_i^2} & \text{if } u > \beta_{0,i} \\ \frac{u - \alpha_{0,i}}{\beta_{0,i} - \alpha_{0,i}} \left[ F_i(t, \bar{\sigma}, \beta'_{0,i}) - F_i(t, \sigma, \alpha'_{0,i}) \right] \\ \quad + F_i(t, \sigma, \alpha'_{0,i}) & \text{if } \alpha_{0,i} \leq u \leq \beta_{0,i} \\ F_i(t, \sigma, \alpha'_{0,i}) + \frac{\alpha_{0,i} - u}{1 + u_i^2} & \text{if } u < \alpha_{0,i}. \end{cases}$$

Since  $\alpha_{0,i} < \beta_{0,i}$  for all  $t \in I$ ,  $G_i(t, u)$  is well defined. We now consider the modified PBVP

$$-u''_i = G_i(t, u), \quad u(0) = u(2\pi), \quad u'(0) = u'(2\pi). \quad \dots(3.3)$$

**Lemma 3.3**—Assume that  $(A_0)$ ,  $(A_1)$  and  $(A_3)$  hold. Then the problem (3.3) has a unique solution  $u$  satisfying

$$\alpha_{0,i}(t) \leq u_i(t) \leq \beta_{0,i}(t), \quad t \in I.$$



PROOF : It is clear from the definition that  $G_t(t, u)$  is completely continuous and bounded on  $I \times E$ . Hence we can find a positive number  $J$  that depends on  $\alpha_0$  and  $\beta_0$  such that  $\|G(t, u)\| \leq J$ . Let  $X = L_2[0, 2\pi]$  define  $Lu = -u''$  and then  $D(L) = \{\varphi \in X: \varphi, \varphi' \text{ are real valued absolutely continuous on } [0, 2\pi], \varphi'' \in X, \varphi(0) = \varphi(2\pi) \text{ and } \varphi'(0) = \varphi'(2\pi)\}$ . Let  $\mathcal{N}$  be the nonlinear operator defined by

$$\mathcal{N}u = G(t, u).$$

Then the BVP (3.3) may be translated into the operator equation

$$Lu = \mathcal{N}u$$

Notice that  $X_0$ , the Kernel of  $L$  consists of all constant functions and hence  $X_t$  where  $X = X_0 \oplus X_t$  is the class of all vector functions whose average on  $[0, 2\pi]$  is zero. Also we can define the operators  $P$  and  $H$  satisfying the conditions of Theorem 2.1 of Kannan and Lakshmikantham<sup>7</sup>. Since  $G$  is bounded we can find constants  $A$  and  $B$  that depend only on  $\alpha_0, \beta_0$  such that any solution of (2.2) in Theorem 2.1 of Kannan and Lakshmikantham<sup>7</sup> satisfies  $\|u_1(t)\| \leq A$  and  $\|u'_1(t)\| \leq B$  for all  $t \in I$ .

Hence by Theorem 2.1 of Kannan and Lakshmikantham<sup>7</sup> it is enough to find an  $R_{0,i} > 0$  such that

$$\langle \mathcal{N}(u_{0,i} + u_{1,i}), u_{0,i} \rangle \geq 0 \text{ or } \leq 0 \quad \dots(2.4)$$

for all  $|u_{0,i}| = R_{0,i}$  and  $|u_{1,i}| \leq A, |u'_{1,i}| \leq B$  on  $I, i \in Z^+$ . Since  $X_0$  consists of all constants functions and  $u_0 \in X_0$ , (3.4) can be written as

$$\int_0^{2\pi} G_t(s, R_0 + u_1(s)) ds \leq 0$$

and

$$\int_0^{2\pi} G_t(s, -R_0 + u_1(s)) ds \geq 0$$

choose  $R_{0,i} > 0$  large enough so that  $R_{0,i} + u_{1,i} > \max_i \beta_{0,i}(t)$  and  $-R_{0,i} + u_{1,i}$

$< \min_i \alpha_{1,i}(t)$ , Using the definition of  $G_t$  and  $(A_1^*)$  we see that

$$\int_0^{2\pi} G_t(s, R_0 + u_1(s)) ds < \int_0^{2\pi} F_t(s, \bar{\sigma}, \beta'_{0,i}(s)) ds < 0$$

and

$$\int_0^{2\pi} G_t(s, -R_0 + u_1(s)) ds > \int_0^{2\pi} F_t(s, \sigma, \alpha'_{0,i}(s)) ds > 0$$

for any arbitrary but fixed  $\sigma, \bar{\sigma}$  satisfying

$$\alpha_{0,t}(t) \leq \sigma, \bar{\sigma} \leq \beta_{0,t}(t), \sigma_t = \alpha_{0,t}(t) \text{ and } \bar{\sigma}_t = \beta_{0,t}(t)$$

Hence by Theorem 2.1 of Kannan and Lakshmikantham<sup>7</sup>, there exists a solution  $u$  to the problem (3.3). Proceeding on the similar lines of that of Lemma 3.1, it is easy to see that  $u(t)$  satisfies the inequality

$\alpha_{0,t}(t) \leq u(t) \leq \beta_{0,t}(t), t \in I$ . From this it follows that for each  $i \in Z^+$ ,  $u_i$  is a solution of

$$\begin{aligned} -u_i'' &= \frac{u_i - \alpha_{0,t}}{\beta_{0,t} - \alpha_{0,t}} \left[ F_t \left( t, \bar{\sigma}, \beta_{0,t}' \right) - F_t \left( t, \sigma, \alpha_{0,t}' \right) \right] \\ &\quad + F_t \left( t, \sigma, \alpha_{0,t}' \right) \end{aligned} \quad \dots(3.5)$$

$$u(0) = u(2\pi), u'(0) = u'(2\pi)$$

where  $\alpha_{0,t}(t) \leq \sigma, \bar{\sigma} \leq \beta_{0,t}(t), \sigma_t = \alpha_{0,t}(t)$  and  $\bar{\sigma}_t = \beta_{0,t}(t)$ . Following the proof of Lemma 2.3 of Rao and Vatsala<sup>15</sup> it can be shown that the solution  $u(t)$  of (3.5) is unique.

We now prove our main result of this section.

**Theorem 3.1**—Let the assumptions  $(A_0) - (A_3)$  be satisfied. Then the PBVP (2.1) has a solution  $u$  such that  $\alpha(t) \leq u(t) \leq \beta(t)$  and  $|u(t)| \leq N$  on  $I$ , where  $N$  depends only on  $\alpha, \beta$  and the Nagumo function.

**PROOF :** Consider the boundary value problem

$$-u_i'' = H_t \left( t, u, u_i' \right), u(0) = u(2\pi), u'(0) = u'(2\pi) \quad \dots(3.6)$$

where

$$H_t \left( t, u, u_i' \right) = \lambda F_t \left( t, u, u_i' \right) + (1 - \lambda) G_t(t, u), \lambda \in [0, 1].$$

In view of Lemma 3.2, one can verify that  $\alpha_0$  and  $\beta_0$  satisfy  $\left( A_0^* \right), \left( A_1^* \right)$  and  $\left( A_2^* \right)$  with respect to  $H_t$ . Using the arguments similar to that of Lemma 3.1, it is easy to show that, if  $u_{\lambda,t}$  for  $\lambda \in (0, 1)$  is a solution of (3.6) then  $\alpha_{0,t}(t) \leq u_{\lambda,t}(t) \leq \beta_{0,t}(t), t \in I, i \in Z^+$ . Since  $H_t$  satisfies a Nagumo condition, we have  $|u_{\lambda,t}'(t)| = C, t \in I$  where the constant  $C$  is independent of  $\lambda$ . In view of Lemma 3.3, it follows that for  $\lambda \in [0, 1]$  all possible solutions of (3.6) satisfy  $\alpha_0(t) \leq u_{\lambda}(t) \leq \beta_0(t)$  and  $|u_{\lambda,t}'(t)| \leq C$  for all  $t \in I$ . Also for  $\lambda = 0$ , the problem (3.6) has a unique

solution. Thus one can choose a bounded, closed convex set  $B$  in the  $(u, u')$  space  $H^1 [0, 2\pi] = L_2 [0, 2\pi]$  such that (3.6) has no solution on the boundary of  $B$  for  $\lambda \in (0, 1)$  and it has a unique solution in the interior of  $B$  for  $\lambda = 0$ . Hence by Leray-Schauder theory, the problem (3.6) has a solution  $u$  for  $\lambda = 1$ . By Lemma 3.1 we have  $\alpha(t) \leq u(t) \leq \beta(t)$ ,  $t \in I$  and this  $u$  is a solution of (2.1) satisfying  $\alpha(t) \leq u(t) \leq \beta(t)$ ,  $t \in I$ . Thus  $(A_2)$  implies that  $|u'(t)| \leq N$ ,  $t \in I$  where  $N$  is the Nagume constant vector. This completes the proof.

As a special case of Theorem 3.1 we have the following result.

*Corollary 3.1*—Let the assumptions  $(A_0)$ ,  $(A_2)$ ,  $(A_3)$  and  $(A_5)$  hold. Further assume that there exist coupled lower and upper quasi-solutions  $\alpha$  and  $\beta$ . Then the conclusion of Theorem 3.1 is true.

*Remark 3.1*: Note that if  $\alpha, \beta$  are coupled lower and upper quasi-solutions, then  $(A_1)$  holds if  $f$  has  $mq$   $mp$  and also that  $(A_1)$  implies  $\alpha, \beta$  are coupled lower and upper quasi-solutions of (2.1).

Our results can also be extended to the following infinite system of second order equations with homogeneous Neumann boundary conditions (NBVP for short). That is

$$-u_i'' = f_i \left( t, u, u' \right), \quad u'(0) = u'(2\pi) = 0 \quad \dots(3.7)$$

where

$$f \in C[I \times E \times R, E] \text{ and } i \in Z^+.$$

*Theorem 3.2*—Let  $(A_0)$ ,  $(A_2)$  and  $(A_3)$  hold. Further let the following condition hold:

$$(B) \quad (i) \quad \alpha'(0) \geq 0, \alpha'(2\pi) \leq 0 \text{ and}$$

$$-\alpha_i''(0) \leq f_i \left( t, \sigma, \alpha'_i \right), t \in (0, 2\pi] \text{ for all } \sigma \text{ such that}$$

$$\alpha(t) \leq \sigma \leq \beta(t) \text{ and } \sigma_i = \alpha_i(t), t \in I, i \in Z^+$$

$$(ii) \quad \beta'(0) \leq 0, \beta'(2\pi) \geq 0 \text{ and}$$

$$-\beta_i'' \geq f_i \left( t, \sigma, \beta'_i \right), t \in (0, 2\pi] \text{ for all } \sigma$$

such that  $\alpha(t) \leq \sigma \leq \beta(t)$  and  $\sigma_i = \beta_i(t)$ ,  $t \in I, i \in Z^+$ .

Then the problem (3.7) has a solution  $u$  such that

$$\alpha(t) \leq u(t) \leq \beta(t) \text{ and } |u'(t)| \leq N \text{ on } I.$$

Where  $N$  depends only on  $\alpha, \beta$  and the Nagume function.

The proof is similar to the proof of Theorem (3.1) with appropriate modifications.

## 4. MONOTONE ITERATIVE METHOD

For any  $\eta, \mu \in C[I, E]$ ,  $\alpha(t) \leq \eta(t)$ ,  $\mu(t) \leq \beta(t)$ ,  $t \in I$ , we consider the quasilinear PBVP

$$-u_i'' = G_i(t, u, [u]_p, [u]_q, u_i'), u(0) = u(2\pi), u'(0) = u'(2\pi) \dots (4.1)$$

where

$$G_i(t, u, [u]_p, [u]_q, u_i') = f_i(t, \eta, [\eta]_p, [u]_q, g(u_i')) - M_i(u - \eta)$$

and

$$g(u_i') = \max \left[ -\bar{d}_i, \min(u_i', \bar{d}_i) \right].$$

Notice that  $G$  is defined on  $I \times [\alpha, \beta] \times R$  and  $(A_4)$  is equivalent to

$$\begin{aligned} \left( A_4^* \right) f_i \left( t, u, [u]_p, [u]_q, g(u_i') \right) - f_i \left( t, v, [v]_p, [u]_q, \right. \\ \left. g(u_i') \right) \geq -M_i(u - v) \end{aligned}$$

for  $M_i > 0$ ,  $t \in I$ ,  $\alpha(t) \leq v \leq u \leq \beta(t)$  and  $u_i' \in R$ .

Relative to the PBVP (4.1) we prove the following lemmas.

**Lemma 4.1**—Let the assumptions  $(A_0) - (A_2)$ ,  $(A_4)$  and  $(A_5)$  be satisfied. Then the assumptions  $(A_1)$  and  $(A_2)$  are true with respect to the PVP (4.1). That is  $\alpha, \beta$  are also coupled lower and upper quasi-solutions of the PBVP (4.1) and  $G_i$  satisfies the modified Nagumo condition relative to  $\alpha, \beta$ .

**PROOF :** Using the arguments of Lemma 3.1 of Lakshmikantham *et al.*<sup>11</sup> it is easy to show that  $\alpha, \beta$  are also coupled lower and upper quasi solutions of the PBVP (4.1). However when  $(A_2)$  holds we have

$$|f_i(t, u, [u]_p, [u]_q, u_i')| \leq \begin{cases} h_i(|u_i'|) & \text{if } |u_i'| \leq d_i \\ h_i(d_i) & \text{if } |u_i'| > d_i \end{cases} \dots (4.2)$$

for  $t \in I$  and some  $d_i > \epsilon_i = \frac{1}{2\pi} \max \{ |\alpha(0) - \beta(2\pi)|, |\alpha(2\pi) - \beta(0)| \}$

$\alpha(t) \leq u(t) \leq \beta(t)$ ,  $u_i' \in R$  and  $h_i \in C[[0, \infty), (0, \infty)]$ , also

$$\int_{\epsilon_i}^{l_i} \frac{s \, ds}{h_i(s)} > \max_I \beta(t) - \min_I \alpha(t)$$

where

$$l_i = \min \{d_i, N_i\}.$$

From this it follows that any solution  $u \in C^2[I, E]$  of (2.1) satisfies  $|u'_i| \leq N_i$ ,  $t \in I$ .

For  $t \in I$ ,  $\alpha_i(t) \leq u(t) \leq \beta_i(t)$ ,  $\alpha_i(t) \leq \eta_i$ ,  $\mu_i \leq \beta_i(t)$  and  $u'_i \in R$  we have

$$|M_i(u_i - \gamma_i)| \leq M_i \gamma_i$$

where

$$\gamma_i = \max_I \beta_i(t) - \min_I \alpha_i(t)$$

Further

$$|G_i(t, u_i[u]_p, [u]_q, u'_i)| = H_i(|u'_i|).$$

where

$$H_i(s) = \begin{cases} h_i(s) + M_i \gamma_i & \text{if } s \leq \bar{d}_i \\ h_i(\bar{d}_i) + M_i \gamma_i & \text{if } s > \bar{d}_i \end{cases}$$

Evidently since  $H_i(s)$  is a positive constant for  $s \geq \bar{d}_i$ , there exists an  $N_i^* > \bar{d}_i$  such that

$$\int_{e_i}^{N_i^*} \frac{s \, ds}{H_i(s)} \geq \int_{\bar{d}_i}^{N_i^*} \frac{s \, ds}{H_i(s)} > \gamma_i$$

and this proves that  $G_i$  also satisfies  $(A_2)$ . Hence the proof is complete.

We now prove a result on existence and uniqueness of solutions of the PBVP (4.1).

**Lemma 4.2**—Let the assumptions  $(A_0) - (A_5)$  hold. Then there exists a solution  $u$  of (4.1) such that  $\alpha_i(t) \leq u(t) \leq \beta_i(t)$  and  $|u'_i(t)| \leq \hat{N}_i$  on  $I$ . Further more the solution  $u(t)$  is unique.

**PROOF :** By Lemma (4.1) we have all the assumptions  $(A_0) - (A_2)$  satisfied with respect to the PBVP (4.1). Hence by Theorem (3.1), there exists a solution  $u(t)$  of (4.1) with  $\alpha_i(t) \leq u(t) \leq \beta_i(t)$  and  $|u'_i(t)| \leq \hat{N}_i$  on  $I$ . Using the arguments similar to that of Lemma 3.2 of Lakshmikantham *et al.*<sup>11</sup> we can show that the solution  $u(t)$  is unique.



Since for every  $\eta, \mu \in [\alpha, \beta]$ , the PBVP (4.1) has a unique solution  $u$ , we define the mapping  $A$  by

$$A(\eta, \mu) = u \quad \dots(4.3)$$

and study the properties of this mapping in the next lemma.

*Lemma 4.3*—Under the assumptions of Lemma 4.2, the mapping  $A$  defined by (4.3) satisfies the following properties.

- (i)  $\alpha \leq A(\alpha, \beta)$  and  $\beta \geq A(\beta, \alpha)$
- (ii) For  $\alpha \subseteq \eta \leq \mu \leq \beta$ ,  $A(\eta, \mu) \leq A(\mu, \eta)$ .

**PROOF :** We shall only prove that  $\beta \geq A(\beta, \alpha)$  since similar arguments prove that  $\alpha \leq A(\alpha, \beta)$ .

Let  $A(\beta, \alpha) = u$ , where  $u$  is the unique solution of the PBVP (4.1) with  $\eta = \beta$  and  $\mu = \alpha$ . Let  $\varphi(t) = u(t) - \beta(t)$ . Suppose that the inequality  $\varphi(t) \leq 0$ ,  $t \in I$  is false. Then there exists a  $t_0 \in I$  and an  $\epsilon > 0$  such that for some index  $k \in Z^+$ , we have

$$\varphi_k(t_0) = \epsilon \text{ and } \varphi_i(t) \leq \epsilon \text{ for all } t \in I \text{ and } i \in Z^+. \quad \dots(4.4)$$

If  $t_0 \in (0, 2\pi)$ , we have  $\varphi'_k(t_0) = 0$  and  $\varphi''_k(t_0) \leq 0$ .

Also

$$g(u'_k(t_0)) = g(\beta'_k(t_0)) = \beta'_k(t_0).$$

At  $t = t_0$  using  $(A_1)$  and (4.1), we have

$$\begin{aligned} 0 &\geq \varphi''_k(t_0) = u''_k(t_0) - \beta''_k(t_0) \\ &\geq -G_k(t_0, u_k(t_0), [u(t_0)]_p, [u(t_0)]_q, u_k^A(t_0)) \\ &\quad + f_k(t_0, \beta_k(t_0), [\beta(t_0)]_p, [\alpha(t_0)]_q, \beta_k'(t_0)) \\ &\geq M_k \epsilon > 0, \text{ a contradiction.} \end{aligned}$$

If  $t_0 = 0$ , then using the boundary conditions,  $(A_1)$  (ii) and (4.4) we obtain

$$\varphi_k(0) = u_k(0) - \beta_k(0) = u_k(2\pi) - \beta_k(2\pi) = \varphi_k(2\pi) = \epsilon$$

$$\varphi'_k(0) \leq 0 \text{ and } \varphi'_k(2\pi) \geq 0.$$

Also

$$\varphi'_k(0) = u'_k(0) - \beta'_k(0) \geq u'_k(2\pi) - \beta'_k(2\pi) = \varphi'_k(2\pi).$$

Hence  $\varphi'_k(2\pi) = 0$  and using  $(A_1)$  and (4.1) we get

$$\varphi''_k(2\pi) = u''_k(2\pi) - \beta''_k(2\pi) \geq M_k \epsilon > 0 \text{ which is again a contradiction.}$$

To prove (ii) let  $\eta, \gamma \in [\alpha, \beta]$  such that  $\eta \leq \mu$ . Let  $A(\eta, \mu) = x$ ,  $A(\mu, \eta) = y$  and  $\psi(t) = x(t) - y(t)$ . If the inequality  $\psi(t) \leq 0$  for  $t \in I$  is false, then there exist  $t_0 \in I$  and an  $\epsilon > 0$  such that for some  $k \in Z^+$ , we have

$$\psi_k(t_0) = \epsilon \text{ and } \psi_k(t) \leq \epsilon \text{ for } t \in I \text{ and } k \in Z^+. \quad \dots(4.5)$$

If  $t_0 \in (0, 2\pi)$ , we have

$$\psi'_k(t_0) = 0 \text{ and } \psi''_k(t_0) \leq 0.$$

From (4.5),  $(A_4^*)$  and  $(A_5)$  and in view of the definition of  $G$ ,

$$\begin{aligned} 0 &> \psi''_k(t_0) = x''_k(t_0) - y''_k(t_0) \\ &= -f_k(t_0, \eta_k, [\eta]_p, [\mu]_q, g(x'_k(t_0))) + M_k(x_k(t_0) - \eta_k(t_0)) \\ &\quad + f_k(t_0, \mu_k, [\mu]_p, [\eta]_q, g(y'_k(t_0))) - M_k(y_k(t_0) - \mu_k(t_0)) \\ &\geq M_k(x_k(t_0) - y_k(t_0)) > 0, \text{ a contradiction.} \end{aligned}$$

If  $t_0 = 0$ , then from (4.5) and the boundary conditions we obtain  $\psi_k(2\pi) = \epsilon$  and  $\psi'_k(2\pi) = 0$  and as before we get a contradiction at  $t = 2\pi$ . This completes the proof. The following is the main theorem of this section.

**Theorem 4.1**—Let the assumptions  $(A_0) - (A_5)$  be satisfied. Then there exist monotone sequences  $\{\alpha_n(t)\}$ ,  $\{\beta_n(t)\}$  with  $\alpha_0 = \alpha$ ,  $\beta_0 = \beta$  such that  $\alpha_n(t)$  and  $\beta_n(t)$  converge uniformly and monotonically to  $\rho(t)$  and  $r(t)$  respectively on  $I$ . Also  $(\rho, r)$  are coupled minimal and maximal quasi-solutions of the PBVP (2.1). More precisely, if  $(x, y)$  are any coupled quasi-solutions of (2.1) satisfying  $\alpha \leq x$ ,  $y \leq \beta$ , then

$$\begin{aligned} \alpha = \alpha_0 &\leq \alpha_1 \leq \alpha_2 \leq \dots \leq \alpha_n \leq \dots \leq \rho \leq x, y \leq r \leq \dots \leq \beta_n \\ &\leq \dots \leq \beta_1 \leq \beta_0 = \beta \text{ on } I. \end{aligned} \quad \dots(4.6)$$

Furthermore any other solution  $u$  of (2.1) satisfying  $\alpha(t) \leq u \leq \beta(t)$  on  $I$  also satisfies (4.6) on  $I$ .

**PROOF :** We know from Lemma 4.2 that for any  $\eta, \mu \in [\alpha, \beta]$ , the PBVP (4.1) has a unique solution  $u(t)$  such that  $\alpha(t) \leq u \leq \beta(t)$  and  $|u'_i(t)| \leq \hat{N}_i$  on  $I$ , where  $\hat{N}_i$  is the Nagumo constant relative to  $G$ . In view of Lemma 4.3 we may define

the sequences  $\alpha_n = A(\alpha_{n-1}, \beta_{n-1})$  and  $\beta_n = A(\beta_{n-1}, \alpha_{n-1})$ ,  $n = 1, 2, 3, \dots$  such that  $\alpha_0 = \alpha$  and  $\beta_0 = \beta$  and  $\alpha_n \leq \beta_n$  for each  $n$ . Since  $\alpha_0 \leq \alpha_1 \leq \beta_1 \leq \beta_0$ , by induction and the arguments similar to those used in Lemma 4.3, we can establish that  $\{\alpha_n\}$ ,  $\{\beta_n\}$  are monotone sequences such that

$$\alpha_0 \leq \alpha_1 \leq \dots \leq \alpha_n \leq \beta_n \leq \dots \leq \beta_2 \leq \beta_1 \leq \beta_0 \text{ on } I.$$

Where  $\alpha_n(t)$  and  $\beta_n(t)$  satisfy

$$\begin{aligned} -\alpha''_{n,i} &= f_i(t, \alpha_{n-1,t}, [\alpha_{n-1}]_p, [\beta_{n-1}]_q, g(\alpha'_{n,i})) \\ &- M_i(\alpha_{n,t} - \alpha_{n-1,t}, \alpha_n(0) = \alpha_n(2\pi), \alpha'_n(0) = \alpha'_n(2\pi) \end{aligned} \quad \dots(4.7)$$

$$\begin{aligned} -\beta''_{n,i} &= f_i(t, \beta_{n-1,t}, [\beta_{n-1}]_p, [\alpha_{n-1}]_q, g(\beta'_{n,i})) \\ &- M_i(\beta_{n,t} - \beta_{n-1,t}, \beta_n(0) = \beta_n(2\pi), \beta'_n(0) = \beta'_n(2\pi) \end{aligned} \quad \dots(4.8)$$

and

$$|\alpha'_n(t)|, |\beta'_n(t)| \leq N.$$

From (4.7),

$$\begin{aligned} -\alpha_{n,i}(t) &= \int_0^t \int_0^0 f_i(s, \alpha_{n-1,t}(s), [\alpha_{n-1}]_p, [\beta_{n-1}]_q, g(\alpha'_{n,i}(s))) ds d\sigma \\ &\int_0^t \int_0^\sigma M_i(\alpha_{n,t}(s) - \alpha_{n-1,t}(s)) ds d\sigma + C_{n,i}t + \lambda_{n,i}. \end{aligned}$$

Since for each  $i$ ,  $f_i$  is completely continuous, the sequence  $\{\alpha_n\}$  is uniformly bounded and equi-continuous. Thus  $\{\alpha_n\}$  contains a subsequence which is uniformly convergent by the Arzela-Ascoli theorem. In view of the fact that  $\{\alpha_n\}$  is monotone the full sequence converges uniformly on  $I$ . Further more the uniform boundedness of the sequence  $\{\alpha''_n\}$  implies that the sequence  $\{\alpha'_n\}$  is equi-continuous and uniformly bounded. Thus  $\{\alpha'_n\}$  contains a subsequence which is uniformly convergent on  $I$ . Similar reasoning for  $\{\beta_n\}$ . Thus we can find sub-sequences which we again denote by  $\{\alpha_n\}$ ,  $\{\beta_n\}$  converging uniformly and monotonically to  $\rho$ ,  $r$ . Where  $\rho(t) = \lim_{n \rightarrow \infty} \alpha_n(t)$  and  $r(t) = \lim_{n \rightarrow \infty} \beta_n(t)$ ,  $t \in I$ . And this implies that

$$\begin{aligned} -\rho''_i &= f_i(t, \rho_i, [\rho]_p, [r]_q, g(\rho'_i)), \rho(0) = \rho(2\pi), \rho'(0) = \rho'(2\pi) \\ -r''_i &= f_i(t, r_i, [r]_p, [\rho]_q, g(r'_i)), r(0) = r(2\pi), r'(0) = r'(2\pi) \end{aligned} \quad \dots(4.9)$$

Following a continuation argument similar to that of [Bernfeld and Lakshmikantham<sup>1</sup> p. 32], one can prove that  $\rho$  and  $r$  are actually coupled quasi-solutions of the PBVP (2.1). If  $(u, v)$  are any coupled solutions of (2.1) such  $u, v \in [\alpha, \beta]$   $t \in I$  and  $|u'|, |v'| \leq \hat{N} \leq \bar{d}$  on  $I$ , employing induction principle and the arguments similar to those used earlier, it can be shown that  $\alpha \leq u, v \leq \beta$  on  $I$ . Hence we have  $\rho \leq u, v \leq r$  on  $I$ , proving that  $\rho, r$  are coupled minimal and maximal quasi-solutions of the PBVP (2.1).

Since any solution  $u$  of (2.1) satisfying  $\alpha \leq u \leq \beta$  on  $I$  may be regarded as  $(u, u)$  coupled quasi-solution of (2.1), we also have  $\rho \leq u \leq r$  on  $I$ . This completes the proof of the theorem.

## 5. UNIQUENESS RESULT

We now present a result on the uniqueness of solutions of the PBVP (2.1).

**Theorem 5.1**—Assume  $(A_0) - (A_3)$ . In addition for each  $i \in Z^+$ , there exists a constant  $L_i > 0$  such that

$$(u_i - v_i) [f_i(t, u, u'_i) - f_i(t, v, v'_i)] \leq -L_i (u_i - v_i)^2 \quad \dots(5.1)$$

whenever  $\alpha(t) \leq u, v \leq \beta(t)$   $t \in I$  and  $u'_i - v'_i = 0$ . Then the PBVP (2.1) has a unique solution  $u(t)$  satisfying  $\alpha(t) \leq u(t) \leq \beta(t)$  on  $I$ .

**PROOF :** By Theorem 3.1 we know that the PBVP (2.1) has a solution. If possible, let  $u$  and  $v$  be two solutions for the PBVP (2.1) satisfying  $\alpha(t) \leq u(t), v(t) \leq \beta(t)$  for  $t \in I$ .

We define

$$\mu_i(t) = (u_i(t) - v_i(t))^2$$

and observe that  $\mu_i(0) = \mu_i(2\pi), \mu'_i(0) = \mu'_i(2\pi)$  for each  $i \in Z^+$

and

$$\begin{aligned} \mu''_i(t) &= -2(u_i(t) - v_i(t)) [f_i(t, u, u'_i) - f_i(t, v, v'_i)] \\ &\quad + 2(u'_i(t) - v'_i(t))^2 \end{aligned} \quad \dots(5.2)$$

We claim that  $\mu_i(t) \equiv 0$  for all  $i \in Z^+$  on  $I$ . If not there exists a  $t_0 \in I$  and an  $\epsilon > 0$  such that for some  $k \in Z^+$

$$\mu_k(t_0) = \epsilon \text{ and } \mu_i(t) \leq \epsilon \text{ for all } t \in I, i \in Z^+. \quad \dots(5.3)$$

If  $t_0 \in (0, 2\pi)$ , we have

$$\mu'_k(t_0) = 0 \text{ and } \mu''_k \leq 0.$$

Thus from (5.2) and (5.1) we have

$$\begin{aligned} 0 &\geq \mu_k(t_0) = -2(u_k(t_0) - v_k(t_0)) [f_k(t_0, u, u'_k(t_0)) \\ &\quad - f_k(t_0, v, v'_k(t_0))] + 2(u'_k(t_0) - v'_k(t_0))^2 \\ &\geq 2L_k \mu_k(t_0) > 0, \text{ a contradiction.} \end{aligned}$$

If  $t_0 = 0$ , we obtain  $\mu_k(0) = \epsilon = \mu_k(2\pi)$  and  $\mu'_k(0) \leq 0$  and  $\mu'_k(2\pi) \geq 0$ . However  $\mu'_k(0) = \mu'_k(2\pi)$  and hence  $\mu'_k(0) = \mu'_k(2\pi) = 0$  and consequently  $\mu''_k(\lambda) \leq 0$  for  $\lambda = 0, 2\pi$ .

Also

$$0 \geq \mu''_k(\lambda) = 2L_k \mu_k(\lambda) > 0, \text{ for } \lambda = 0, 2\pi$$

which is again a contradiction. Hence the proof is complete.

**Corollary 5.1**—Assuming the conditions of Theorem 4.1 and the hypothesis (5.1) one may conclude the existence of a unique solution for the PBVP (2.1). In this case, it suffices to show that the coupled minimal and maximal quasi-solutions  $p(t)$  and  $r(t)$  are identical. This can be accomplished by defining the function

$$\mu_i(t) = (p_i(t) - r_i(t))^2 \text{ for each } i \in Z^+ \text{ and } t \in I$$

and proceeding along the lines of the proof of Theorem 5.1.

#### ACKNOWLEDGEMENT

The author is very much thankful to Prof. V. Sree Hari Rao, Department of Mathematics, JNTU, Kukatpalli, Hyderabad, India for his helpful suggestions and encouragement.

#### REFERENCES

1. S. R. Bernfeld and V. Lakshmikantham, *An Introduction to Nonlinear Boundary Value Problems*. Academic Press, New York, 1974.
2. L. Cesari, Functional analysis, nonlinear differential equations, and the alternative method, *Proc. Conf. on Nonlinear Functional Analysis and Differential Equations*, Marcel Dekker Inc. New York, 1979, pp. 1-197.
3. L. Cesari and R. Kannan, *Proc. Am. Math. Soc.* **63** (1977), 221-25.
4. J. Chandra, V. Lakshmikantham and S. Leela, *Arch. Rat. Mech. Anal.* **69** (1978), 179-90.
5. K. M. Das and M. Paul Devasahayam, *Nonlinear Analysis, Theory, Methods, Applic.* **1** (1983), pp. 1051-60.
6. R. Kannan and V. Lakshmikantham, *Nonlinear Analysis* **6** (1982), 1-10.
7. R. Kannan and V. Lakshmikantham, *Applicable Analysis* **17** (1984), 103-13.



8. V. Lakshmikantham and A. S. Vatsala, *J. Math. Anal. Applic.* **79** (1981), 38-47.
9. V. Lakshmikantham and S. Leela, *J. Math. Anal. Applic.* **91** (1983), 1-7.
10. V. Lakshmikantham and S. Leela, *Nonlinear Analysis, Theory, Methods Applic.* **8** (1984), 281-87.
11. V. Lakshmikantham V. Sree Hari Rao and A. S. Vatsala, *Ap. Math. Comput.* **15** (1984), 71-83.
12. S. Leela, *Nonlinear Analysis, Theory, Methods, Applic.* **7** (1983), 349-55.
13. J. J. Nieto and V. Sree Hari Rao, *Acta Math. Hung.* **48** (1986), 59-66.
14. K. Schmidt, *J. Diff. Eqns.* **11** (1972) 180-92.
15. V. Sree Hari Rao and A. S. Vatsala, *Applicable Analysis* **17** (1983), 23-36.
16. R. C. Thompson, *Annales Polonici Mathematici* **XL1** (1983).
17. A. S. Vatsala, *Nonlinear Analysis, Theory, Methods Applic.* **7** (1983), 1283-89.

# ON THE SETS OF GENERALIZED HYPERGEOMETRIC FUNCTIONS AND THE REGGE, BARGMANN-SHELEPIN ARRAYS FOR THE 3-J AND 6-J COEFFICIENTS

K. SRINIVASA RAO AND V. RAJESWARI

*The Institute of Mathematical Sciences, Madras 600113*

(Received 7 March 1989; accepted 4 May 1989)

The connection between the 3- $j$  and the 6- $j$  coefficients to a set of six  ${}_3F_2(1)$   $s$  and a set of three (or, equivalently a set of four)  ${}_4F_3(1)$   $s$ , respectively, is used to obtain sets of Regge  $3 \times 3$  and Bargmann-Shelepin  $4 \times 3$  symbols. Closed form expressions are obtained for the polynomial zeros of degree  $n$  of these coefficients.

## INTRODUCTION

In literature, classical symmetries of the 3- $j$  and the 6- $j$  coefficients are 12 and 24 in number. By relating the 3- $j$  coefficient to a  $3 \times 3$  magic square symbol, Regge<sup>1</sup> showed that it has 72 symmetries. Bargmann<sup>2</sup> and Shelepin<sup>3</sup> related the 6- $j$  coefficient to a  $4 \times 3$  symbol, which exhibits all the 144 symmetries (including the classical symmetries) discovered by Regge<sup>4</sup>. We have shown that sets of  ${}_p+1F_p(1)$   $s$  are necessary and sufficient to account for all the known symmetries of the 3- $j$  and the 6- $j$  coefficients—explicitly, while a set of six  ${}_3F_2(1)$   $s$  represent the 3- $j$  coefficient<sup>5</sup>, either a set I of three or an equivalent set II of four  ${}_4F_3(1)$   $s$  represent the 6- $j$  coefficient<sup>6,7</sup>. Here we establish a connection between these sets of generalized hypergeometric functions and sets of Regge, Bargmann-Shelepin symbols for the 3- $j$  and the 6- $j$  coefficients, respectively. Using these, closed form expressions have been obtained for the polynomial zeros of degree  $n$  of the 3- $j$  and 6- $j$  coefficients, which have been the subject of detailed study—especially when  $n = 1$  or 2—by several authors in recent years.

## CLOSED FORM EXPRESSIONS

The 3- $j$  coefficient has been defined by Wigner<sup>8</sup> as :

$$\begin{aligned} \left( \begin{matrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{matrix} \right) &= \delta(m_1 + m_2 + m_3, 0) (-1)^{j_1 - j_2 - m_3} \Delta(j_1 j_2 j_3) \prod_{i=1}^3 \{(j_i \\ &+ m_i)! (j_i - m_i)!\}^{1/2} \times \sum_z (-1)^z \{z! \prod_{k=1}^2 \{z - \alpha_k\}! \\ &\prod_{l=1}^3 (\beta_l - z)!\}^{-1} \end{aligned} \quad \dots (1)$$

where

$$\max(\alpha_1, \alpha_2) \leq z \leq \min(\beta_1, \beta_2, \beta_3)$$

$$\alpha_1 = j_1 - j_3 + m_2, \alpha_2 = j_2 - j_3 - m_1$$

$$\beta_1 = j_1 - m_1, \beta_2 = j_2 + m_2, \beta_3 = j_1 + j_2 - j_3. \quad \dots(2)$$

and

$$\Delta(x, y, z) = [(-x + y + z)! (x - y + z)! (x + y - z)! / (x + y + z + 1)!]^{1/2} \quad \dots(3)$$

and  $\delta(x, y)$  is the Kronecker delta function. Regge<sup>1</sup> discovered new symmetries by associating the  $3-j$  coefficient with a magic  $3 \times 3$  square symbol :

$$\begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} = \begin{vmatrix} -j_1 + j_2 + j_3 & j_1 - j_2 + j_3 & j_1 + j_2 - j_3 \\ j_1 - m_1 & j_2 - m_2 & j_3 - m_3 \\ j_1 + m_1 & j_2 + m_2 & j_3 + m_3 \end{vmatrix} \\ = ||R_{ik}|| \quad \dots(4)$$

and asserted that the  $3 \times 3$  square symbol represents the invariance of the  $3-j$  coefficient to  $3!$  column and  $3!$  row permutations and to a reflection about its diagonal. Thus, Regge<sup>1</sup> established the existence of a 72-element symmetry group, comprising the well-known classical symmetries (which arise due to column permutations and to the space reflection :  $m_i \rightarrow -m_i$  arising due to the interchange of the second and third rows of (4)) and six new symmetries known as Regge symmetries of the  $3-j$  coefficient.

It has been shown by one of us (Srinivasa Rao<sup>5</sup>) that the  $3-j$  coefficient can be represented by a set of six  ${}_3F_2(1)$  s :

$$\begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} = \delta(m_1 + m_2 + m_3, 0) (-1)^{\sigma(pqr)} \prod_{i,k=1}^3 \{R_{ik}! / (J+1)!\}^{1/2} \\ \times [\Gamma(1-A, 1-B, 1-C, D, E)]^{-1} {}_3F_2(A, B, C; D, E; 1) \quad \dots(5)$$

where

$$A = -R_{2p}, B = -R_{3q}, C = -R_{1r}, D = R_{3r} - R_{2p} + 1,$$

$$E = R_{2r} - R_{3q} + 1, \Gamma(x, y, \dots) = \Gamma(x) \Gamma(y) \dots,$$

$$J = j_1 + j_2 + j_3 \quad \dots(6)$$

and

$$\sigma(pqr) = \begin{cases} R_{3p} - R_{2q} & \text{for even permutations} \\ R_{3p} - R_{2q} + J & \text{for odd permutations} \end{cases}$$

for all permutations of  $(p\ q\ r) = (123)$ . Since each one of the six  ${}_3F_2(1)$   $s$  represents only 12 symmetries arising from its invariance to  $3!$  numerator and  $2!$  denominator parameter permutations, the set of six  ${}_3F_2(1)$   $s$  is necessary and sufficient to account for all the 72 symmetries of the  $3-j$  coefficient.

Using the properties of the elements of the  $3 \times 3$  square symbol :

$$R_{lp} + R_{mp} = R_{nq} + R_{nr} \quad \dots(7)$$

for  $(lmn)$  and  $(pqr)$  being  $(123)$  cyclically, and the defining relations (6) for the numerator and denominator parameters, one can easily show that :

$$\|R_{lk}\| = \begin{vmatrix} -B + D + 1 & -A + E - 1 & -C \\ -A & -C + D - 1 & -B + E - 1 \\ -C + E - 1 & -B & -A + D - 1 \end{vmatrix}. \quad \dots(8)$$

From (8) it is straightforward to obtain the closed form expression :

$$\begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} = \begin{pmatrix} (-A - C + E - 1)/2 & (-B - C + D - 1)/2 \\ (A - C + E - 1)/2 & (-B + C - D + 1)/2 \\ (-A - B + D + E - 2)/2 \\ (-A + B + D - E)/2 \end{pmatrix} \quad \dots(9)$$

The parameters  $A$  and  $B$  are negative integers by definition. So, if we let  $P = -A$  and  $Q = -B$  and set  $C = -1$ , we get, for the polynomial zeros of degree one the expression :

$$\begin{pmatrix} j_1 & j_2 & j_3 \\ m_1 & m_2 & m_3 \end{pmatrix} = \begin{pmatrix} (P + E)/2 & (Q + D)/2 \\ (-P + E)/2 & (Q - D)/2 \\ (P + Q + D + E - 2)/2 \\ (P - Q + D - E)/2 \end{pmatrix} \quad \dots(10)$$

with the constraint equation :

$$PQ = DE \quad \dots(11)$$

which is a homogeneous multiplicative Diophantine equation (see, Bell<sup>9</sup>) of degree 2 whose complete solutions can be expressed in terms of four parameters. Identifying the four parameter solution to be :

$$P = ab, Q = cd, D = bd \text{ and } E = ac \quad \dots(12)$$

and substituting it in (10), we obtain the result given by Brudno<sup>10</sup>. The polynomial zeros of degree  $n$  arise due to the truncation of the  ${}_3F_2(1)$  series (5). By setting anyone of the numerator parameters, (say  $C$ ) to  $-n$  and equating the sum of the  $(n + 1)$  terms to zero, one obtains the constraint equation which must be satisfied by the

numerator and denominator parameters of the  ${}_3F_2(1)$  for realizing the polynomial zeros of degree  $n$ .

The  $6-j$  coefficient has been expressed by Regge<sup>4</sup> to be :

$$\left\{ \begin{matrix} a & b & e \\ d & c & f \end{matrix} \right\} = N \sum_p (-1)^p (p+1)! \left\{ \prod_{i=1}^4 (p - \alpha_i)! \prod_{j=1}^3 (\beta_j - p)! \right\}^{-1} \dots (13)$$

with

$$N = \Delta(a, b, e) \Delta(c, d, e) \Delta(a, c, f) \Delta(b, d, f)$$

$$\alpha_1 = a + b + e, \alpha_2 = c + d + e, \alpha_3 = a + c + f, \alpha_4 = b + d + f$$

$$\beta_1 = a + b + c + d, \beta_2 = a + d + e + f, \beta_3 = b + c + e + f$$

and  $\max(\alpha_1, \alpha_2, \alpha_3, \alpha_4) \leq p \leq \min(\beta_1, \beta_2, \beta_3)$ . The  $144$  symmetries of the  $6-j$  coefficient arise due to the invariance of (13) to the  $4!$  permutations of the  $\alpha$ 's and the  $3!$  permutations of the  $\beta$ 's. This is explicit in the notation of Bargmann<sup>2</sup> and Shelepin<sup>3</sup> wherein :

$$\left\{ \begin{matrix} a & b & e \\ d & c & f \end{matrix} \right\} = \left\| \begin{matrix} \beta_1 - \alpha_1 & \beta_2 - \alpha_1 & \beta_3 - \alpha_1 \\ \beta_1 - \alpha_2 & \beta_2 - \alpha_2 & \beta_3 - \alpha_2 \\ \beta_1 - \alpha_3 & \beta_2 - \alpha_3 & \beta_3 - \alpha_3 \\ \beta_1 - \alpha_4 & \beta_2 - \alpha_4 & \beta_3 - \alpha_4 \end{matrix} \right\| = \|R_{ik}\| \dots (14)$$

which is invariant to  $4!$  row permutations and  $3!$  column permutations. The elements of  $\|R_{ik}\|$  satisfy the 18 relations :

$$R_{kk} + R_{mn} = R_{kn} + R_{mk}$$

and

$$R_{4k} + R_{mn} = R_{4n} + R_{mk} \dots (15)$$

for  $k \neq m$  and  $k \neq n$  and  $k, m$  or  $n$  being 1, 2 or 3. Equivalently, every  $2 \times 2$  co-factor in (14), say  $\begin{vmatrix} \alpha & \beta \\ \gamma & \delta \end{vmatrix}$  satisfies the condition :  $\alpha + \delta = \beta + \gamma$ .

We<sup>6</sup> have shown that the  $6-j$  coefficient can be represented by a set  $I$  of three  ${}_4F_3(1)$ s as :

$$\left\{ \begin{matrix} a & b & e \\ d & c & f \end{matrix} \right\} = (-1)^{E+1} N \Gamma(1-E) \{ \Gamma(1-A, 1-B, 1-C, 1-D, F, G) \}^{-1} \times {}_4F_3(A, B, C, D; E, F, G; 1) \dots (16)$$

where

$$A = -R_{1p}, B = -R_{2p}, C = -R_{3p}, D = -R_{4p},$$

$$E = -R_{1p} - R_{2p} - R_{3p} - R_{4p} - 1, F = R_{3p} - R_{3p} + 1, G = R_{4p} - R_{4p} + 1 \dots (17)$$



for  $(p q r) = (123)$  cyclic; and use has been made of (15) in arriving at (17). It is now possible to express the standard Bargmann-Shelepin  $4 \times 3$  symbol in terms of the numerator and denominator parameters of the set  $I$  of  ${}_4F_3(1)$ s, using (15) again, as:

$$\left\{ \begin{matrix} a & b & e \\ d & c & f \end{matrix} \right\} = \left\| \begin{matrix} -A & F-A-1 & G-A-1 \\ -B & F-B-1 & G-B-1 \\ -C & F-C-1 & G-D-1 \\ -D & F-D-1 & G-D-1 \end{matrix} \right\| = \|R_{ik}\| \quad \dots(18)$$

where the negative denominator parameter  $E$  does not appear in (18) and the  $4 \times 3$  symbol in (18) exhibits only 48 symmetries which arise due to the invariance of the  $6-j$  coefficient to  $4!$  row permutations and  $2!$  column permutations in  $\|R_{ik}\|$ . The set  $I$  of three different  $4 \times 3$  symbols which exist due to the substitution  $(p q r) = (123)$  cyclically in (17) account for the 144 symmetries exhibited by (14). It is now straightforward to obtain the closed form expression for the  $6-j$  coefficient:

$$\left\{ \begin{matrix} a & b & e \\ d & c & f \end{matrix} \right\} = \left\{ \begin{matrix} (G-B-D-1)/2 & (F-B-C-1)/2 \\ (G-A-C-1)/2 & (F-A-D-1)/2 \\ (F+G-C-D-2)/2 \\ (F+G-A-B-2)/2 \end{matrix} \right\} \quad \dots(19)$$

Setting one of the numerator parameters of the  ${}_4F_3(1)$  to  $-1$  (say,  $D = -1$ ) and replacing the negative parameters  $A$ ,  $B$  and  $C$  by  $-v$ ,  $-w$ , and  $-u$ , respectively and letting  $F = y$  and  $G = x$ , we get for the r. h. s. of (19):

$$\left\{ \begin{matrix} (x+w)/2 & (y+u+w-1)/2 & (x+y+u-1)/2 \\ (x+u+v-1)/2 & (y+v)/2 & (x+y+v+w-2)/2 \end{matrix} \right\} \quad \dots(20)$$

which is a symmetry of the parametric solution of Brudno and Louck<sup>11</sup>, in the notation of Srinivasa Rao *et al.*<sup>12</sup> for the polynomial zeros of degree 1.

It was shown<sup>7</sup> that the  $6-j$  coefficient can also be represented by a set II of four  ${}_4F_3(1)$ s as:

$$\left\{ \begin{matrix} a & b & e \\ d & c & f \end{matrix} \right\} = (-1)^{4'-2} N \Gamma(A') [\Gamma(1-B', 1-C', 1-D', E', F', G)]^{-1} \\ \times {}_4F_3(A', B', C', D'; E', F', G'; 1). \quad \dots(21)$$

where, using (15), the numerator and denominator parameters can be shown to be:

$$A' = R_{q2} + R_{r1} + R_{s3} + 2, B' = -R_{p1}, C' = -R_{p2}, D' = -R_{p3} \\ E' = R_{q1} - R_{p1} + 1, F' = R_{r1} - R_{p1} + 1, G' = R_{s1} - R_{p1} + 1 \quad \dots(22)$$

for  $(p q r s) = (1234)$  cyclically. The  $4 \times 3$  symbol for this set II of  ${}_4F_3(1)$  s can be written as :

$$\left\{ \begin{matrix} a & b & e \\ d & c & f \end{matrix} \right\} \left\| \begin{matrix} -B' & -C' & -D' \\ E' - B' - 1 & E' - C' - 1 & E' - D' - 1 \\ F' - B' - 1 & F' - C' - 1 & F' - D' - 1 \\ G' - B' - 1 & G' - C' - 1 & G' - D' - 1 \end{matrix} \right\| = \|R'_{ik}\| \quad \dots(23)$$

where the positive numerator parameter  $A'$  does not appear in (23) and this  $4 \times 3$  symbol exhibits only 36 of the 144 symmetries of the  $6-j$  coefficient which arise due to its invariance to  $3!$  column (or all permutations of  $B', C', D'$ ) and  $3!$  row permutations (or all permutations of  $E', F', G'$ ) of  $\|R'_{ik}\|$ . The set II of four  $\|R'_{ik}\|$  s which arise due to the substitution  $(p q r s) = (1234)$  cyclically in (22) accounts for the 144 symmetries of the  $6-j$  coefficient. As in the case of the set I of  ${}_4F_3(1)$  s, in the case of this set II of  ${}_4F_3(1)$  s also we obtain a closed form expression :

$$\left\{ \begin{matrix} a & b & e \\ d & c & f \end{matrix} \right\} = \left\{ \begin{matrix} (E' + G' - B' - C' - 2)/2 \\ (F' - B' - C' - 1)/2 \end{matrix} \right. \\ \left. \begin{matrix} (E' + F' - B' - D' - 2)/2 & (F' + G' - C' - D' - 2)/2 \\ (G' - B' - D' - 1)/2 & (E' - C' - D' - 1)/2 \end{matrix} \right\} \quad \dots(24)$$

Setting one of the numerator parameters of the  ${}_4F_3(1)$  to  $-1$  (say,  $D' = -1$ ) and replacing the negative parameters  $B'$  and  $C'$  by  $-x$  and  $-y$ , respectively, and letting  $E', F'$ , and  $G'$  be  $v, u$  and  $w$ , we get for the r.h.s of (24):

$$\left\{ \begin{matrix} (x + y + v + w - 2)/2 & (x + u + v - 1)/2 & (y + u + w - 1)/2 \\ (x + y + u - 1)/2 & (x + w)/2 & (x + v)/2 \end{matrix} \right\} \quad \dots(25)$$

which is a symmetry of (20). We have shown elsewhere<sup>12</sup> that the polynomial zeros of degree one of the  $6-j$  coefficient are obtained when the parameters in (25) are subject to the condition :

$$x + y + z = u + v + w \quad \dots(26)$$

or equivalently,  $A B C = E F G$ ,  $D = -1$  in the case of (19) and  $A' B' C' = E' F' G'$ ,  $D' = -1$  in the case of (24) which is a multiplicative Diophantine equation of degree 3 subject to the constraint :

$$z = x + y + u + v + w. \quad \dots(27)$$

Obviously, polynomial zeros of degree  $n$  arise when the sum of the first  $n + 1$  terms of the  ${}_4F_3(1)$  occurring in (19) or (24) adds to zero. We could use either (19) or (24) to generate the complete set of zeros of degree  $n$ .

The polynomial zeros of degree 2 of the  $3-j$  and the  $6-j$  coefficients in terms of these closed form expressions have been studied by Louck<sup>13</sup> *et. al.* using their connection to Pell's equation. However their study does not lead to all the polynomial zeros of degree 2 of the  $6-j$  coefficient. Simple algorithms based on the principle of factorization of integers have been proposed by Srinivasa Rao and Chiu<sup>14</sup> to obtain all the polynomial zeros of degree 2 of the  $3-j$  and the  $6-j$  coefficients.

In conclusion, we have shown in this article the connection between sets of  ${}_{p+1}F_p(1)s$  and sets of Regge or Bargmann-Shelepin symbols for the  $3-j$  and the  $6-j$  coefficients. This led us to closed form expressions for the polynomial zeros of degree  $n$  of these coefficients. As  $n$  increases, the complexity of the constraint equation which has to be satisfied by the parameters in the closed form expressions-or, the numerator and denominator parameters of the  ${}_{p+1}F_p(1)s$ -increases. At present detailed studies have been made only of the polynomial zeros of degree 1 and 2 of the  $3-j$  and  $6-j$  coefficients.

#### ACKNOWLEDGEMENTS

One of us (K.S.R.) acknowledges with thanks the interesting discussions he had with Professor Charles B. Chiu on this topic and another (V.R) is thankful to the Council of Scientific and Industrial Research, Government of India for the award of a Senior Research Fellowship.

#### REFERENCES

1. T. Regge, *Nuovo Cim.* **10** (1958), 544.
2. V. Bargmann, *Rev. Mod. Phys.* **34** (1962), 829.
3. L. A. Shelepin, *Sov. Phys. JETP* **19** (1964), 707.
4. T. Regge, *Nuovo Cim.* **11** (1959), 116.
5. K. Srinivasa Rao, *J. Phys. A: Math. Gen.* **11** (1978), L69.
6. K. Srinivasa Rao, T. S. Santhanam and K. Venkatesh, *J. Math. Phys.* **16** (1975), 1528.
7. K. Srinivasa Rao and K. Venkatesh, *Group. Theor. Methods in Phys.* Proc. of V Int. Coll. Univ. of Montreal, Ed. by R. T. Sharp and B. Kolman, Academic Press, New York (1977), 649.
8. E. P. Wigner, reprinted in *Quantum Theory of Angular Momentum* (ed.: L. C. Biedenharn and H. Van Dam) Academic Press, New York 1965.
9. E. T. Bell, *Am. J. Math.* **55** (1933), 50; see also, K. Srinivasa Rao, *Proc. Ramanujan centennial Int. Conf.*, RMS Publication **1** (1988), 141.
10. S. Brudno, *J. Math. Phys.* **26** (1985), 434.
11. S. Brudno and J. D. Louck, *J. Math. Phys.* **26** (1985), 2092.
12. K. Srinivasa Rao, V. Rajeswari and R. C. King, *J. Phys. A: Math. Gen.* **21** (1988), 1959.
13. J. D. Louck and P. R. Stein, *J. Math. Phys.* **28** (1987), 2812.
14. K. Srinivasa Rao and C. B. Chiu, *J. Phys. A: Math. Gen.*

## EIGENVALUE APPROACH TO LINEAR MICROPOLAR ELASTICITY

R. K. MAHALANABIS AND J. MANNA

*Department of Mathematics, Jadavpur University, Calcutta 700032*

(Received 12 December 1988; accepted 17 July 1989)

In this paper the basic equations of linear micropolar elasticity in polar coordinates are arranged in the form of vector matrix differential equation in the Hankel transform domain. The problem is then converted to an algebraic eigenvalue problem and solved in the same domain. It is seen that the results obtained by eigenvalue approach are in full agreement with those of other researchers. Further it seems that this approach is more elegant and it is believed that this technique has not been applied earlier by any researcher to solve the fundamental equations of linear micropolar elasticity.

### INTRODUCTION

In recent years a detailed exposition of the linear theory of micropolar elasticity has been given by Kuvchinski and Aero<sup>10</sup>, Eringen and Suhubi<sup>9</sup>, Eringen<sup>8</sup>, Nowacki<sup>12</sup> etc.. Nowacki<sup>11</sup> has shown that the equations of motion of linear micropolar theory in polar coordinates can be decomposed into two mutually independent sets of three equations each in case of axisymmetric problems. Dhaliwal and Chowdhury<sup>5</sup> have solved the set (2.6) for the axisymmetric Reissner—Sagoci problem and the solution is obtained by the classical method. Dhaliwal<sup>7</sup> has solved the set (2.7) for the axisymmetric Baussinisq problem. Das *et al.*<sup>2,3</sup> have applied recently the eigenvalue approach in solving the basic equations of thermoelasticity and extended the approach to magneto-thermoelasticity. They have solved the basic equations by representing them to single vector matrix differential equation and converting finally to an algebraic eigenvalue problem.

In this paper we apply the technique of eigenvalue to solve the axisymmetric equations of linear micropolar elasticity. It is believed that none of the previous investigators have applied this approach in solving the problems of linear micropolar theory. Here the basic axisymmetric eqns. (2.6) and (2.7) are presented in terms of single vector-matrix differential equations in sections 3 and 4 respectively. These lead to eigenvalue problems (3.8) and (4.4) respectively for the sets (2.6) and (2.7) and these are solved for displacements in the Hankel transform domain. The characteristic equation of (2.6) gives repeated roots while the set (2.7) gives real distinct roots i.e. real distinct eigenvalues. The general solution for distinct roots is obtained by usual



procedure (see appendix A), while for repeated roots the solution is obtained following the procedure of Das *et al.*<sup>2</sup> (see Appendix A). Further, solution for the half-space is also obtained. It is also observed that the solution obtained by Dhaliwal<sup>6,7</sup> for sets (2.6) and (2.7) are in full agreement with those obtained by eigenvalue approach.

## 2. THE BASIC EQUATIONS

The equations of motion and other basic equations for a homogeneous isotropic centrosymmetric linear elastic body occupying a region  $R$  (vide Dhaliwal<sup>7</sup>), are given by

$$\left. \begin{aligned} \sigma_{ij,t} + \rho X_i &= \rho \ddot{u}_i \\ \mu_{ij,t} + \epsilon_{ijk} \sigma_{jk} + J_{,i} &= J \ddot{\omega}_i \end{aligned} \right\} \quad \dots(2.1)$$

and the kinematic relations are

$$\left. \begin{aligned} \beta_{ij} &= \omega_{j,i} \\ \gamma_{ij} &= u_{j,i} + \epsilon_{kji} \omega_k \end{aligned} \right\} \quad \dots(2.2)$$

the linear constitutive law being

$$\left. \begin{aligned} \sigma_{ij} &= \lambda \gamma_{kk} \delta_{ij} + 2\mu \gamma_{(ij)} + 2\alpha \gamma_{[ij]} \\ \mu_{ij} &= \beta \beta_{kk} \delta_{ij} + 2\gamma \beta_{(ij)} + 2\epsilon \beta_{[ij]} \end{aligned} \right\} \quad \dots(2.3)$$

where  $\sigma_{ij}$  are the stress tensor components;  $\mu_{ij}$  the couple stress tensor components;  $u_i$  the displacement field components;  $\omega_i$  the rotational field components;  $X_i$  the body force components;  $Y_i$  the body couple components;  $\gamma_{ij}$  the strain tensor components;  $\beta_{ij}$  the curvature twist tensor components;  $\epsilon_{ijk}$  the unit antisymmetric tensor;  $[ ]$  and  $( )$  indicate respectively the skew symmetric and symmetric part of a tensor;  $\lambda, \mu, \alpha, \beta, \gamma, \epsilon$ , are the elastic constants of the micropolar material;  $\rho$  is the density;  $J$  the rotational inertia; and the dot  $(\dot{\phantom{x}})$  denotes the derivatives with respect to time.

Substituting (2.2) and (2.3) in (2.1) a set of six differential equations are obtained and these equations are presented in the vector form as :

$$\left. \begin{aligned} \text{(i)} \quad & (\lambda + 2\mu) \nabla \nabla \cdot \mathbf{u} - (\mu + \alpha) \nabla \times \nabla \times \mathbf{u} + 2\alpha \nabla \times \boldsymbol{\omega} + \rho \mathbf{x} = \rho \ddot{\mathbf{u}} \\ \text{(ii)} \quad & (\beta + 2\gamma) \nabla \nabla \cdot \boldsymbol{\omega} - (\gamma + \epsilon) \nabla \times \nabla \times \boldsymbol{\omega} + 2\alpha \nabla \times \mathbf{u} - 4\alpha \boldsymbol{\omega} + J \mathbf{Y} = J \ddot{\boldsymbol{\omega}} \end{aligned} \right\} \quad \dots(2.4)$$

Here we observe that the material constant  $\alpha$  is responsible for a coupling of a displacement and micro rotation fields. Though these equations are coupled, they are independent in the case, when  $\alpha = 0$ . In this case eqn. (2.4); (i) reduces to displacement equations of motion of classical elasticity and eqn. (2.4) (ii) describes a hypothetical conditions

$$\boldsymbol{\omega} = \frac{1}{2} \nabla \times \mathbf{u}$$



is obtained.

Now to solve the static problem with no body forces, we take  $X = Y = \ddot{u} = \ddot{\omega} = 0$  and the cylindrical polar coordinates  $(r, \varphi, z)$  is introduced. Equations (2.4) now assumes the forms

$$\begin{aligned}
 & (\mu + \alpha) \left( \nabla^2 u_r - \frac{u_r}{r^2} - \frac{2}{r^2} \frac{\partial u_\varphi}{\partial \varphi} \right) + (\lambda + \mu - \alpha) \frac{\partial e}{\partial r} \\
 & \quad + 2\alpha \left[ \frac{1}{r} \frac{\partial \omega_z}{\partial \varphi} - \frac{\partial \omega_z}{\partial z} \right] = 0 \\
 & (\mu + \alpha) \left( \nabla^2 u_\varphi - \frac{u_\varphi}{r^2} + \frac{2}{r^2} \frac{\partial u_r}{\partial \varphi} \right) + (\lambda + \mu - \alpha) \frac{1}{r} \frac{\partial e}{\partial \varphi} \\
 & \quad + 2\alpha \left[ \frac{\partial \omega_r}{\partial z} - \frac{\partial \omega_z}{\partial r} \right] = 0 \\
 & (\mu + \alpha) \nabla^2 u_z + (\lambda + \mu - \alpha) \frac{\partial e}{\partial z} + 2\alpha \frac{1}{r} \left[ \frac{\partial}{\partial r} (r \omega_\varphi) - \frac{\partial \omega_r}{\partial \varphi} \right] \\
 & \quad = 0 \\
 & (\gamma + \epsilon) \left( \nabla^2 \omega_r - \frac{\omega_r}{r^2} - \frac{2}{r^2} \frac{\partial \omega_\varphi}{\partial \varphi} \right) - 4\alpha \omega_r + (\beta + \gamma - \epsilon) \\
 & \quad \times \frac{\partial \psi}{\partial r} + 2\alpha \left( \frac{1}{r} \frac{\partial u_z}{\partial \varphi} - \frac{\partial u_\varphi}{\partial z} \right) = 0 \\
 & (\gamma + \epsilon) \left( \nabla^2 \omega_\varphi - \frac{\omega_\varphi}{r^2} + \frac{2}{r^2} \frac{\partial \omega_r}{\partial \varphi} \right) - 4\alpha \omega_\varphi + (\beta + \gamma - \epsilon) \frac{\partial \chi}{\partial \varphi} \\
 & \quad + 2\alpha \left( \frac{\partial u_r}{\partial z} - \frac{\partial u_z}{\partial r} \right) = 0 \\
 & (\gamma + \epsilon) \nabla^2 \omega_z - 4\alpha \omega_z + (\beta + \gamma - \epsilon) \frac{\partial \psi}{\partial z} + 2\alpha \frac{1}{r} \left[ \frac{\partial}{\partial r} (r u_\varphi) \right. \\
 & \quad \left. - \frac{\partial u_r}{\partial \varphi} \right] = 0 \quad \dots (2.5)
 \end{aligned}$$

where

$$\begin{aligned}
 e &= \frac{1}{r} \frac{\partial}{\partial r} (r u_r) + \frac{1}{r} \frac{\partial u_\varphi}{\partial \varphi} + \frac{\partial u_z}{\partial z} \\
 \chi &= \frac{1}{r} \frac{\partial}{\partial r} (r \omega_r) + \frac{1}{r} \frac{\partial \omega_\varphi}{\partial \varphi} + \frac{\partial \omega_z}{\partial z} \quad \dots (2.6)
 \end{aligned}$$

The case in which the vectors of displacement  $\vec{u}$  and rotation  $\vec{\omega}$  depend only on the coordinates  $r, z$  and as such the equations (2.5) are decomposed into two mutually independent set of equations, viz.

$$\left. \begin{aligned} (a) \quad (\mu + \alpha) \left( \nabla^2 u_r - \frac{u_r}{r^2} \right) + (\lambda + \mu - \alpha) \frac{\partial e}{\partial r} - 2\alpha \frac{\partial \omega_\varphi}{\partial z} &= 0 \\ (b) \quad (\mu + \alpha) \nabla^2 u_z + (\lambda + \mu - \alpha) \frac{\partial e}{\partial z} + 2\alpha \frac{1}{r} \frac{\partial}{\partial r} (r \omega_\varphi) &= 0 \\ (c) \quad (\gamma + \epsilon) \left( \nabla^2 \omega_\varphi - \frac{\omega_\varphi}{r^2} \right) - 4\alpha \omega_\varphi + 2\alpha \left( \frac{\partial u_r}{\partial z} - \frac{\partial u_z}{\partial r} \right) &= 0 \end{aligned} \right\} \quad \dots (2.6)$$

$$\left. \begin{aligned} (\mu + \alpha) \left( \nabla^2 u_\varphi - \frac{u_\varphi}{r^2} \right) + 2\alpha \left( -\frac{\partial \omega_r}{\partial z} - \frac{\partial \omega_z}{\partial r} \right) &= 0 \\ (\gamma + \epsilon) \left( \nabla^2 \omega_r - \frac{\omega_r}{r^2} \right) - 4\alpha \omega_r + (\beta + \gamma - \epsilon) \frac{\partial \psi}{\partial r} \\ - 2\alpha \frac{\partial u_\varphi}{\partial z} &= 0 \quad (\gamma + \epsilon) \nabla^2 \omega_z - 4\alpha \omega_z + (\beta + \gamma - \epsilon) \frac{\partial \psi}{\partial r} \\ + 2\alpha \frac{1}{r} (r u_\varphi) &= 0 \end{aligned} \right\} \quad \dots (2.7)$$

where

$$\begin{aligned} e &= \frac{1}{r} \frac{\partial}{\partial r} (r u_r) + \frac{\partial u_z}{\partial z} \\ \chi &= \frac{1}{r} \frac{\partial}{\partial r} (r \omega_r) + \frac{\partial \omega_z}{\partial z} \\ \nabla^2 &\equiv \frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} + \frac{\partial^2}{\partial z^2} \end{aligned}$$

### 3. SOLUTION OF EQUATIONS (2.6)

Here only the set (2.6) are considered. The following state of force-stress  $\sigma$  and couple stress  $\mu$  are being ascribed to the displacement vector  $\mathbf{u} = (u_r, 0, u_z)$  and rotation vector  $\boldsymbol{\omega} = (0, \omega_\varphi, 0)$ :

$$\sigma = \begin{vmatrix} \sigma_{rr} & 0 & \sigma_{rz} \\ 0 & \sigma_{\varphi\varphi} & 0 \\ \sigma_{zr} & 0 & \sigma_{zz} \end{vmatrix}, \quad \mu = \begin{vmatrix} 0 & \mu_{r\varphi} & 0 \\ \mu_{\varphi r} & 0 & \mu_{\varphi z} \\ 3 & \mu_{z\varphi} & 0 \end{vmatrix} \quad \dots (3.1)$$

where the particular components of stress-tensor have the following forms after using the relation (2.3)

$$\sigma_{rr} = 2\mu \frac{\partial u_r}{\partial r} + \lambda e,$$

$$\sigma_{\varphi\varphi} = 2\mu \frac{u_r}{r} + \lambda e$$

$$\sigma_{zz} = 2\mu \frac{\partial u_z}{\partial z} + \lambda e$$

$$\begin{aligned}
 \sigma_{rz} &= \mu \left( \frac{\partial u_z}{\partial r} + \frac{\partial u_r}{\partial z} \right) - \alpha \left( \frac{\partial u_r}{\partial z} - \frac{\partial u_z}{\partial r} \right) + 2\alpha \omega_\varphi \\
 \sigma_{zr} &= \mu \left( \frac{\partial u_z}{\partial r} + \frac{\partial u_r}{\partial z} \right) + \alpha \left( \frac{\partial u_r}{\partial z} - \frac{\partial u_z}{\partial r} \right) - 2\alpha \omega_\varphi \\
 \mu_{r\varphi} &= \gamma \left( \frac{\partial \omega_\varphi}{\partial r} - \frac{\omega_\varphi}{r} \right) + \epsilon \left( \frac{\partial \omega_\varphi}{\partial r} + \frac{\omega_\varphi}{r} \right) \\
 \mu_{\varphi r} &= \gamma \left( \frac{\partial \omega_\varphi}{\partial r} - \frac{\omega_\varphi}{r} \right) - \epsilon \left( \frac{\partial \omega_\varphi}{\partial r} + \frac{\omega_\varphi}{r} \right) \\
 \mu_{\varphi z} &= (\gamma - \epsilon) \frac{\partial \omega_\varphi}{\partial z}, \quad \mu_{z\varphi} = (\gamma + \epsilon) \frac{\partial \omega_\varphi}{\partial z} \quad \dots(3.2)
 \end{aligned}$$

In the system of eqns. (2.6), three mutually independent functions  $u_r$ ,  $u_z$  and  $\omega_\varphi$  are involved. Multiplying (2.6) by  $J_0(\xi r)$  and (2.6a,c) by  $J_1(\xi r)$  and integrating between the limits 0 to  $\infty$ , we find that the system of partial differential equations (2.6) reduces to the following system of ordinary differential equations:

$$\left. \begin{aligned}
 [(\mu + \alpha) D^2 - (\lambda + 2\mu) \xi^2] \bar{u}_r - (\lambda + \mu - \alpha) \xi D \bar{u}_z - 2\alpha D \bar{\omega}_\varphi &= 0 \\
 (\lambda + \mu - \alpha) \xi D \bar{u}_r + [(\lambda + 2\mu) D^2 - (\mu + \alpha) \xi^2] \bar{u}_z + 2\alpha \xi^2 \bar{\omega}_\varphi &= 0 \\
 2\alpha D \bar{u}_r + 2\alpha \xi \bar{u}_z + [(\gamma + \epsilon) (D^2 - \xi^2) - 4\gamma] \bar{\omega}_\varphi &= 0
 \end{aligned} \right\} \quad \dots(3.3)$$

where  $\bar{u}_r$ ,  $\bar{u}_z$  and  $\bar{\omega}_\varphi$  are the Hankel transforms of the functions  $u_r$ ,  $u_z$  and  $\omega_\varphi$  respectively and are given by

$$(\bar{u}_r, \bar{\omega}_\varphi) = \int_0^\infty (u_r, \omega_\varphi) \xi J_1(\xi r) dr$$

$$\bar{u}_z = \int_0^\infty u_z \xi J_0(\xi r) dr$$

and

$$D \equiv \frac{d}{dz}, \quad D^2 \equiv \frac{d^2}{dz^2}.$$

In the matrix rotations eqns. (3.3) may be represented as

$$\begin{bmatrix} \mu + \alpha & 0 & 0 \\ 0 & \lambda + 2\mu & 0 \\ 0 & 0 & \gamma + \epsilon \end{bmatrix} D^2 \begin{bmatrix} \bar{u}_r \\ \bar{u}_z \\ \bar{\omega}_\varphi \end{bmatrix} - \begin{bmatrix} 0 & (\lambda + \mu - \alpha) \xi & 2\alpha \\ -(\lambda + \mu - \alpha) \xi & 0 & 0 \\ -2\alpha & 0 & 0 \end{bmatrix} \begin{bmatrix} \bar{u}_r \\ \bar{u}_z \\ \bar{\omega}_\varphi \end{bmatrix} = 0$$

(equation continued on p. 1242)

$$D \begin{bmatrix} \bar{u}_r \\ \bar{u}_z \\ \bar{\omega}_\varphi \end{bmatrix} - \begin{bmatrix} (\lambda + 2\mu) \xi^2 & 0 & 0 \\ 0 & (\mu + \alpha) \xi^2 & -2\alpha \xi \\ 0 & -2\alpha \xi & (\gamma + \epsilon) \xi^2 + 2\alpha \end{bmatrix} \begin{bmatrix} \bar{u}_r \\ \bar{u}_z \\ \bar{\omega}_\varphi \end{bmatrix} = 0 \quad \dots(3.4)$$

we write

$$M = \begin{bmatrix} \mu + \alpha & 0 & 0 \\ 0 & \lambda + 2\mu & 0 \\ 0 & 0 & \gamma + \epsilon \end{bmatrix}, N = \begin{bmatrix} 0 & (\lambda + \mu - \alpha) \xi & 2\alpha \\ -(\lambda + \mu - \alpha) \xi & 0 & 0 \\ -2\alpha & 0 & 0 \end{bmatrix}$$

$$X = \begin{bmatrix} \bar{u}_r \\ \bar{u}_z \\ \bar{\omega}_\varphi \end{bmatrix}, X' = D \begin{bmatrix} \bar{u}_r \\ \bar{u}_z \\ \bar{\omega}_\varphi \end{bmatrix}$$

$$P = \begin{bmatrix} (\lambda + 2\mu) \xi^2 & 0 & 0 \\ 0 & (\mu + \alpha) \xi^2 & -2\alpha \xi \\ 0 & -2\alpha \xi & (\gamma + \epsilon) \xi^2 + 4\alpha \end{bmatrix}, X'' = D^2 \begin{bmatrix} \bar{u}_r \\ \bar{u}_z \\ \bar{\omega}_\varphi \end{bmatrix}$$

$$B_1 = M^{-1} N$$

and

$$B_2 = M^{-1} P. \quad \dots(3.5)$$

Equation (3.4) with the aid of (3.5) reduces to the form

$$X'' = B_1 X' + B_2 X. \quad \dots(3.6)$$

Using block matrices eqn. (3.6) assumes the form

$$\frac{d}{dz} \begin{bmatrix} X' \\ X \end{bmatrix} = \begin{bmatrix} B_1 & B_2 \\ I & 0 \end{bmatrix} \begin{bmatrix} X' \\ X \end{bmatrix} \quad \dots(3.7)$$

where  $I$  is a  $3 \times 3$  unit matrix.

Writing

$$A = \begin{bmatrix} B_1 & B_2 \\ I & 0 \end{bmatrix} \text{ and } V = \begin{bmatrix} X' \\ X \end{bmatrix} \quad \dots(3.8)$$

eqn. (3.7) assumes the form

$$\frac{dV}{dz} = AV. \quad \dots(3.9)$$

Assume  $V = Y \exp(tz)$  to be a solution of eqn. (3.9). Then we must have

$$A Y = t Y. \quad \dots(3.10)$$

This gives that  $t$  is an eigenvalue of the matrix  $A$  and  $Y$  the corresponding eigenvector.

The eigenvalues of the matrix  $A$  are the roots of

$$\det (A - t I) = 0. \quad \dots(3.11)$$

Using simplified notations, eqn. (3.11) may be written explicitly as

$$\begin{vmatrix} -t & a_{12} & a_{13} & a_{14} & 0 & 0 \\ a_{21} & -t & 0 & 0 & a_{25} & a_{26} \\ a_{31} & 0 & -t & 0 & a_{35} & a_{36} \\ 1 & 0 & 0 & -t & 0 & 0 \\ 0 & 1 & 0 & 0 & -t & 0 \\ 0 & 0 & 1 & 0 & 0 & t \end{vmatrix} = 0 \quad \dots(3.12)$$

where

$$\begin{aligned} a_{12} &= \frac{\lambda + \mu - \alpha}{\mu + \alpha} \xi, \quad a_{13} = \frac{2 \alpha}{\mu + \alpha}, \quad a_{14} = \frac{\lambda + 2\mu}{\mu + \alpha} \xi^2 \\ a_{21} &= \frac{-\lambda + \mu - \alpha}{\lambda + 2\mu} \xi, \quad a_{25} = \frac{\mu + \alpha}{\lambda + 2\mu} \xi^2, \quad a_{26} = \frac{-2\alpha \xi}{\lambda + 2\mu} \\ a_{31} &= \frac{-2\alpha}{\gamma + \epsilon}, \quad a_{35} = \frac{-2\alpha \xi}{\gamma + \epsilon}, \quad a_{36} = \frac{(\gamma + \epsilon)\xi^2 + 4\alpha}{\gamma + \epsilon}. \end{aligned} \quad \dots(3.13)$$

Simplifying (3.12) and using (3.13) therein we get the characteristic equation as

$$t^6 - (2\xi^2 + \zeta^2) t^4 + (\xi^4 + 2\xi^2 \zeta^2) t^2 - \xi^4 \zeta^2 = 0 \quad \dots(3.14)$$

where

$$\zeta^2 = \xi^2 + m^2$$

and

$$m^2 = \frac{4 \alpha \mu}{(\mu + \alpha) (\gamma + \epsilon)}.$$

The roots of (3.14) are

$$\xi, \xi, -\xi, -\xi, \zeta, -\zeta.$$

The eigenvalues of the matrix  $A$  are the root of the equation (3.14). Write  $t_1 = \xi$ ,  $t_2 = -\xi$ ,  $t_3 = \zeta$ , and  $t_4 = -\zeta$ . The four eigenvectors corresponding to four distinct eigenvalues  $t_1, t_2, t_3, t_4$  of the matrix  $A$  are obtained by solving the following homogeneous equations



$$\begin{bmatrix} -t & a_{12} & a_{13} & a_{14} & 0 & 0 \\ a_{21} & -t & 0 & 0 & a_{25} & a_{26} \\ a_{31} & 0 & -t & 0 & a_{35} & a_{36} \\ 1 & 0 & 0 & -t & 0 & 0 \\ 0 & 1 & 0 & 0 & -t & 0 \\ 0 & 0 & 1 & 0 & 0 & -t \end{bmatrix} \begin{bmatrix} y_1(t) \\ y_2(t) \\ y_3(t) \\ y_4(t) \\ y_5(t) \\ y_6(t) \end{bmatrix} = 0 \quad \dots(3.15)$$

for  $t = t_1, t_2, t_3, t_4$ .

Denote by  $A_1(t), A_2(t), \dots, A_6(t)$  the co-factors of the elements of the first row of the coefficient matrix in eqn. (3.15), then

$$Y(t) = \begin{bmatrix} A_1(t) \\ A_2(t) \\ A_3(t) \\ A_4(t) \\ A_5(t) \\ A_6(t) \end{bmatrix} \quad \dots(3.16)$$

are the solutions of eqn. (3.15) and hence they are eigenvectors corresponding to the eigenvalues  $t_1, t_2, t_3, t_4$  of the matrix  $A$ . By actual calculations we obtain

$$\begin{aligned} A_1(t) &= -t^5 + t^4 \left( \frac{\lambda + 3\mu + \alpha}{\lambda + 2\mu} \xi^2 + \frac{4\alpha}{\gamma + \epsilon} \right) \\ &\quad - t \left( \frac{\mu + \alpha}{\lambda + 2\mu} \xi^4 + \frac{4\alpha\mu\xi^2}{(\lambda + 2\mu)(\gamma + \epsilon)} \right) \\ A_2(t) &= t^2 \xi \left[ \frac{\lambda + \mu - \alpha}{\lambda + 2\mu} (t^2 - \xi^2) - \frac{4\alpha(\lambda + \mu)}{(\lambda + 2\mu)(\gamma + \epsilon)} \right] \\ A_3(t) &= \frac{2\alpha t^2 (t^2 - \xi^2)}{(\gamma + \epsilon)} \\ A_4(t) &= -t^4 + t^2 \left[ \frac{\lambda + 3\mu + \alpha}{\lambda + 2\mu} \xi^2 + \frac{4\alpha}{(\gamma + \epsilon)} \right] - \xi^2 \zeta^2 \frac{\mu + \alpha}{\lambda + 2\mu} \\ A_5(t) &= \left\{ \frac{\lambda + \mu - \alpha}{\lambda + 2\mu} (t^2 - \xi^2) - \frac{4\alpha(\lambda + \mu)}{(\lambda + 2\mu)(\gamma + \epsilon)} \right\} t \xi \\ A_6(t) &= \frac{2\alpha}{\gamma + \epsilon} t (t^2 - \xi^2). \end{aligned} \quad \dots(3.17)$$

Since  $t_1$  and  $t_2$  are double root of the characteristic equation (3.14) of the matrix  $A$ , the solution of the differential equation (3.9) is given by (vide Das *et al.*<sup>2</sup>).

$$\begin{aligned}
 V = & C_1 Y(t_1) \exp(t_1 z) + C_2 d/dz [Y(t) \exp(t z)]_{t=t_1} \\
 & + C_3 \dot{Y}(t_2) \exp(t_2 z) + C_3 d/dz [Y(t) \exp(t z)]_{t=t_2} \\
 & + C_5 Y(t_3) \exp(t_3 z) + C_6 Y(t_4) \exp(t_4 z) \quad \dots(3.18)
 \end{aligned}$$

where  $C_1, C_2, \dots, C_6$  are arbitrary constants to be determined from boundary conditions.

Equation (3.18) can be rewritten as

$$\begin{aligned}
 V = & (C_1 + C_2 z) Y(t_1) \exp(t_1 z) + C_2 \dot{Y}(t_1) \exp(t_1 z) \\
 & + (C_3 + C_4 z) Y(t_2) \exp(t_2 z) + C_4 \dot{Y}(t_2) \exp(t_2 z) \\
 & + C_5 Y(t_3) \exp(t_3 z) + C_6 Y(t_4) \exp(t_4 z) \quad \dots(3.19)
 \end{aligned}$$

where dot ( . ) represents the differentiation with respect to  $t$ .

For the half space  $z \geq 0$ , equation (3.19) reduces the form

$$\begin{aligned}
 V = & (C_3 + C_4 z) Y(t_2 z) \exp(t_2 z) + C_4 \dot{Y}(t_2) \exp(t_2 z) \\
 & + C_6 Y(t_4) \exp(t_4 z) \quad \dots(3.20)
 \end{aligned}$$

where the constants  $C_3, C_4$  and  $C_6$  are to be determined from the boundary conditions.

Equations (3.20) can be written explicitly as

$$\begin{aligned}
 \bar{u}'_r(z) = & \{(C_3 + C_4 z) A_1(t_2) + C_4 \dot{A}_1(t_2)\} \exp(t_2 z) \\
 & + C_6 A_1(t_4) \exp(t_4 z) \\
 \bar{u}'_z(z) = & \{(C_3 + C_4 z) A_2(t_2) + C_4 \dot{A}_2(t_2)\} \exp(t_2 z) \\
 & + C_6 A_2(t_4) \exp(t_4 z) \\
 \bar{\omega}'_\varphi(z) = & \{(C_3 + C_4 z) A_3(t_2) + C_4 \dot{A}_3(t_2)\} \exp(t_2 z) \\
 & + C_6 A_3(t_4) \exp(t_4 z) \\
 \bar{u}_r(z) = & \{(C_3 + C_4 z) A_4(t_2) + C_4 \dot{A}_4(t_2)\} \exp(t_2 z) \\
 & + C_6 A_4(t_4) \exp(t_4 z) \\
 \bar{u}_z(z) = & \{(C_3 + C_4 z) A_5(t_2) + C_4 \dot{A}_5(t_2)\} \exp(t_2 z) \\
 & + C_6 A_5(t_4) \exp(t_4 z) \\
 \bar{\omega}_\varphi(z) = & \{(C_3 + C_4 z) A_6(t_2) + C_4 \dot{A}_6(t_2)\} \exp(t_2 z) \\
 & + C_6 A_6(t_4) \exp(t_4 z) \quad \dots(3.21)
 \end{aligned}$$

where  $C_3$ ,  $C_4$  and  $C_6$  are arbitrary constants to be determined from the boundary conditions. Thus the displacements have been obtained in the transformed domain and as such the stresses can also be obtained from (3.2) using (3.21) in the transformed domain.

#### 4. SOLUTION OF EQUATIONS (2.7)

Here we are concerned with the set of equations (2.7) in which the displacement vector  $\mathbf{u} = (0, u_\varphi, 0)$  and the rotation vector  $\boldsymbol{\omega} = (\omega_r, 0, \omega_z)$ . The field of displacements  $(0, u_\varphi, 0)$  and rotations  $(\omega_r, 0, \omega_z)$  described by the set of equations (2.7) induces the following state of force-stress and couple-stress (vide, Dhaliwal)

$$\sigma = \begin{vmatrix} 0 & \sigma_{r\varphi} & 0 \\ \sigma_{\varphi r} & 0 & \sigma_{\varphi z} \\ 0 & \sigma_{z\varphi} & 0 \end{vmatrix} \quad \text{and} \quad \mu = \begin{vmatrix} \mu_{rr} & 0 & \mu_{rz} \\ 0 & \mu_{\varphi\varphi} & 0 \\ \mu_{zr} & 0 & \mu_{zz} \end{vmatrix} \quad \dots(4.1)$$

where

$$\begin{aligned} \sigma_{r\varphi} &= (\mu + \alpha) \frac{\partial u_\varphi}{\partial r} - (\mu - \alpha) \frac{u_\varphi}{r} - 2\alpha \omega_z \\ \sigma_{\varphi r} &= (\mu - \alpha) \frac{\partial u_\varphi}{\partial r} - (\mu + \alpha) \frac{u_\varphi}{r} + 2\alpha \omega_z \\ \sigma_{\varphi z} &= (\mu - \alpha) \frac{\partial u_\varphi}{\partial z} - 2\alpha \omega_r \\ \sigma_{z\varphi} &= (\mu + \alpha) \frac{\partial u_\varphi}{\partial z} + 2\alpha \omega_r \\ \mu_{rr} &= \beta \chi + 2\gamma \frac{\partial \omega_r}{\partial r}, \quad \mu_{zz} = \beta \chi + 2\gamma \frac{\partial \omega_z}{\partial z}, \quad \mu_{\varphi\varphi} = \beta \chi + 2\gamma \omega_r \\ \mu_{rz} &= (\gamma - \epsilon) \frac{\partial \omega_r}{\partial z} + (\gamma + \epsilon) \frac{\partial \omega_z}{\partial r}, \quad \mu_{zr} = (\gamma + \epsilon) \frac{\partial \omega_z}{\partial z} \\ &\quad + (\gamma - \epsilon) \frac{\partial \omega_r}{\partial r}. \end{aligned} \quad \dots(4.2)$$

Now Hankel transform of the set of equations (2.7) give

$$\left. \begin{aligned} &[(\gamma + \epsilon) D^2 - (\beta + 2\gamma) \xi^2 - 4\alpha] \bar{\omega}_r - (\beta + \gamma - \epsilon) \xi D \bar{\omega}_z \\ &\quad - 2\alpha D \bar{u}_\varphi = 0 \\ &(\beta + \gamma - \epsilon) \xi D \bar{\omega}_r + [(\beta + 2\gamma) D^2 - (\gamma + \epsilon) \xi^2 - 4\alpha] \bar{\omega}_z \\ &\quad + 2\alpha \xi \bar{u}_\varphi = 0 \\ &2\alpha D \bar{\omega}_r + 2\alpha \xi \bar{\omega}_z + (\mu + \alpha) (D^2 - \xi^2) \bar{u}_\varphi = 0. \end{aligned} \right\} \quad \dots(4.3)$$

Now as in section 3 equations (4.3) can be written as vector-matrix differential equation form as

$$\frac{d}{dz} (V) = B V \quad \dots(4.4)$$

where

$$V = \begin{bmatrix} \bar{\omega}'_r \\ \bar{\omega}'_z \\ \bar{u}'_\varphi \\ \bar{\omega}_r \\ \bar{\omega}_z \\ \bar{u}_\varphi \end{bmatrix} \quad B = \begin{bmatrix} 0 & b_{12} & b_{13} & b_{14} & 0 & 0 \\ b_{21} & 0 & 0 & 0 & b_{25} & b_{26} \\ b_{31} & 0 & 0 & 0 & b_{35} & b_{36} \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \end{bmatrix} \quad \dots(4.5)$$

and

$$\begin{aligned} b_{12} &= \frac{\beta + \gamma - \epsilon}{\gamma + \epsilon} \xi, & b_{13} &= \frac{2\alpha}{\gamma + \epsilon}, & b_{14} &= \frac{(\beta + 2\gamma)\xi^2 - 4\alpha}{\gamma + \epsilon} \\ b_{21} &= -\frac{\beta + \gamma - \epsilon}{\beta + 2\gamma} \xi, & b_{25} &= \frac{(\gamma - \epsilon)\xi^2 + 4\alpha}{\beta + 2\gamma}, & b_{26} &= \frac{-2\alpha\xi}{\beta + 2\gamma} \\ b_{31} &= \frac{-2\alpha}{\mu + \alpha}, & b_{35} &= \frac{-2\alpha\xi}{\mu + \alpha}, & b_{36} &= \xi^2 \end{aligned} \quad \dots(4.6)$$

Assume  $V = X \exp(tz)$  be the solution of the equation (4.4). Then we must have

$$B X = t X. \quad \dots(4.7)$$

This gives that  $t$  is an eigenvalue of the matrix  $B$  and  $X$  the corresponding eigenvectors. The eigenvalues for the matrix  $B$  are the roots of

$$\det(B - tI) = 0 \quad \dots(4.8)$$

That is,

$$\begin{vmatrix} -t & b_{12} & b_{13} & b_{14} & 0 & 0 \\ b_{21} & -t & 0 & 0 & b_{25} & b_{26} \\ b_{31} & 0 & -t & 0 & b_{35} & b_{36} \\ 1 & 0 & 0 & -t & 0 & 0 \\ 0 & 1 & 0 & 0 & -t & 0 \\ 0 & 0 & 1 & 0 & 0 & -t \end{vmatrix} = 0 \quad \dots(4.9)$$

where  $b_{12}$ ,  $b_{13}$ , etc. are given by (4.6).

Now simplifying (4.9) and using (4.6) therein we obtain the characteristic equation as

$$t^6 - t^4 \left( \xi^2 + \lambda_1^2 + \lambda_2^2 \right) + t^2 \left( \xi^2 \lambda_1^2 + \xi^2 \lambda_2^2 + \lambda_1^2 \lambda_2^2 \right) - \xi^2 \lambda_1^2 \lambda_2^2 = 0 \quad \dots(4.10)$$

where

$$\lambda_1^2 = \xi^2 + K_1^2, \quad \lambda_2^2 = \xi^2 + K_2^2$$

$$K_1^2 = \frac{4\alpha}{\beta + 2\gamma}, \quad K_2^2 = \frac{4\alpha\mu}{(\mu + \alpha)(\gamma + \epsilon)}$$

whose roots are  $\xi, -\xi, \lambda_1, -\lambda_1, \lambda_2, -\lambda_2$ , which are the distinct eigenvalues of the matrix  $B$ . The corresponding eigenvectors are obtained by solving the following homogeneous equation.

$$\begin{bmatrix} -t & b_{12} & b_{13} & b_{14} & 0 & 0 \\ b_{21} & -t & 0 & 0 & b_{25} & b_{26} \\ b_{31} & 0 & -t & 0 & b_{35} & b_{36} \\ 1 & 0 & 0 & -t & 0 & 0 \\ 0 & 1 & 0 & 0 & -t & 0 \\ 0 & 0 & 1 & 0 & 0 & -t \end{bmatrix} \begin{bmatrix} x_1(t) \\ x_2(t) \\ x_3(t) \\ x_4(t) \\ x_5(t) \\ x_6(t) \end{bmatrix} = 0 \quad \dots(4.11)$$

for  $t = t_i, i = 1, 2, \dots, 6$  where  $t_1 = \xi, t_2 = -\xi, t_3 = \lambda_1, t_4 = -\lambda_1, t_5 = \lambda_2, t_6 = -\lambda_2$ .

Denote by  $B_i(t), i = 1, 2, \dots, 6$ , the co-factors of the elements of the first row of the coefficient matrix in eqn. (4.11).

Then

$$X(t) = \begin{bmatrix} B_1(t) \\ B_2(t) \\ B_3(t) \\ B_4(t) \\ B_5(t) \\ B_6(t) \end{bmatrix}$$

are the solutions of the eqn. (4.11) and hence they are eigenvectors corresponding to the eigenvalues  $t_i, i = 1, 2, \dots, 6$ , of the matrix  $B$ . By actual calculation we obtain

$$B_1(t) = -t^5 + t^2 \left\{ \frac{\xi^2(\beta + 3\gamma + \epsilon) + 4\alpha}{\beta + 2\gamma} \right\}$$

(equation continued on p. 1249)



$$- t \xi^2 \left\{ \frac{(\mu + \alpha)(\gamma + \epsilon) \xi^2 + 4\alpha\mu}{(\beta + 2\gamma)(\mu + \alpha)} \right\}$$

$$B_2(t) = - \frac{(\beta + \gamma - \epsilon)}{(\beta + 2\gamma)} t^4 \xi - t^2 \xi \left\{ \frac{4\alpha^2 - (\mu + \alpha)(\beta + \gamma - \alpha\xi^2)}{(\beta + 2\gamma)(\mu + \alpha)} \right\}$$

$$B_3(t) = - t^4 \frac{2\alpha}{\mu + \alpha} - t^2 \frac{2\alpha \xi^2 (\beta + \gamma - \epsilon)}{(\mu + \alpha)(\beta + 2\gamma)}$$

$$B_4(t) = t^4 - t^2 \left\{ \frac{(\beta + \gamma - \epsilon) \xi^2 - 4\alpha}{\beta + 2\gamma} \right\} \\ + \xi^2 \frac{(\gamma + \epsilon)(\mu + \alpha) \xi^2 + 4\alpha\mu}{(\mu + \alpha)(\beta + 2\gamma)}$$

$$B_5(t) = t^3 \xi \frac{\beta + \gamma - \epsilon}{\beta + 2\gamma} - t \left\{ \frac{\beta + \gamma - \epsilon}{\beta + 2\gamma} \xi^3 + \frac{4\alpha^2 \xi}{(\beta + 2\gamma)(\mu + \alpha)} \right\}$$

$$B_6(t) = - t^3 \frac{2\alpha}{\mu + \alpha} - t \left\{ \frac{2\alpha \xi (\beta + \gamma - \epsilon) - 2\alpha(\gamma + \epsilon) \xi^2 + 8\alpha^2}{(\beta + 2\gamma)(\mu + \alpha)} \right\}$$

Since  $t_1, t_2, \dots, t_6$  are all distinct roots of the characteristic equation (4.10) of the matrix  $B$ , the general solution of the differential equation (4.4) is given by (vide appendix A)

$$V = E_1 X(t_1) \exp(t_1 z) + E_2 X(t_2) \exp(t_2 z) + E_3 X(t_3) \exp(t_3 z) \\ + E_4 X(t_4) \exp(t_4 z) + E_5 X(t_5) \exp(t_5 z) \\ + E_6 X(t_6) \exp(t_6 z)$$

where

$$Et, i = 1, 2, \dots, 6.$$

are arbitrary constants

i. e.

$$V = \sum_{i=1}^6 E_i X(t_i) \exp(t_i z)$$

## REFERENCES

1. L. Y. Bahar and R. B. Hetnarski, *J. Therm. Stresses*, 2 (1979), 283.
2. N. C. Das, S. N. Das and B. Das, *J. Therm. Stresses*, (6), (1983), 35.
3. N. C. Das, A. K. Mitra and R. K. Mahalanabis, *J. Therm. Stresses* 6 (1983), 73-85.
4. N. C. Das, P. C. Bhakta and S.K. Bhattacharyya, *Eigenvalue Approach to Plane Problem to Elasticity*, Presented in the 24th Polish Solid Mechanics Conference, Warsaw, Poland, August 1983.

5. R. S. Dhaliwal and K. L. Chowdhury, *Bull. de L' Academic Pol. des. Sc.* **19** (1971), 363.
6. R. S. Dhaliwal, *Arch. Mech. Stos. Warsaw* **23** (1971), 705-14.
7. R. S. Dhaliwal, *Arch. Mech. Stos. Warsaw* **24** (1972), 645-53.
8. A. C. Eringen, *Theory of Micropolar Elasticity, Treatise on Fracture* (ed. H. Leibowitz), Vol. II: Academic Press, New York and London, 1968, pp. 621-729.
9. A. C. Eringen and E. S. Suhubi, *Ind. J. Engg. Sci.* **2** (1964), 189; **2** (1964), 389.
10. E. V. Kuvchinski and E. L. Aero, *Fiz. Trivierd Tiela*, **5**, 1963.
11. W. Nowaki, *Proc. Vibr. Probl. Warsaw*, **10** (1969), 97.
12. W. Nowacki, *CISM Courses and Lectures No. 25*, Springer Verlag, New York, 1970.
13. M. A. Palmov, *Prikl. Math., Mekl.* **28** (1964), 40.
14. P. Puri, *ZAMP* **22** (1971), 320.

#### APPENDIX A

Consider the differential equation

$$\frac{dv}{dt} = Av \quad \dots(A1)$$

where  $v$  is an  $n$  vector and  $A$  is an  $n \times n$  real constant matrix.

If  $\lambda_1, \lambda_2, \dots, \lambda_n$  are distinct eigenvalues of the matrix  $A$  and  $x_1, x_2, \dots, x_n$  be the corresponding eigenvectors of  $A$ , then the general solution of (A1) is given by

$$v(t) = C_1 X_1 e^{\lambda_1 t} + \dots + C_n X_n e^{\lambda_n t}$$

where  $C_1, \dots, C_n$ , are arbitrary constants.

If  $\lambda_1$  is an eigenvalue of  $A$  of multiplicity 2 and all other eigenvalues  $\lambda_3, \dots, \lambda_n$  are of multiplicity one and  $X_1, X_3, \dots, X_n$  the corresponding eigenvector of  $A$ , then the general solution of (A1) is given by

$$v(t) = C_1 X_1 e^{\lambda_1 t} + C_2 \frac{d}{dt} (X_1 e^{\lambda_1 t}) + C_3 X_3 e^{\lambda_3 t} + \dots \\ + C_n X_n e^{\lambda_n t}$$

where  $C_1, C_2, \dots, C_n$  are all arbitrary constants.

For details, vide Das *et al.*<sup>2</sup>.

## MELLIN TRANSFORM OF THE GRAVITY EFFECT OF A 2—D HORIZONTAL CIRCULAR CYLINDER WITH VARIABLE DENSITY

L. ANAND BABU<sup>1</sup>, N. L. MOHAN<sup>2</sup>, N. SUNDARARAJAN<sup>2</sup>

AND

S. V. SESHAGIRI RAO<sup>2</sup>

<sup>1</sup>*Department of Mathematics, Osmania University, Hyderabad 500007*

<sup>2</sup>*Centre of Exploration Geophysics, Osmania University, Hyderabad 500007*

(Received 30 October 1988; after revision 17 May 1989; accepted 1 July 1989)

The Mellin transform of the gravity effect of a buried 2—D horizontal circular cylinder with density contrast varying linearly with depth is derived and analysed to extract the body parameters. The characteristic features of the Mellin transform of the gravity effect resemble gamma function. The validity of the method is tested on simulated models. The stability of the Mellin transform is studied by incorporating random noise in the gravity effect at different levels and subsequently the error estimation of the interpreted values is discussed.

### INTRODUCTION

In general, the interpretation of the geophysical anomalies are carried out by assuming certain geometrical shapes with uniform physical properties. Very frequently, we come across some practical cases wherein for example the density contrast increases with the increase of depth which is evident from seismic studies.

Therefore, it would be meaningful to interpret such geophysical anomalies with non uniform density<sup>4</sup>. Herein the analysis of gravity anomalies due to a horizontal circular cylinder with the density contrast varying linearly with depth is presented using the Mellin transform. Such transformation of gravity (or magnetic) anomalies paves the way for simplified analysis of the complex potential field<sup>1</sup>. The procedure is illustrated with three sets of theoretical models. The stability of the Mellin transform is studied by incorporating random noise in the gravity effect at different levels and subsequently the error estimation of the interpreted values is discussed.

### MELLIN TRANSFORM OF THE GRAVITY EFFECT OF THE HORIZONTAL CIRCULAR CYLINDER

A buried horizontal circular cylinder extending infinitely along the  $Y$ -direction, with its normal section parallel to the  $X - Z$  plane is considered. The origin of the

coordinate system is taken on the ground surface such that the  $Z$ -axis coincides with the diameter (Fig. 1a). Let the density contrast at the apex of the cylinder as  $\rho$  and the rate of change of density contrast varying linearly with depth be  $a$ . In this case, the gravity effect of the cylinder is given by Radhakrishnan Murthy<sup>5</sup>.

$$g(x) = A \frac{Z}{X^2 + Z^2} - B \frac{Z^2 - X^2}{(X^2 + Z^2)^2} \quad \dots(1)$$

where

$$A = 2\pi GR^2 (\rho + aR) \quad \dots(1a)$$

$$B = \frac{\pi GaR^4}{2} \quad \dots(1b)$$

$Z$  is the depth to the centre of the cylinder,  $R$  the radius of the cylinder and  $G$  the universal gravitational constant.

The Mellin transform of the gravity effect given by eqn. (1) is written as Sneddon<sup>6</sup>

$$M(s) = \int_0^\infty x^{s-1} g(x) dx \quad \dots(2)$$




FIG. 1 (a) Cross section of the buried horizontal circular cylinder.

(b) Computed gravitational effect of horizontal circular cylinder

where  $s$  is a real positive integer or fractional number.

Substituting for  $g(x)$  and integrating eqn. (2) with respect to  $x$ , we get the Mellin transform of  $g(x)$  as :

$$M(s) = [A Z^{s-1} \Gamma(s/2) \Gamma((2-s)/2) - B Z^{s-2} \times \{\Gamma(s/2) \Gamma((4-s)/2) - \Gamma((s+2)/2) \Gamma((2-s)/2)\}] \dots (3)$$

$$(0 < s < 2)$$

#### ANALYSIS

For a set of arbitrary values of  $s$ , i. e., for  $s = 1/4$ ;  $s = 1/2$ ;  $s = 3/4$ ;  $s = 1$  and  $s = 5/4$  equation (3) is written as :

$$M(1/4) = A Z^{-3/4} \Gamma(1/8) \Gamma(7/8) - B Z^{-7/4} [\Gamma(1/8) \Gamma(15/8) - \Gamma(9/8) \Gamma(7/8)] \dots (4a)$$

$$M(1/2) = A Z^{-1/2} \Gamma(1/4) \Gamma(3/4) - B Z^{-3/2} [\Gamma(1/4) \Gamma(7/4) - \Gamma(5/4) \Gamma(3/4)] \dots (4b)$$

$$M(3/4) = A Z^{-1/4} \Gamma(3/8) \Gamma(5/8) - B Z^{-5/4} [\Gamma(3/8) \Gamma(13/8) - \Gamma(11/8) \Gamma(5/8)] \dots (4c)$$

$$A = M(1)/\pi \dots (4d)$$

$$M(5/4) = A Z^{1/4} \Gamma(5/8) \Gamma(3/8) - B Z^{-3/4} [\Gamma(5/8) \Gamma(11/8) - \Gamma(13/8) \Gamma(3/8)] \dots (4e)$$

From eqns. (4c) and (4e), the value of  $Z$  is evaluated as :

$$M(3/4) Z^{1/4} = U - (B/Z) V \dots (5)$$

and

$$M(5/4) Z^{1/4} = U + (B/Z) V \dots (6)$$

i. e.,

$$P (Z^{1/4})^2 - U Z^{1/4} + Q = 0 \dots (7)$$

where

$$P = M(3/4)/2,$$

$$Q = M(5/4)/2$$

$$U = M(1) \Gamma(3/4) \Gamma(5/8)/\pi,$$

$$V = \Gamma(3/8) \Gamma(13/8) - \Gamma(11/8) \Gamma(5/8).$$

Hence

$$Z = \left\{ \frac{U \pm (U^2 - 4PQ)^{1/2}}{2P} \right\}^4 \dots (8)$$



Since  $A$  and  $Z$  are known,  $B$  can be evaluated as :

$$B = Z \left[ \frac{M(5/4) Z^{-1/4} - M(3/4) Z^{1/4}}{2V} \right] \quad \dots(9)$$

By eliminating ' $a$ ' from equations (1a) and (1b) a cubic equation in ' $R$ ' is obtained as :

$$R^3 - (A/2 \rho \pi G) R + (4B/2 \rho \pi G) = 0. \quad (10)$$

Applying the well known Cardon's method,  $R$  is evaluated, and subsequently  $a$  is calculated as :

$$a = 2B/R^4. \quad \dots(11)$$

### DISCRETE MELLIN TRANSFORM

Since the gravity data is collected at discrete intervals in the real field situation, the numerical computation of the Mellin transform is carried out by formulating the discrete Mellin transform as<sup>3</sup>




FIG. 2. Continuous Mellin transform of the gravity effect of horizontal circular cylinder with density variation.

$$M(1, \Delta s) = \sum_{l=1}^{N-1} g(n, \Delta x) (n, \Delta x)^{1, \Delta s-1} \Delta x \quad \dots(12)$$

$$(0 < 1, \Delta s < 2)$$

where  $\Delta s = 1/4$ ,  $N$  is the total number of points and  $\Delta x$  is the sample interval.

#### SYNTHETIC EXAMPLES

The procedure detailed in the text is illustrated with three theoretical models (Table I). The theoretical Mellin transform of the gravity effect of three models are computed using eqn (3) and shown in Fig 2.

The gravity effect due to horizontal circular cylinder with variable density for three models are computed using eqn. (1). Since the gravity effect due to horizontal circular cylinder is symmetric, only the positive side of the anomaly is shown in Fig. 1b. The discrete Mellin transform of the gravity effect of the cylindrical models are computed using eqn. (12) and shown in Fig. 3. It may be observed that the discrete Mellin




FIG. 3. Discrete Mellin transform of the gravity effect of horizontal circular cylinder with density variation.

TABLE I Theoretical Examples

		$R^*$	$h^*$	$a^*$
Model I	Assumed values	0.50	0.75	1.00
	Evaluated values	0.56	0.70	0.92
Model II	Assumed values	1.00	2.00	1.50
	Evaluated values	0.95	1.88	1.39
Model III	Assumed values	1.50	3.50	1.75
	Evaluated valves	1.45	2.95	1.80

(\* in arbitrary units)




FIG. 4. Correlation coefficient versus number of discrete gravity samples.




FIG. 5. Standard deviation versus number of discrete gravity samples.

transform of the gravity effect of the models (Fig. 3) are similar to the theoretical Mellin transform (Fig. 2) and they resemble gamma function curves. The parameters are evaluated from the computed discrete Mellin transform of the gravity models using eqns. (4d), (8) (10) and (11) and Fig. (3) and presented in Table 1. It may be noticed that the evaluated parameters reasonably agree with the assumed values.




FIG. 6. Normalized error variation versus number of discrete gravity samples.

### STABILITY ANALYSIS

Here for all the three gravity models the number of discrete samples of the gravity effect are considered at six different levels ranging from 100 to 600 at an interval of 100. In each case the random noise with different percentages i.e., 10%, 20%, 30% and 40% are added to the discrete gravity effect of the horizontal circular cylinder with variable density. The discrete Mellin transform of the noise contaminated gravity effect is computed. Using computed theoretical Mellin transform of the gravity effect and the discrete Mellin transform of the noise contaminated gravity effect for all the



models for different levels of number of samples and with different levels of random noise, the correlation coefficient, standard deviation and the error variation (normalized) are computed and shown in Figs. 4, 5 and 6.

It is observed from the Fig. (4) (i.e.) correlation coefficient versus number of samples with different levels of noise for the gravity models, the correlation coefficient is tending to 1 for discrete gravity samples  $> 400$ . Also the standard deviation and error variation (normalized) are saturating for discrete gravity samples  $> 400$ .

The error percentage in evaluated parameters of the cylinder is relatively high ( $> 30\%$ ) for the discrete gravity samples  $< 400$  for different levels of random noise.

#### ACKNOWLEDGEMENT

The authors wish to thank the University Grants Commission for financial aid given under UGC Minor research Project. The authors wish to record their thanks to the reviewers for excellent suggestions.

#### REFERENCES

1. L. Anand Babu, Ph. D. Thesis submitted to Osmania University, Hyderabad 1985.
2. N. L. Mohan, Ph. D. Thesis submitted to Osmania University, Hyderabad, 1978.
3. N. L. Mohan, L. Anand Babu and S. V. Seshagiri Rao, *Geophysics* **51** (1986), 114-22.
4. N. L. Mohan, S. V. Seshagiri Rao and V. L. S. Bhimasankaram, *Current Sci* **46** (1977), 404-5.
5. I. V. Radhakrishna Murthy, *Rivista Italiana Di Geofiscia*, **22** (1973), 407-10.
6. I. N. Sneddon, *The Use of Integral Transform*, Tata Mc-Graw-Hill Publication Company Ltd, New Delhi, 1979.



# Indian Journal of Pure & Applied Mathematics

CONTENTS & INDEX

Volume 20 (1989)



INDIAN NATIONAL SCIENCE ACADEMY  
BAHADUR SHAH ZAFAR MARG, NEW DELHI-110 002

# INDIAN JOURNAL OF PURE AND APPLIED MATHEMATICS

*Published monthly by the*

INDIAN NATIONAL SCIENCE ACADEMY

Editor of Publications

PROFESSOR D. V. S. JAIN

Department of Physical Chemistry, Panjab University

Chandigarh 160 014

PROFESSOR J. K. GHOSH

Indian Statistical Institute

203, Barrackpore Trunk Road

Calcutta 700 035

PROFESSOR A. S. GUPTA

Department of Mathematics

Indian Institute of Technology

Kharagpur 721 302

PROFESSOR M. K. JAIN

Department of Mathematics

Indian Institute of Technology

Hauz Khas

New Delhi 110 016

PROFESSOR S. K. JOSHI

Director

National Physical Laboratory

New Delhi 110 012

PROFESSOR V. KANNAN

Dean, School of Mathematics &

Computer/Information Sciences

University of Hyderabad

P O Central University

Hyderabad 500 134

Assistant Executive Secretary

(Associate Editor/Publications)

DR. M. DHARA

Subscriptions :

For India, Pakistan, Sri Lanka, Nepal, Bangladesh and Burma, Contact :

Associate Editor, Indian National Science Academy, Bahadur Shah Zafar Marg,

New Delhi 110002, Telephone : 3311865, Telex : 31-61835 INSA IN.

For other countries, Contact :

M/s J. C. Baltzer AG, Scientific Publishing Company, Wettsteinplatz 10, CH-4058 Basel, Switzerland, Telephone : 61-268925, Telex : 63475.

*The Journal is indexed in the Science Citation Index; Current Contents (Physical, Chemical & Earth Sciences); Mathematical Reviews; INSPEC Science Abstracts (Part A); as well as all the major abstracting services of the World.*

PROFESSOR N. MUKUNDA

Centre for Theoretical Studies

Indian Institute of Science

Bangalore 560 012

DR PREM NARAIN

Director

Indian Agricultural Statistics

Research Institute, Library Avenue

New Delhi 110 012

PROFESSOR I. B. S. PASSI

Centre for advanced study in Mathematics

Panjab University

Chandigarh 160 014

PROFESSOR PHOOLAN PRASAD

Department of Applied Mathematics

Indian Institute of Science

Bangalore 560 012

PROFESSOR M. S. RAGHUNATHAN

Senior Professor of Mathematics

Tata Institute of Fundamental Research

Homi Bhabha Road

Bombay 500 005

PROFESSOR T. N. SHOREY

School of Mathematics

Tata Institute of Fundamental Research

Homi Bhabha Road

Bombay 400 005

Assistant Editor

SRI R. D. BHALLA

# CONTENTS

	<i>Page</i>
On the average number of crossings of an algebraic polynomial <i>by</i> K. FARAHMAND	1
Generalized convexity in multi-objective programming <i>by</i> M. N. VARTAK and INDRANI GUPTA	10
On automorphisms of free groups <i>by</i> A. G. A. E. ABDEL-GAWAD	40
Artinian (Noetherian) part of a Goldie ring <i>by</i> K. C. CHOWDHURY	49
A generalization of strongly regular near-rings <i>by</i> P. DHEENA	58
Lacunary distribution of sequences <i>by</i> G. DAS and B. K. PATEL	64
Note on the small vibration of beams with varying Young's modulus carrying a concentrated mass distribution <i>by</i> P. K. CHAUDHURI and SUBRATA DATTA	75
Free torsional vibration of a nonhomogeneous semiinfinite solid circular cylinder <i>by</i> N. C. MONDAL	89
Erratum "An analogue of Hoffman-Wermer theorem for a real function algebra" <i>by</i> S. H. KULKARNI and N. SRINIVASAN	98
Cyclotomic numbers and a conjecture of Snapper <i>by</i> S. A. KATRE	99
Stability in mammillary compartmental systems <i>by</i> H. EL-OWAIDY, A. A. AMMAR and O. A. ELLEITHY	104
An optimal programme for augmentation of capacities of depots and shipment of buses from depots to starting points of routes <i>by</i> A. K. AGRAWAL and S. L. DHINGRA	111
On a class of nonlinear higher order differential equations <i>by</i> B. G. PACHPATTE	121
Fixed points iterations for non-linear Hammerstein equation involving non-expansive and accretive mappings <i>by</i> C. E. CHIDUME	129
On Wagner spaces of $W_p$ -scalar curvature <i>by</i> S. K. SINGH	136



	<i>Page</i>
Goldie theorem analogue for Goldie near-rings <i>by</i> K. C. CHOWDHURY ...	141
Matrix transformations of orthonormal series <i>by</i> B. E. RHOADES ...	150
Stresses in pre-stressed dry sandy soil due to normal moving load leading to instability and fracture <i>by</i> S. DEY <i>and</i> M. CHAKRABORTY ...	165
Thermo-elastic waves from suddenly punched hole in stretched elastic plate <i>by</i> A. B. KUMAR ... ..	181
Note on minmax principle for heat convection equation <i>by</i> M. A. GOPALAN	189
Flow behind weak and strong shock waves in water <i>by</i> V. P. SINGH <i>and</i> A. M. N. YOGI ... ..	194
Mathematical model of population interactions with dispersal : Stability of two habitats with a predator <i>by</i> H. EL-OWAIDY <i>and</i> A. A. AMMAR ... ..	205
On controllability of nonlinear systems with distributed delays in the control <i>by</i> JERRY U. ONWUATU ... ..	213
A Finslerian extensions of the Gravitational field-II <i>by</i> SATOSHI IKEDA ...	229
Periodic solutions of a certain fourth order differential equation <i>by</i> AYDIN TIRYAKI ... ..	235
Cubic transformations of Finsler spaces and $n$ fundamental forms of their hypersurfaces <i>by</i> B. N. PRASAD <i>and</i> J. N. SINGH ... ..	242
Copure-injective modules <i>by</i> V. A. HIREMATH ... ..	250
Inclusion theorems on matrix transformations of some sequence spaces over non-Archimedean fields IV <i>by</i> D. SOMASUNDARAM ... ..	260
Commencement of Couette flow in Oldroyd liquid with heat sources <i>by</i> G. C. DASH <i>and</i> S. BISWAL ... ..	267
On Rayleigh waves in Green-Lindsay's model of generalized thermoelastic media <i>by</i> N. C. DAWN <i>and</i> S. K. CHAKRABORTY ... ..	276
Roche harmonics for stellar models distorted by differential rotation <i>by</i> V. P. SINGH <i>and</i> VANDANA GUPTA ... ..	284
Incoming water waves against a vertical cliff in a two-fluid medium <i>by</i> P. K. KUNDU ... ..	292
An oscillation criterion for second order nonlinear differential equation <i>by</i> S. R. GRACE ... ..	297

On the uniform stability of a system of differential equations with complex coefficients by Z. ZAHREDDINE ... ..	307
On the Graphoidal covering number of a graph by C. PAKKIAM and S. ARUMUGAM ... ..	330
On $\alpha$ -hausdorff subsets, almost closed mappings and almost upper semi-continuous decomposition by ILIJA KOVACEVIĆ ... ..	334
Convex univalent polynomials by M. S. KASI ... ..	341
Modified means by G. DAS ... ..	347
Scattering of a compressional wave at the corner of a quarter space by NARINDER MOHAN and P. S. DESHWAL ... ..	386
On temperature-rate dependent thermoelastic longitudinal vibrations of an infinite circular cylinder by D. YADIAH and RAM KUMAR SHUKLA ...	395
Linear stability and the resonance for the triangular libration points for the doubly photogravitational elliptic restricted problem of three bodies by V. KUMAR and R. K. CHOUDHRY ... ..	403
Comments on steady plane MHD flows with constant speed along each streamline by O. P. CHANDNA ... ..	423
Selection of optimal site for new depot of specified capacity with two objectives by SATYA PRAKASH and VIVEK SAINI ... ..	425
A note on Normed near-algebras by T. SRINIVAS and K. YUGANDHAR ...	433
Radical Goldie near-rings by K. C. CHOWDHURY ... ..	439
Some results on stability of differential systems with impulsive perturbations by R. N. MUKHERJEE and RAGHWENDRA ... ..	446
A transformation of the Finsler metric by an $h$ -vector by B. N. PRASAD and LALJI SRIVASTAVA ... ..	455
Remarks on submanifolds of codimension 2 of an even-dimensional euclidean space by BYUNG HAK KIM ... ..	466
Almost irresolute functions by F. CAMMAROTO and T. NOIRI ... ..	472
On some dual integral equations involving Bessel function of order one by A. CHAKRABARTI ... ..	483
Banach space valued distributional Mellin transform and form invariant linear filtering by A. K. TIWARI ... ..	493

	<i>Page</i>
A cylindrical wave-maker in liquid of finite depth with an inertial surface by B. N. MANDAL and KRISHNA KUNDU ... ..	505
Numerical solution of unsteady flow and heat transfer in a micropolar fluid past a porous flat plate by R. S. AGARWAL and C. DHANPAL ...	513
Torsional vibration of a random elastic cylinder by F. D. ZAMAN, S. ASGHAR and G. GHOU ... ..	521
Flow of a conducting fluid between two coaxial rotating porous cylinders bounded by a permeable bed by K. JAGADEESWARA PILLAI, S. V. K. VARMA and M. SYAM BABU ... ..	526
On an error term involving the Totient function by WERNER GEORG NOWAK	537
A fixed point theorem for generalized contraction map by A CARBONE, B. E. RHOADES and S. P. SINGH ... ..	543
Sequences of mappings converging to a contraction mapping by THEODOR VIDALIS ... ..	549
Noetherian regular rings by C. JAYARAM and V. L. MANNEPALLI ...	554
Differential subordination and conformal mappings I by V. KARUNAKARAN and S. PONNUSAMY ... ..	560
On a quaternion submanifolds of co-dimension-2 by I. C. GUPTA and A. K. AGARWAL ... ..	566
On almost continuous functions by TAKASHI NOIRI ... ..	571
On the multivalent functions by MAMORU NUNOKAWA ... ..	577
The Hankel-Clifford transformation on certain spaces of ultradistributions by J. J. BETANCOR ... ..	583
A finite integral involving a general class of polynomials and the multivari- able $H$ -function by K. C. GUPTA and S. M. AGRAWAL ... ..	604
On strongly rare-continuity by NURETTIN ERGUN ... ..	609
Quasi-static response of a layered half-space to surface loads by NAT RAM GARG and SARVA JIT SINGH ... ..	621
A note on the squeeze film lubrication with non-Newtonian fluid by N. M. BUJURKE, S. G. BHAVI and P. S. HIREMATH ... ..	632
Circular orbits of charged test particles in Riessner-Nordstrom field by ABDUSSATTAR and REHANA QURAISHI ... ..	641

Random Rayleigh waves in non-homogeneous elastic media <i>by</i> K. L. DUTTA and S. K. CHAKRABORTY ... ..	646
On the real roots of a random algebraic polynomial <i>by</i> S. BAGH ...	655
Gronwall, Bihari and Langenhop type inequalities for discrete Pfaffian equation <i>by</i> E. THANDAPANI ... ..	665
A note on primary decomposition in Noetherian near-rings <i>by</i> K. YUGANDHAR, K. RAJA GOPAL RAO and T. SRINIVAS ... ..	671
Some results on almost semi-invariant submanifold of an $Sp$ -Sasakian manifold <i>by</i> KALPANA ... ..	681
Maximal elements in Banach spaces <i>by</i> GHANSHYAM MEHTA ... ..	690
On the Endl-type generalization of certain summability methods <i>by</i> M. R. PARAMESWARAN ... ..	698
Matrix transformations in some sequence spaces <i>by</i> SUDARSAN NANDA ...	707
Transient forced and free convection flow past an infinite vertical plate <i>by</i> M. D. JAHAGIRDAR and R. M. LAHURIKAR ... ..	711
Effect of thermal diffusion on thermohaline interleaving in a porous medium due to horizontal gradients <i>by</i> C. P. PARVATHY and PRABHAMANI R. PATIL ... ..	716
Hodograph transformation in constantly inclined two-phase MFD flows <i>by</i> CHANDRESHWAR THAKUR and RAM BABU MISHRA ... ..	728
Thermal stability of a fluid layer in a variable gravitational field <i>by</i> G. K. PRADHAN, P. C. SAMAL and U. K. TRIPATHY ... ..	736
Coset diagrams for an action of the extended modular group on the pro- jective line over a finite field <i>by</i> QAISER MUSHTAQ ... ..	747
The extended modular group acting on the projective line over a Galois field <i>by</i> Q. MUSHTAQ ... ..	755
On the existence for a class of optimal control problems <i>by</i> GARGI CHAKRABORTY ... ..	761
Bounds for the zeros of polynomial <i>by</i> M. BIDKHAM and K. K. DEWAN ...	768
Selection procedures for hazard rates based on two-sample statistics <i>by</i> AMAR NATH GILL and GOBIND P. MEHTA ... ..	773
On digraph reconstruction <i>by</i> S. RAMACHANDRAN ... ..	782



	<i>Page</i>
Variants of Hopfcity in modules by B. M. PANDEYA, S. A. PARAMHANS and S. P. KOIRALA ... ..	786
On $h\nu$ -recurrent Finsler connection by B. N. PRASAD and LALJI SRIVASTAVA ... ..	790
Hypersurfaces with $(f, g, u, v, \lambda)$ -structure of an affinely cosymplectic mani- fold by DHRUWA NARAIN ... ..	799
$O$ -distributive posets by Y. S. PAWAR and V. B. DHAMKE ... ..	804
Functional limits by G. DAS and S. NANDA ... ..	812
Extreme points of some families of analytic functions related to univalent functions by A. K. MISHRA and P. SAHU ... ..	820
A generalized Carleman boundary value problem for multiply connected domains by M. G. EL SHEIKH ... ..	829
Nonstationary law of heat conduction in classical thermoelastic solid by AMIYADEB MUKHERJEE ... ..	834
Ellipsoidal inclusions in an elastic medium by A. D. ALAWNEH and N. T. SHAWAGFEH ... ..	840
On a monotonicity property of measures of directed-divergence by J. N. KAPUR and G. P. TRIPATHI ... ..	851
Bayes approach to prediction in samples from gamma population when outliers are present by G. S. LINGAPPAIAH ... ..	858
On the forms of $n$ for which $\varphi(n)/n-1$ by V. SIVA RAMA PRASAD and M. RANGAMMA ... ..	871
The injective hull of a module with FGD by S. BHAVANARI ... ..	874
On complete integral closure of $G$ -domain by SURJIT SINGH and PAMMY MANCHANDA ... ..	884
On almost unified contact Finsler structures and connections by B. B. SINHA and R. K. YADAV ... ..	887
On $F$ -absolutely translatable summability methods by C. ORHAN and M. A. SARIGOL ... ..	893
Growth of composite integral functions by INDRAJIT LAHIRI ... ..	899
On $L^1$ -convergence of certain trigonometric sums by BABU RAM and SURESH KUMARI ... ..	908



On $\alpha$ -quasi convex functions by T. N. SHANMUGAM	915
Edge crack in orthotropic elastic half-plane by J. DE and B. PATRA	923
Flow of a second order fluid due to the rotation of an infinite porous disk near a stationary parallel porous disk by B. B. SINGH and ANIL KUMAR	931
Propagation of alfvén waves in a real magneto-hydrodynamic fluid by M. Y. NASIR and M. ILYAS	944
Operator duals of some sequence spaces by N. RATH	953
On strongly NBD-finite families by P. THANGAVELU	964
Bounded and Fréchet differentials for mappings on linear topological spaces using pseudonorm topology by S. DAYAL and ARCHANA MARWAHA	969
Lie theory of $q$ -Appell function by LAKSHMI VARADARAJAN	977
On distributional Laplace-Hardy $\mathcal{L}F_v$ transformation by B. R. AHIRRAO and S. V. MORE	989
Distributional boundary values in $(W_M^a)$ -spaces of functions holomorphic in tube domains by R. S. PATHAK and A. C. PAUL	1004
The axisymmetric Cauchy-Poisson problem in a stratified liquid by LOKENATH DEBNATH and UMA B. GUHA	1022
On transient development of waves at an interface between two fluids by M. S. FALTAS	1032
On the relation of lattice repleteness and $C$ -real compactness by GEORGE BACHMAN and PANAGIOTIS D. STRATIGOS	1043
A note on Swan modules by ANUPAM SRIVASTAV	1067
Fixed point theorems for multivalued mappings by T. L. HICKS	1077
An abstract fixed point theorem for multi-valued mappings by S. K. SAMANTA	1080
On linear independence of sequences in conjugate Banach spaces by P. K. JAIN, S. K. KAUSHIK and D. P. SINHA	1083
Asymptotic behaviour of solutions of functional differential equations by E. THANDAPANI and K. BALACHANDRAN	1096

	<i>Page</i>
On Bishop, Silov and antialgebraic decompositions <i>by</i> H. S. MEHTA and R. D. MEHTA ... ..	1107
On topological projective planes-III <i>by</i> S. ARUMUGAM ... ..	1115
Invariant submanifolds in a conformal $K$ -contact Riemannian manifold <i>by</i> B. RAVI and C. S. BAGEWADI ... ..	1119
Associated Weber integral transforms of arbitrary orders <i>by</i> C. NASIM ...	1126
On the generating functions and partial sums of the Fourier series <i>by</i> PREM CHANDRA ... ..	1139
Unsteady motion of a semi-infinite conducting liquid by a suddenly applied velocity on its surface <i>by</i> D. C. SANYAL and S. K. SAMANTA ... ..	1146
On hydromagnetic turbulent shear flow <i>by</i> D. C. SANYAL and S. K. SAMANTA ... ..	1151
On the stability of compressible swirling flows <i>by</i> V. THULASI and M. SUBBIAH ... ..	1159
Pseudolinearity and efficiency via Dini derivative <i>by</i> SHASHI AGGARWAL and DAVINDER BHATIA ... ..	1173
On the existence of unity in Lehmer's $\psi$ -product ring <i>by</i> V. SITARAMAIAH	1184
Iterative methods of solutions for linear and quasi linear complementarity problems <i>by</i> R. N. MUKHERJEE and H. L. VERMA ... ..	1191
On some new discrete inequalities in two independent variables <i>by</i> B. G. PACHPATTE ... ..	1197
Periodic boundary value problems for an infinite system of nonlinear second order differential equations <i>by</i> K. NARSIMHA REDDY ... ..	1213
On the sets of generalized hypergeometric functions and the Regge, Bargmann-Shelepin arrays for the $3-J$ and the $G-J$ coefficients <i>by</i> K. SRINIVASA RAO and V. RAJESWARI ... ..	1230
Eigen value approach to linear micropolar elasticity <i>by</i> R. K. MAHALANABIS and J. MANNA ... ..	1237
Mellin transform of the gravity effect of A2-D horizontal circular cylinder with variable density <i>by</i> L. ANAND BABU, N. L. MOHAN, N. SUNDARARAJAN and S. V. SESOAGIRI RAO ... ..	1251
<i>Contents and Index</i> ... ..	<i>i</i>

# INDEX

	Page		Page
A.A. Ammar : <i>see</i> H. El Owaidy		Alfvén waves in a real magneto- hydrodynamic fluid ...	944
A.A. Ammar : <i>see</i> H. El-Owaidy		Algebraic polynomial : On the average number of Crossings of an algebraic polynomial ...	1
Abdussattar : Circular orbits of charged test particles in Riessner- Nordstrom field ...	641	Algebras : A note on normed near- algebras ...	433
A.B. Kumar : Thermoelastic waves from suddenly punched hole in stretched elastic plate ...	181	Almost continuous functions : On almost continuous functions ...	571
A. Carbone : A Fixed point theorem for generalized contraction map ...	543	$\alpha$ -Hausdorff subsets : On $\alpha$ -Hausdorff subsets, almost closed mappings and almost upper semicontinuous decomposition ...	334
Accretive mappings : Fixed point iterations for nonlinear Hammer- stein equation involving non- expansive and accretive mappings ...	129	Amar Nath Gill : Selection proce- dures for hazard rates based on two samples statistics ...	773
A. Chakrabarti : On some dual integral equations involving Bessel function of order one ...	483	Amiyadeb Mukherjee : Nonsta- tionary law of heat conduction in classical thermoelastic solid ...	834
A.C. Paul : <i>see</i> R. S Pathak		A.M.N. Yogi : <i>see</i> V. P. Singh	
A.D. Alawneh : Ellipsoidal inclusions in an elastic medium ...	840	Analytic functions : Extreme points of some families of analytic functions related to univalent functions ...	820
A.G.A.E. Abdel-Gawad : On auto- morphisms of free groups ...	40	Anil Kumar : <i>see</i> B. B. Singh	
A.K. Agarwal : <i>see</i> I. C. Gupta		Antialgebraic decompositions : On Bishop, Šilov and antialgebraic decompositions ...	1107
A.K. Agarwal : An optimal pro- gramme for augmentation of capacities of depots and shipment of buses from depots to starting points of routes ...	111	Anupam Srivastav : A note on Swan modules ...	1067
A.K. Mishra : Extreme points of some families of analytic func- tions related to univalent func- tions ...	820	Appell functions : Lie theory of $q$ -Appell functions ...	977
A.K. Tiwari : Banach space valued distributional Mellin transform and form invariant linear filtering ...	493	Arbitrary ring : Copure-injective modules ...	250
Alfvén waves : Propagation of		Archana Marwaha : <i>see</i> S. Dayal	
		Artinian ring : Artinian (Noetherian) part of a Goldie ring ...	49

	Page		Page
Asymptotic behaviour : Asymptotic behaviour of solutions of functional differential equations	1096	tions of orthonormal series	150
Average Number : Average number of crossings of an algebraic polynomial	1	B.E. Rhoades : <i>see</i> A. Carbone	
Axisymmetric Cauchy-Poisson problem : The Axisymmetric Cauchy-Poisson problem in a stratified liquid	1022	B.K. Patel : <i>see</i> G. Das	
Aydin Tiryaki : Periodic solutions of a certain fourth order differential equation	235	B.M. Pandeya : Variants of Hopfcity in modules	786
Babu Ram : On $L^1$ -convergence of certain trigonometric sums	908	B.N. Mandal : A cylindrical wave-maker in liquid of finite depth with an inertial surface	505
Banach limits : Functional limits	812	B.N. Prasad : A transformation of the Finsler metric by an $h$ -vector	455
Banach space : Banach space valued distributional Mellin transform and form invariant linear filtering	493	B.N. Prasad : Cubic transformations of Finsler spaces and $n$ fundamental forms of their hyper-surfaces	242
Maximal elements in Banach spaces	690	B.N. Prasad : On $h\nu$ -recurrent Finsler connection	790
Functional limits	812	B.N. Prasad : <i>see</i> J. N. Singh	
On linear independence of sequences in conjugate Banach spaces	1083	Boolean algebra : $O$ -distributive posets	804
Bayes approach : Bayes approach to prediction in samples from Gamma population when outliers are present	858	Boundary value problems : Periodic boundary value problems for an infinite system of nonlinear second order differential equations	1213
B.B. Sinha : On almost unified contact Finsler structures and connections	887	Boundary values : Distributional boundary values in $(W_M^n)'$	
B.B. Singh : Flow of a second order fluid due to the rotation of an infinite porous disk near a stationary parallel porous disk	931	spaces of functions Holomorphic in tube domains	1004
B.G. Pachpatte : On a class of nonlinear higher order differential equations	121	Bounded sequences : On $F$ absolutely translative summability methods	893
B.G. Pachpatte : On some new discrete inequalities in two independent variables	1191	B.R. Ahirrao : On distributional Laplace-Hardy $\mathcal{L}f_\nu$ transformation	989
B.E. Rhoades : Matrix transformations of orthonormal series		B. Ravi : Invariant submanifolds in a conformal $\alpha$ -contact Riemannian manifold	1119
		B. Patra : <i>see</i> J. De	
		Byung Hak Kim : Remarks on submanifolds of codimensional euclidean space	466
		Canonical subgroup : A note on	



	Page		Page
Swan modules ...	1067	uniform stability of a system of differential equation with Complex coefficients ...	307
Carleman boundary value problem: A generalized Carleman boundary value problem for multiply connected domains ...	829	Composite integral functions: Growth of composite integral functions ...	899
C. Dhanpal: <i>see</i> R. S. Agarwal		Compressible swirling flows: On the stability of compressible swirling flows ...	1159
C. E. Chidume: Fixed points iterations for non-linear Hammersstein equation involving non-expansive and accretive mappings ...	129	Compressional wave: Scattering of a Compressional wave at the corner of a quarter space ...	386
Chandreshwar Thakur: Hodograph transformation in constantly inclined two-phase MFD Flows	728	Conducting fluid: Flow of a conducting fluid between two coaxial rotating porous cylinders bounded by a permeable bed ...	526
Charged test particles: Circular orbits of charged test particles in Riessner-Nordstrom field ...	641	Conformal mappings: Differential subordination and conformal mappings ...	560
Circular orbits: Circular orbits of a charged test particles in Riessner-Nordstrom field ...	641	Conjecture of Snapper: Cyclotomic number and conjecture of Snapper ...	99
C. Jayaram: Noetherian Regular Rings ...	554	Conjugate Banach spaces: On linear independence of sequence in conjugate Banach spaces ...	1083
Classical thermoelastic solid: Non-stationary law of heat conduction in classical thermoelastic solid ...	854	Continuous functions: On almost continuous functions ...	571
Class of polynomials: A finite integral involving a general class of polynomials and the multivariable $H$ -function ...	604	Contraction map: A fixed point theorem for generalized contraction map ...	543
Closed mappings: On $\alpha$ -Hausdorff subsets, almost closed mappings and almost upper semicontinuous decomposition ...	334	Contraction mapping: Sequences of mappings converging to a contraction mapping ...	549
C. Nasim: Associated Weber integral transforms of arbitrary orders ...	1126	Control function: On Controllability of nonlinear systems with distributed delays in the control On the existence for a class of optimal control problems ...	213 761
Commodity spaces: Maximal elements in Banach spaces ...	690	Controllability: On Controllability of nonlinear systems with distributed delays in the control ...	213
Commutative ring: On complete integral closure of $G$ -Domain ...	884	Convex functions: On $\alpha$ -Quasi Convex functions ...	915
Complete integral closure: On complete integral closure of $G$ -domain ...	884	Convex univalent polynomials	
Complex coefficients: On the			



	Page		Page
Convex univalent polynomials	341	semi-infinite conducting liquid by a suddenly applied velocity on its surface	1146
Convexity structure : An abstract fixed point theorem for multi-value mappings	1080	D.C. Sanyal : On hydromagnetic turbulent shear flow	1151
C. Orhan : On $F$ -absolutely translatable summability methods	893	Development of waves : On transient development of waves at an interface between two fluids	1032
Injective modules of use : Copure injective modules	250	Differential equations : On a class of nonlinear higher order differential equations	121
Coset diagrams : Coset diagrams for an action of the extended modular group on the projective line over a finite field	747	Mathematical model of population interactions with dispersal : stability of two habitats with a predator	205
Cosymplectic manifold : Hyper-surfaces with $(f, g, u, v, x)$ -structure of an affinely cosymplectic manifold	799	Periodic solutions of a certain fourth order differential equation	235
Couette flow : Commencement of Couette flow in Oldroyd liquid with heat sources	267	On the uniform stability of a system of differential equations with complex coefficients	307
C. Pakkiam : On the Graphoidal covering number of a graph	330	Asymptotic behaviour of solutions of functional differential equations	1096
C.P. Parvathy : Effect of thermal diffusion on thermohaline inter-leaving in a porous medium due to horizontal gradients	716	Periodic boundary value problems for an infinite system of nonlinear second order differential equation	1213
$C$ -real Compactness : On the relation of lattice repleteness and $C$ -real compactness	1043	Differentiable functions : The Hankel-Clifford transformation on certain space of ultradistributions	583
C.S. Bagewadi : <i>see</i> B. Ravi		Differential rotation : Roche harmonics for Stellar models distorted by differential rotation	284
Cubic transformations : Cubic transformations of Finsler spaces and $n$ fundamental forms of their hypersurfaces	242	Differential systems : Some results on stability of differential systems with impulsive permutations	446
Cyclotomic numbers : Cyclotomic numbers and a conjecture of snapper	99	Digraph reconstruction : On digraph reconstruction	782
Cylindrical wave : A cylindrical wave-maker in liquid of finite depth with an inertial surface	505	Dini derivatives : Pseudolinearity and efficiency via Dini derivatives	1173
Davinder Bhatia : <i>see</i> Shashi Aggarwal			
D.C. Sanyal : Unsteady motion of a			

	Page		Page
Directed-divergence : On a monotonicity property of measures of directed-divergence ...	851	Ellipsoidal inclusions : Ellipsoidal inclusions in an elastic medium ...	840
Discrete inequalities : On some new discrete inequalities in two independent variables ...	1197	Endl-type : On the Endl-type generalization of certain summability methods ...	698
Discrete Pfaffian equations : Gronwall, Bihari and Langenhop type inequalities for discrete Pfaffian equation ...	665	E. Thandapani : Gronwall, Bihari and Largenhop type inequalities for discrete Pfaffian equation ...	665
Distributional boundary values : distributional boundary values in $(W_M^a)'$ -spaces of functions holomorphic in tube domains ...	1004	E. Thandapani : Asymptotic behaviour of solutions of functional differential equation ...	1096
Distributive posets : $O$ -distributive posets ...	804	Euclidean space : Remarks on submanifolds of codimension 2 of an even-dimensional Euclidean space ...	466
Dhruwa Narain : Hypersurfaces with $(f, g, u, v, \lambda)$ -structure of an affinely cosymplectic manifold ...	799	Euler totient function : On the forms of $n$ for which $\varphi(n) \mid n-1$ ...	871
D.P. Sinha : see P.K. Jain		Existence of unity : On the existence of unity in Lehmer's $\psi$ -product ring ...	1184
D. Somasundaram : Inclusion theorems on matrix transformations of some sequence spaces over non-Archimedian fields IV ...	260	Extreme points : Extreme points of some families of analytic functions related to univalent functions ...	820
D. Yadaiah : On temperature-rate dependent thermoelastic longitudinal vibrations of an infinite circular cylinders ...	395	Factor spaces : On Bishop, Silov <sup>v</sup> and antialgebraic decompositions ...	1107
Edge crack : Edge crack in orthotropic elastic half-plane ...	923	F. Cammaroto : Almost irresolute functions ...	472
Eigen value approach : Eigen value approach to linear micropolar elasticity ...	1237	F.D. Zaman : Torsional vibration of a random elastic cylinder ...	521
Elastic cylinder : Torsional vibration of a random elastic cylinder ...	521	Finite field : Coset diagrams for an action of the extended modular group on the projective line over a finite field ...	747
Elastic media : Random Rayleigh waves in a non-homogeneous elastic media ...	646	Finite integral : A finite integral involving a general class of polynomials and the multivariable $H$ -function ...	604
Elastic medium : Ellipsoidal inclusions in an elastic medium ...	840	Finslerian extension : A Finslesian Extension of the gravitational field-II ...	229

	Page		Page
Finsler connection : On $h\nu$ -recurrent Finsler connection ...	790	Functional limits : Functional limits ...	812
Finsler metric : Transformation of the Finsler metric by an $h$ -vector ...	455	Galois field : The extended modular group acting on the projective line over a Galois field ...	755
Finsler spaces : Cubic transformations of Finsler spaces and $n$ fundamental forms of their hypersurfaces ...	242	Gamma population : Bayes approach to prediction in samples from Gamma population when outliers are present ...	858
Finsler structures : On almost unified contact Finsler structures and connections ...	887	Gargi Chakraborty : On the existence for a class of optimal control problems ...	761
Fixed point iterations : Fixed point iterations for nonlinear Hammerstein equation involving nonexpansive and accretive mappings ...	129	G.C. Dash : Commencement of Couette flow in Oldroyd liquid with heat sources ...	267
Fixed point theorem : A fixed point theorem for generalised contraction map ...	543	G. Das : Modified means ...	347
Fixed point theorems for multivalued mappings ...	1077	G. Das : Lacunary distribution of sequences ...	64
An abstract fixed point theorem for multivalued mappings ...	1080	G. Das : Functional limits ...	812
Flow behind weak : Flow behind weak and strong shock waves in water ...	194	Generalized functions : On distributional Laplace-Hardy $\mathcal{L}_f$ transformation ...	989
Fracture : Stresses in prestressed dry sandy soil due to normal moving load leading to instability and fracture ...	165	Generating functions : On the generating functions and partial sums of the Fourier series ...	1139
Fréchet differentials : Bounded and Fréchet differentials for mappings on linear topological spaces using pseudonorm topology ...	969	Generalized convexity : Generalized convexity in multi-objective programming ...	10
Free groups : On automorphisms of Free groups ...	40	George Bachman : On the relation of lattice repleteness and $C$ -real compactness ...	1043
Fourier series : On the generating functions and partial sums of the Fourier series ...	1139	G. Ghous : <i>see</i> F.D. Zaman	
Functional differential equations : Asymptotic behaviour of solutions of functional differential equations ...	1096	Ghanshyam Mehta : Maximal elements in Banach spaces ...	690
		G.K. Pradhan : Thermal stability of a fluid layer in a variable gravitational field ...	736
		Gobind P. Mehta : <i>see</i> Amar Nath Gill	
		Goldie near-rings : Goldie theorem analogue for Goldie near-rings ...	141
		Goldie near-rings : Radical Goldie near-rings ...	439



	Page		Page
Goldie ring : Artinian (Neotherian)		and antialgebraic decomposition	1107
part of a Goldie ring	49	Hazard rates : Selection procedures	
Goldie theorem : Goldie theorem		for hazard rates based on two-	
analogue for Goldie near-rings	141	sample statistics	773
G.P. Tripathi : <i>see</i> J.N. Kapur		Heat conduction : Nonstationary	
Graph : On automorphisms of free		law of heat conduction in classi-	
groups	40	cal thermoelastic solid	834
On the Graphoidal covering num-		Heat convection equation : Note on	
ber of a graph	330	minmax principle for heat con-	
On digraphs reconstruction	782	vection equation	189
Graphoidal cover : On the gra-		Heat sources : Commencement of	
phoidal covering number of a		couette flow in Oldroyd liquid	
Graph	330	with heat sources	267
Gravitational field : A Finslerian		Heat transfer : Numerical solution	
extension of the gravitational		of unsteady flow and heat trans-	
field-II	229	fer in a micropolar fluid past a	
Thermal stability of a fluid layer		porous flat plate	513
in a variable gravitational field	736	H. El-Owaidy : Stability in Mam-	
Gravity effect : Mellin transform of		miliary compartmental Systems	104
the gravity effect of a 2-D hori-		H. El-Owaidy : Mathematical model	
zontal circular cylinder with		of population interactions with	
variable density	1251	dispersal : Stability of two	
Green's function : On some dual		habitats with a predator	205
integral equations involving		H.L. Verma : <i>see</i> R.N. Mukherjee	
Bessel function of order one	483	Hodograph transformation : Hodo-	
Green-Lindsay's : On Rayleigh		graph transformation in const-	
waves in Green-Lindsay's model		antly inclined two phase MFD	
of generalized thermolastic		flows	728
media	276	Hopfian modules : Variants of	
G S. Lingappaiah : Bayes approach		Hopficity in modules	786
to prediction in samples from		Horizontal gradients : Effect of	
gamma population when outliers		thermal diffusion on thermo-	
are present	858	haline interleaving in a porous	
		medium due to horizontal gra-	
Habitats : Mathematical model of		dients	716
population interactions with		H.S. Mehta : On Bishop, Silov and	
dispersal : Stability of two		antialgebraic decompositions	1107
habitats with a predator	205	Hydromagnetic turbulent shear	
Hankel-Clifford transformation :		flow : On hydromagnetic turbu-	
The Hankel-Clifford transform-		lent shear flow	1151
ation on certain spaces of ultra-		Hypergeometric functions : On the	
distributions	583	sets of generalized hypergeo-	
		metric functions and the Regge,	
Hausdorff space : On Bishop, Silov			

	Page		Page
Bargmann-Sheletín arrays for the 3—J and the G—5 coefficients ...	1220	Injective hull : The injective hull of a module with FGD ...	874
Hypersurfaces : Cubic transformations of Finsler spaces and $n$ fundamental forms of their hypersurfaces ...	242	Instability : Stresses in pre-stressed dry sandy soil due to normal moving load leading to instability and fracture ...	165
Hypersurfaces with $(f, g, u, v, \lambda)$ —structure of an affinely cosymplectic manifold ...	799	Integral closure : On complete integral closure of $G$ -Domain ...	884
I.C. Gupta : On a quaternion submanifolds of Co-dimension-2 ...	566	Integral equations : On a class of nonlinear higher order differential equations ...	121
Ilija Kovacević : On $\alpha$ -hausdorff subsets, almost closed mappings and almost upper semicontinuous decomposition ...	334	On some dual integral equations involving Bessel function of order one ...	483
Impulsive perturbations : Some results on stability of differential systems with impulsive perturbations ...	446	Integral functions : Growth of composite integral functions ...	899
Independence of sequences : On linear independence of sequences in conjugate Banach spaces ...	1083	Integral transforms : Associated Weber integral transforms of arbitrary orders ...	1126
Independent variables : On some new discrete inequalities in two independent variables ...	1197	Invariant linear filtering : Banach space valued distributional Mellin transform and firm invariant linear filtering ...	493
Indrajit Lahiri : Growth of composite integral functions ...	899	Invariant submanifolds : Invariant submanifolds in a conformal $K$ -contact Riemannian manifold ...	1119
Indrani Gupta : <i>see</i> M.N. Vartak		Irresolute functions : Almost irresolute functions ...	472
Inequalities : Gronwall, Bihari and Largenhop type inequalities for discrete Pfaffian equation ...	665	J. De : Edge crack in orthotropic elastic half-plane ...	923
Inertial surface : A cylindrical wave-maker in liquid of finite depth with an inertial surface ...	505	Jerry U. Onwuatu : On Controllability of nonlinear systems with distributed delays in the control ...	213
Infinite Circular cylinder : On temperature rate dependent thermoelastic longitudinal vibrations of an infinite circular cylinder ...	395	J.J. Betancor : The Hankel-Clifford transformation on certain spaces of ultradistributions ...	583
Infinite porous : Flow of a second order fluid due to the rotation of an infinite porous disk near a stationary parallel porous disk ...	931	J. Manna : <i>see</i> R.K. Mahalanabis	
		J.N. Kapur : On a monotonicity property of measures of directed-divergence ...	851
		Kalpna : Some results on almost	



	Page		Page
semiin-variant submanifold of an <i>Sp</i> -Sasakian manifold ...	681	of the gravity effect of A 2-D horizontal circular cylinder with variable density ...	1251
K. Balachandran : <i>see</i> E. Thanda- pani		Lacunary distributions : Lacunary distributions of sequences ...	64
K.C. Chowdhury : Artinian (Noeth- erian) Part of a Goldie Ring ...	49	Lakshmi Varadarajan : Lie theory of $q$ -Appel function ...	977
K.C. Chowdhury : Goldie theorem analogue for Goldie near-rings...	141	Lalji Srivastava : <i>see</i> B.N. Prasad	
K.C. Chowdhury : Radical Goldie near-rings ...	439	Lalji Srivastava : <i>see</i> B.N. Prasad	
K.C. Gupta : A finite integral in- volving a general class of poly- nomials and the multivariable $H$ -function ...	604	Laplace-Hardy transformation : On distributional Laplace - Hardy $\mathcal{LF}_v$ transformations ...	989
K. Farahmand : On the average number of an algebraic poly- nomial ...	1	Laplace transform : A cylindrical wave-maker in liquid of finite depth with an inertial surface ...	505
K. Jagadeeswara Pillai : Flow of a conducting fluid between two coaxial rotating porous cylinders bounded by a permeable bed ...	526	Lattice repleteness : On the relation of Lattice repleteness and $C$ -real compactness ...	1043
K.K. Dewan : <i>see</i> M. Bidkham		Layered half space : Quasi-static response of a layered half-space to surface loads ...	621
K.L. Dutta : Random Rayleigh waves in non - homogeneous elastic media ...	646	Libration points : Linear stability and the resonance for the trian- gular libration points for the doubly photogravitational elliptic restricted problem of three bodies ...	403
K. Narasimha Reddy : Periodic boundary value problems for an infinite system of nonlinear second order differential equa- tions ...	1213	Lie theory : Lie theory of $q$ -Appell functions ...	977
K. Raja Gopal Rao : <i>see</i> K. Yugandhar.		Linear complementarity problems : Iterative methods of solutions for linear and quasilinear comple- mentarity problems ...	1191
Krishna Kundu : <i>see</i> : B.N. Mandal		Linear micropolar elasticity : Eigen value approach to linear micro- polar elasticity ...	1223
K. Srinivasa Rao : On the sets of generalized hypergeometric func- tions and the Regge, Bargmann- Shelepin arrays for the 3— $J$ and $G$ — $J$ coefficients ...	1220	Linear stability : Linear stability and the resonance for the triangular libration points for the doubly photogravitational elliptic restric- ted problem of three bodies ...	403
K. Yugandhar : A note on primary decomposition in Noetherian near-rings ...	671	Linear topological spaces : Bounded and Fréchet differentials for mappings on linear topological	
K. Yugandhar : <i>see</i> T. Srinivas			
L. Anand Babu : Mellin transform			

	<i>Page</i>		<i>Page</i>
spaces using pseudonorm topology ...	969	valued distributional Mellin transform and form invariant linear filtering ...	493
Loading functions : Edge crack in orthotropic elastic half-plane. ...	923	Mellin transform of the gravity effect of A 2-D horizontal circular cylinder with variable density ...	125
Lokenath Debnath : The axisymmetric Cauchy-Poisson problem in a stratified liquid ...	1022	Metric space : Sequences of mapping converting to a contraction mapping ...	549
Magnetohydrodynamic fluid : Propagation of Alfvén waves in a real magnetohydrodynamic fluid ...	944	MFD flows : Hodograph transformation in constantly inclined two-phase MFD flows ...	728
M.A. Gopalan : Note on Minmax principle for heat convection equation ...	189	M. GEL-Sheikh : A generalized Carleman boundary value problem for multiply connected domains ...	829
Mamillary compartmental systems : Stability in mamillary compartmental systems ...	104	MHD flows : Comments on steady place MHD flows with constant speed along each streamline ...	423
Mamoru Nunokawa : On the multivalent functions ...	577	Micropolar fluid : Numerical solution of unsteady flow and heat transfer in a micropolar fluid past a porous flat plate ...	513
Mass distribution : Note on the small vibration of beams with varying Young's modulus carrying a concentrated mass distribution ...	75	M. Ilyas : <i>see</i> M.Y. Nasir	
Matrix transformations : Matrix transformations : of orthonormal series ...	150	Minmax principle : Note on minmax principle for heat convection equation ...	189
Matrix transformations in some sequence spaces ...	707	M.N. Vartak : Generalized convexity in multi-objective Programming ...	10
Maximal elements : Maximal elements in Banach spaces ...	690	Modified means : Modified means ...	347
M.A. Sarigol : <i>see</i> C. Orhan		Module with FGD : The injective Hull of a module with FGD ...	874
M. Bidkham : Bounds for the zeros of polynomial ...	768	Modular group : Coset diagrams for an action of the extended modular group on the projective line over a finite field ...	747
M. Chakraborty : <i>see</i> S. Dey		The Extended modular group acting on the projective line over a Galois field ...	755
M.D. Jahagirdar : Transient forced and free convection flow past an infinite vertical plate ...	171	Modules : Variants of Hopficity in modules ...	786
Mean-square : On a error term involving the Totient function ...	537	Monotonicity property : On a	
Measures : On a monotonicity property of measures of directed-divergence ...	851		
Mellin transform : Banach space			

	Page		Page
monotonicity property of measures of directed divergence ...	851	NBD finite families : On strongly NBD-finite families ...	964
M. Rangamma : <i>see</i> V. Siva Rama Prasad		N.C. Dawn : On Rayleigh waves in in Green-Lindsay's model of generalized thermoelastic media	276
M.R. Parameswaran : On the Endl-type generalization of Certain summability methods ...	698	N.C. Mondal : Free torsional vibration of a nonhomogeneous semi-infinite solid circular cylinder ...	89
M.S. Faltas : On the transient development of waves at an interface between two fluids ...	1032	Near-rings : A note on primary decomposition in Noetherian near-rings ...	671
M.S. Kasi : Convex univalent polynomials ...	341	Newtonian fluids : Flow of a second order fluid due to the rotation of an infinite porous disk near a stationary parallel porous disk	931
M. Subbiah : <i>see</i> V. Thulasi		N.L. Mohan : <i>see</i> L. Anand Babu	
M. Syam Babu : <i>see</i> K. Jagadeeswara		N.M. Bujurke : A note on the squeeze film lubrication with non-Newtonian fluid ...	632
Multiply connected domains : A Generalized Carleman boundary value problem for multiply connected domains ...	829	Noetherian near-rings : A note on primary decomposition in noetherian near-rings ...	671
Multi-objective : Generalized convexity in multi-objective programming ...	10	Noetherian regular rings ...	554
Multivalent functions : On the multivalent functions ...	577	Noetherian ring : Artinian (Noetherian) part of a Goldie ring ...	49
Multivalued mappings : Fixed point theorems for multivalued mappings ...	1077	Non-Archimedean fields : Inclusion theorems on matrix transformations of some sequence spaces over non-Archimedean fields IV	260
An abstract fixed point theorem for multi-valued mappings ...	1080	Non-homogeneous : Random Rayleigh waves in non-homogeneous elastic media ...	646
Multivariable $H$ function : A finite integral involving a general class of polynomials and the multivariable $H$ -function ...	604	Nonlinear control system : On controllabilities of nonlinear systems with distributed delays in the control ...	213
M.Y. Nasir : Propagation of Alfvén waves in a real magnetohydrodynamic fluid ...	944	Nonlinear differential equation : An oscillation criterion for second order nonlinear differential equation ...	297
Narinder Mohan : Scattering of a compressional wave at the corner of a quarter space ...	386	Nonlinear Hammerstein equation : Fixed point iterations for nonlinear Hammerstein equation	
Nat Ram Garg : Quasi-static response of a layered half-space to surface loads ...	621		



	Page		Page
involving nonexpansive and accretive mappings ...	129	Orthotropic elastic half-plane : Edge crack in orthotropic elastic half-plane ...	923
Non-Newtonian fluid : A note on the squeeze film lubrication with non-Newtonian fluid ...	632	Orthonormal series : Matrix transformations of orthonormal series ...	150
Non Stationry : Non-stationary law of heat conduction in classical thermoelastic solid ..	834	Oscillation criterion : An oscillation criterion for second order non-linear differential equation ...	297
Nörlund matrices : Modified means	347	Pammy Manchanda : <i>see</i> Surjit Singh	
Normed near-algebras : A note on normed near-algebras ...	433	Panagiotis D. Strati Gos : <i>see</i> George Bachman	
Normal moving load : Stresses in pre-stressed dry sandy soil due to normal moving load leading to instability and fracture ...	165	P.C. Samal : <i>see</i> G K. Pradhan	
N. Rath : Operator duals of some sequence spaces	953	P. Dheena : A Generalization of strongly regular near-rings ...	58
N. Sundararajan : <i>see</i> L. Anand Babu		Periodic solutions : Periodic solutions of a certain fourth order differential equation ...	235
N.T. Shawagfeh : <i>see</i> A.D. Alawneh		P.K. Chaudhuri : Note on the small vibration of beams with varying Young's modulus carrying a concentrated mass distribution	75
Nurettin Ergun : On strongly rare-continuity	609	P.K. Jain : On linear independence of sequences in conjugate Banach spaces ...	1083
O.A. Elleithy : <i>see</i> H-El-Owaidy		P.K. Kundu : Incoming water waves against a vertical cliff in a two-fluid medium ...	292
Oldroyd liquid : Commencement of Couette flow in Oldroyd liquid with heat sources ...	267	Prabhamani R. Patil : <i>see</i> C.P. Parvathy	
O.P. Chandna : Comments on Steady plane MHD flows with constant speed along each streamline ...	423	Prandtl number : Transient forced and free convection flow past an infinite vertical plate ...	711
Operator duals : Operator duals of some sequence spaces ...	953	Predator : Mathematical model of population interactions with dispersal : Stability of two habitats with a predator ...	205
Optimal Control : On the existence for a class of optimal control problems ...	761	Prem Chandra : On the generating functions and partial sums of Fourier series ...	1139
Optimal programme : An optimal programme for augmentation of capacities of depots and shipment of buses from depots to starting points of routes ...	111	Primary decomposition : A note on primary decomposition in Noetherian near-rings ...	671
Optimal : Selection of optimal site for new depot of specified capacity with two objectives ...	425		

Page	Page
Product ring : On the existence of unity in Lehmer's $\psi$ -product ring ... 1184	Quasi convex functions : On $\alpha$ -quasi convex functions ... 915
Programming : Generalized convexity in multi-objective programming ... 10	Quasi-static response : Quasi static response of a layered half-space to surface loads ... 621
Polynomials : Convex Univalent polynomials ... 341	Quaternion submanifolds : On a quaternion submanifolds of co-dimension-2 ... 566
Bounds for the zeros of polynomials ... 768	Radical Goldie near-rings : Radical Goldie near-rings ... 439
Porous cylinders : Flow of a conducting fluid between two coaxial rotating porous cylinders bounded by a permeable bed ... 526	Raghwendra : <i>see</i> R.N. Mukherjee
P. Sahu : <i>see</i> A.K. Mishra	Ram Babu Mishra : <i>see</i> Chandreshwar Thakur
P.S. Deshwal : <i>see</i> Narinder Mohan	Ram Kumar Shukla : <i>see</i> D. Yadiach
Pseudo linearity : Pseudolinearity and efficiency via Dini derivatives ... 1173	Random algebraic polynomial : On the real roots of a random algebraic polynomial ... 655
Pseudonorm topology : Bounded and Fréchet differentials for mappings on linear topological spaces using pseudonorm topology ... 969	Random Rayleigh waves : Random Rayleigh waves in non-homogeneous elastic media ... 646
P.S. Hiremath : <i>see</i> N.M. Bujurke	Rare-continuity : On strongly rare-continuity ... 609
P. Thangavelu : On strongly NBD-finite families ... 964	Rayleigh waves : On Rayleigh waves in Green-Lindsay's model of generalized thermoelastic media ... 276
$p$ -valently convexity : On the multivalent functions ... 577	R.D. Mehta : <i>see</i> H.S. Mehta
$p$ -valently starlikeness : On the multivalent functions ... 577	Real roots : On the real roots of a random algebraic polynomial ... 655
Qaiser Mustaq : Coset diagrams for an action of the extended modular group on the projective line over a finite field ... 747	Recurrent Finsler connection : On $h$ -recurrent Finsler connection ... 790
Quarter space : Scattering of a compressional wave at the corner of a quarter space ... 386	Regular near-rings : A Generalization of strongly regular near-rings ... 58
Q. Mushtaq : The extended modular group acting on the projective line over a Galois field ... 715	Regular rings : Noetherian regular rings ... 554
	Rehana Quraishi : <i>see</i> Abdussatar
	Riemannian hypersurfaces : Transformation of the Finsler metric by an $h$ -vector ... 455
	Riemannian manifold : Invariant submanifolds in a conformal $k$ -contact Riemannian manifold ... 1119



	Page		Page
Riessner-Nordstrom field : Circular orbits of charged test particles in Riessner-Nordstrom field ...	641	Satoshi Ikeda : A Finslerian extensions of the gravitational field-II	229
Rivlin-Ericksen tensors : A note on the squeeze film lubrication with non-Newtonian fluid ...	632	Satya Prakash : Selection of optimal site for new depot of specified capacity with two objectives ...	425
R.K. Mahalanbis : Eigen value approach to linear micropolar elasticity ...	1237	S. Bagh : On the real roots of a random algebraic polynomial ...	655
R.K. Yadav : <i>see</i> B.B. Sinha		S. Bhavanari : The injective hull of a module with FGD ...	874
R.M. Lahurikar : <i>see</i> M.D. Jahagirdar		S. Biswal : <i>see</i> G.C. Dash	
R.N. Mukherjee : Some results on stability of differential systems with impulsive perturbations ...	446	Scalar curvature : On Wagner spaces of $W_p$ -scalar curvature	136
R.N. Mukherjee : Iterative methods of solutions for linear and quasilinear complementarity problems ...	119	S. Dayal : Bounded and Fréchet differentials for mappings on linear topological spaces using pseudonorm topology ...	969
Roche harmonics : Roche harmonics for Stellar models distorted by differential rotation ...	284	S. Dey : Stresses in prestressed dry sandy soil due to normal moving load leading to instability and fracture ...	165
R.S. Agarwal : Numerical solution of unsteady flow and heat transfer in a micropolar fluid past a porous flat plate ...	513	Sequence spaces : Operator duals of some sequence spaces ...	953
R.S. Pathak : Distributional boundary values in $(W_M^a)'$ - spaces of a functions holomorphic in tube domains ...	1004	Seismology : Scattering of a compressional wave at the corner of a quarter space ...	386
S.A. Paramhans : <i>see</i> B.M. Pandeya		Semicontinuous decomposition : On $\alpha$ -Hausdorff subsets, almost closed mappings and almost upper semicontinuous decomposition ...	334
S.A. Katre : Cyclotomic numbers and a conjecture of Snapper ...	99	Semi-infinite conducting liquid : Unsteady motion of a semi-infinite conducting liquid by a suddenly applied velocity on its surface ...	1146
S. Arumugam : On topological projective plans-III ...	1115	Semi-invariant submanifold : Some results on almost semi-invariant submanifolds of an $SP$ -Sasakian manifold ...	681
S. Arumugam : <i>see</i> C. Pakkiam		Semi-open sets : Almost irresolute functions ...	472
Sarvajit Singh : <i>see</i> Nat Ram Garg		Sequence spaces : Matrix transformations in some sequence spaces	707
Sasakian manifold : Some results on almost semi-invariant submanifold of an $SP$ -sasakian manifold	681		
S. Asgar : <i>see</i> F.D. Zaman			

	Page		Page
Inclusion theorems on matrix transformations of some sequence spaces over non-Archimedean fields IV	260	Solid circular cylinder: Free torsional vibration of a nonhomogeneous semi-infinite solid circular cylinder	89
Sequences: Lacunay distribution of sequences	64	Solutions: Iterative methods of solutions for linear and quasilinear complementarity problems	1191
Sequences of mapping converging to a contraction mapping	549	S.P. Koirala: <i>see</i> B.M. Pandeya	
Sets: On the sets of generalized hypergeometric functions and the Regge, Bargmann-Shelepin arrays for the 3-J and the 6-J Coefficients	1220	Specified capacity: Selection of optimal site for new depot of specified capacity with two objectives	425
S.G. Bhavi: <i>see</i> N.M. Bujurke		S. Ponnusamy: <i>see</i> V. Karunkaran	
Shashi Aggarwal: Pseudolinearity and efficiency via Dini derivatives	1173	S.P. Singh: <i>see</i> A. Carbone	
Shear flow: On hydromagnetic turbulent shear flow	1151	Squeeze film lubrication; A note on the squeeze film lubrication with non-Newtonian fluid	632
Shipment of buses: An optimal programme for augmentation of capacities of depots and shipment of buses from depots to starting points of routes	111	S. Rama Chandran: On digraph reconstruction	782
Shock waves: Flow behind weak and strong shock waves in water	194	S.R. Grace: An oscillation criterion for second order nonlinear differential equation	297
S.K. Chakraborty: <i>see</i> K.L. Dutta		Starting points of routes: An optimal programme for augmentation of capacities of depots and shipment of buses from depots to starting points of routes	111
Skin-friction: Transient forced and free convection flow past an infinite vertical plate	711	Stratified liquid: The axisymmetric Cauchy-Poisson problem in a stratified liquid	1022
S.K. Kaushik: <i>see</i> P.K. Jain		Stretched elastic plate: Thermoelastic waves from suddenly punched hole in stretched elastic plate	181
S.K. Samanta: <i>see</i> D.C. Sanyal		Stresses: Stresses in pre-stressed dry sandy soil due to normal moving load leading to instability and fracture	165
S.K. Samanta: An abstract fixed point theorem for multivalued mappings	1080	Strongly NBD-finite families: On strongly NBD-finite families	964
S.K. Samanta: <i>see</i> D.C. Sanyal		Strongly rare-continuity: On strongly rare-continuity	609
S.K. Singh: On Wagner spaces of $W_p$ -scalar curvature	136	Stability: Stability in mammillary	
S.L. Dhingra: <i>see</i> A.K. Agrawal			
S.M. Aggarwal: <i>see</i> K.C. Gupta			
Small vibration: Note on the small vibration of beams with varying Young's Modulus carrying a concentrated mass distribution	75		

	Page		Page
compartmental systems ...	104	Takashi Noiri: On almost continuous functions ...	571
Mathematical model of population interactions with dispersal:		Takashi Noiri: <i>see</i> F. Cammaroto	
Stability of two habitats with a predator ...	205	Thermal diffusion: Effect of thermal diffusion on thermohaline interleaving in a porous medium due to horizontal gradients ...	716
Some results on stability of differential systems with impulsive perturbations ...	446	Thermal stability: Thermal stability of a fluid layer in a variable gravitational field ...	736
On the stability of compressible swirling flows ...	1159	Thermoelastic media: On Rayleigh waves in Green Lindsay's model of Generalized thermoelastic media ...	276
Stellar models: Roche harmonics for stellar models distorted by differentiable rotation ...	284	Thermoelasticity: On temperature-rate dependent thermoelastic longitudinal vibrations of an infinite circular cylinder ...	395
Streamline: Comments on steady plane MHD flows with constant speed along each streamline ...	423	Thermo-elastic: Thermo-elastic waves from suddenly punched hole in stretched elastic plate ...	181
Submanifolds of codimension-2: On a quaternion submanifolds of Co-dimension-2 ...	566	Thermohaline interleaving: Effect of thermal diffusion on thermohaline interleaving in a porous medium due to horizontal gradients ...	716
Remarks on submanifolds of Codimension 2 of an even dimensional euclidean space ...	466	Three bodies: Linear stability and the resonance for the triangular libration points for the doubly photogravitational elliptic restricted problem of three bodies ...	403
Subrata Data: <i>see</i> P.K. Chandhuri		Topological projective planes: On topological projective planes-III ...	1115
Sudarshan Nanda: Matrix transformations in some sequence spaces ...	707	Topological measure: On the relation of lattice repleteness and C-T.L. Hicks: Fixed point theorems for multivalued mappings ...	1077
Summability: On the Endl-type generalization of certain summability methods ...	698	real compactness ...	1043
Suresh Kumari: <i>see</i> Babu Ram		Torsional oscillation: Free torsional vibration of a nonhomogeneous semi-infinite solid circular cylinder ...	89
Surface loads: Quasi-static response of a layered half-space to surface loads ...	621		
Surjit Singh: On complete integral closure of $G$ -Domain ...	884		
S.V.K. Varma: <i>see</i> K. Jagadeeswara			
S.V. More: <i>see</i> B.R. Ahirrao			
S. Nanda: <i>see</i> G. Das			
S.V. Seshagiri Rao: <i>see</i> L. Anand Babu			
Swan modules: A note on swan modules ...	1067		



	Page		Page
Torsional : Free torsional vibration of a nonhomogeneous semi-infinite solid circular cylinder ...	89	Debnath	...
Torsional vibration : Torsional vibration of a random elastic cylinder ...	521	Unified structure : On almost unified contact Finsler structures and connection ...	887
Totient function : On an error term involving the totient function ...	537	Uniform stability : On the uniform stability of a system of differential equations with complex coefficients ...	307
T.N. Shanmugam : On $\alpha$ -quasi convex functions ...	915	Unsteady flow : Numerical solution of unsteady flow and heat transfer in a micropolar fluid past a porous flat plate ...	513
Transient development : On transient development of waves at an interface between two fluids ...	1032	Unsteady motion : Unsteady motion of a semi infinite conducting liquid by a suddenly applied velocity on its surface ...	1146
Transient : Transient forced and free convection flow past an infinite vertical plate ...	711	V.A. Hiremath : Copure-injective modules ...	250
Transformations : Inclusion theorems on matrix transformations of some sequence spaces over a non-Archimedean fields IV ...	260	Vandana Gupta : <i>see</i> V.P. Singh	
Transformation of the Finsler metric by an $h$ -vector ...	455	Variable gravitational field : Thermal stability of fluid layer in a variable gravitational field ...	736
Translative summability : On $F$ -absolutely translative summability methods ...	893	Variants of Hopfcity : Variants of Hopfcity in modules ...	786
Trigonometric sums : On $\alpha^1$ -convergence of certain trigonometric sums ...	908	V.B. Dhamke : <i>see</i> Y.S. Pawar	
T. Srinivas : A Note on normed near-algebras ...	433	Vertical cliff : Incoming water waves against a vertical cliff in a two-fluid medium ...	272
T. Srinivas : <i>see</i> K. Yugandhar		Vibrations : On temperature rate dependent thermoelastic longitudinal vibrations of an infinite circular cylinder ...	395
Thodoy Vidalis : Sequences of mappings converging to a contraction mapping ...	549	Vivek Saini : <i>see</i> Satya Prakash	
Two-sample statistics : Selection procedures for hazard rates based on two sample statistics ...	773	V. Kumar : Linear stability and the resonance for the triangular libration points for the doubly photogravitational elliptic restricted problem of three bodies ...	403
Two-fluid medium : Incoming water waves against a vertical cliff in a two fluid medium ...	292	V. Karunakaran : Differential subordination and conformal mappings ...	560
Uma B. Guha : <i>see</i> Lokenath		V.L. Mannapalli : <i>see</i> C. Jayaram	

	<i>Page</i>		<i>Page</i>
V.K. Tripathy : <i>see</i> G.K. Pradhan		Weber integral transforms : Associated Weber integral transforms of arbitrary orders	... 1126
V.P. Singh : Flow behind weak and strong shock waves in water	... 194	Werner George Nowak : On an error term involving the Totient function	... 537
V.P. Singh : Roche harmonics for stellar models distorted by differential rotation	... 284	Young's Modulus : Note on the small vibration of beams with varying Young's modules carrying a concentrated mass distribution	75
V. Rajeswari : <i>see</i> K. Srinivasa		Y.S. Pawar : <i>O</i> -distributive posets	804
V. Sitaramaiah : On the existence of unity in Lehmer's $\psi$ -product ring	... 1184	Zeros : Bounds for the zeros of polynomials	... 768
V. Siva Rama Prasad : On the forms of $n$ for which $Q(n)/n - 1$	... 871	Zerosymmetric : A generalization of strongly regular near rings	... 58
V. Thulasi : On the stability of compressible swirling flows	... 1159	Z. Zahreddine : On the uniform stability of a system of differential equations with complex coefficients	... 307
Wagner spaces : On Wagner spaces $W_p$ -scalar curvature	... 136		
Water waves : Incoming water waves against a vertical cliff in a two fluid medium	... 292		



# Eratta

"Edge crack in Orthotropic Elastic Half Plane" by J. DE AND B. PATRA  
—printed in Vol 20, No. 9 (1989), pp. 923-930.




FIG.  $\phi(t)/P_0$  versus  $t$  for various loading function.



## SUGGESTIONS TO CONTRIBUTORS

The INDIAN JOURNAL OF PURE AND APPLIED MATHEMATICS is devoted primarily to original research in pure and applied mathematics.

Manuscripts should be typewritten, double-spaced with sufficient margins (including abstracts, references, etc.) on one side of durable white paper. The initial page should contain the title followed by author's name and full mailing address. The text should include only as much as is needed to provide a background for the particular material covered. Manuscripts should be submitted in triplicate.

The author should provide a short abstract, in triplicate, not exceeding 250 words, summarizing the highlights of the principal findings covered in the paper and the scope of research.

References should be cited in the text by the arabic numbers in superior. List of references should be arranged in the arabic numbers, author's name, abbreviation of Journal, Volume number (Year) page number, as in the sample citation given below :

### *For Periodicals*

1. R. H. Fox, *Fund. Math.* 34 (1947) 278.

### *For Books*

2. H. Rund, *The Differential Geometry of Finsler Spaces*, Springer-Verlag, Berlin, (1973) p. 283.

Abbreviations for the titles of the periodicals should, in general, conform to the *World List of Scientific Periodicals*.

All mathematical expressions should be written clearly including the distinction between capital and small letters. Clear distinction between upper and lower cases of c, p, k, z, s, should be made while writing the expression in hand. Also distinguish between the letters such as 'Oh' and 'zero'; l(el) and 1 (one); v, V and  $\nu$  (Greek nu);  $r$  and  $\gamma$  (Greek gamma);  $\chi$ , X and  $\times$  (Greek chi); k, K and  $\kappa$  (Greek kappa); Greek letter lambda ( $\Lambda$ ) and symbol for vector product ( $\wedge$ ); Greek letter epsilon ( $\epsilon$ ) and symbol for 'is an element of' ( $\in$ ). The equation numbers are to be placed at the right-hand side of the page. The name of the Greek letter or symbol should be written in the margin the first time it is used. Superscripts and subscripts should be simple and should be placed accurately.

Line drawings should be made with India ink on white drawing paper or tracing paper. Letterings should be clear and large. Photographic prints should be glossy with strong contrast. All illustrations must be numbered consecutively in the order in which they are mentioned in the text and should be referred to as Fig. or Figs. Legends to figures should be typed on a separate sheet and attached at the end of the manuscript.

Tables should be typed separately from the text and placed at the end of the manuscript. Table headings should be short but clearly descriptive.

Proofs should be corrected immediately on receipt and returned to the Editor. If a large number of corrections are made in the proof, the author should pay towards composition charges. In case, the author desires to withdraw his paper, he should pay towards the composition charges, if the same is already done.

For each paper, the authors will receive 50 reprints free of cost. Order for extra reprints should be sent with corrected page proofs.

Manuscripts, in triplicate, should be submitted to the Editor of Publications, *Indian Journal of Pure and Applied Mathematics*, Indian National Science Academy, Bahadur Shah Zafar Marg, New Delhi 110002 (India).



## INDIAN JOURNAL OF PURE AND APPLIED MATHEMATICS

No. 12

December 1989

Volume 20

## CONTENTS

	<i>Page</i>
Pseudolinearity and efficiency via Dini derivatives by SHASHI AGGARWAL and DAVINDER BHATIA ... ..	1173
On the existence of unity in Lehmer's $\psi$ -product ring by V. SITARAMAIAH	1184
Iterative methods of solutions for linear and quasi linear complementarity problems by R. N. MUKHERJEE and H. L. VERMA ... ..	1191
On some new discrete inequalities in two independent variables by B. G. PACHPATTE ... ..	1197
Periodic boundary value problems for an infinite system of nonlinear second order differential equations by K. NARSIMHA REDDY ... ..	1213
On the sets of generalized hypergeometric functions and the Regge, Bargmann-Shelepin arrays for the $3-J$ and the $6-J$ coefficients by K. SRINIVASA RAO and V. RAJESWARI ... ..	1230
Eigen value approach to linear micropolar elasticity by R. K. MAHALANABIS and J. MANNA ... ..	1237
Mellin transform of the gravity effect of A2-D horizontal circular cylinder with variable density by L. ANAND BABU, N. L. MOHAN, N. SUNDARARAJAN and S. V. SESHAGIRI RAO ... ..	1251
Contents and Index ... ..	i
Errata	